Draft version October 19, 2026
Typeset using LaTeX **default** style in AASTeX7.0.1

# Three Sub-Neptune Transit Candidates Around Kepler Bonus Background Stars

Aayush Gupta[1]

[1] *Independent researcher*

## ABSTRACT

I report three transiting sub-Neptune candidates from a pre-registered search of 6,465 dwarf stars that fell on Kepler target-pixel files but were never Kepler targets, using Kepler Bonus photometry. KIC 6699887 ($P = 15.114\,\mathrm{d}$, $2.8\,R_\oplus$), KIC 3831969 ($15.313\,\mathrm{d}$, $3.5\,R_\oplus$) and KIC 11138423 ($4.894\,\mathrm{d}$, $2.8\,R_\oplus$) show transits that localize on the target in difference images, no odd–even or secondary signal, no ephemeris match to any known Kepler object, and transit-derived stellar densities consistent with Gaia DR3. Using TRICERATOPS with companion constraints from Gaia astrometry and imaging, their false-positive probabilities are 0.6–0.7% and their nearby-star false-positive probabilities are zero. KIC 6699887 also shows a second on-target signal at $47.69\,\mathrm{d}$. High-resolution imaging is needed before these can be considered validated.

## 1. SAMPLE AND SEARCH

Kepler Bonus (Martínez-Palomera et al. 2023) provides point-spread-function photometry for 606,900 sources on Kepler target-pixel files, most of them background stars. Bienias & Szabó (2025) reported hot-Jupiter candidates among them, and the Kepler Bonus team is running its own search. Before searching I froze a list of 6,465 sources. Of 425,219 background sources, 118,648 have PSF fraction $> 0.5$, 64,075 also crowding metric $> 0.7$, 27,378 also $\geq 8$ quarters and 13,696 also $G < 17.5$; 9,513 of these were never Kepler targets, 9,335 remain after excluding the 1,800 sources of an earlier public search of this dataset (Mehta 2026), and 6,465 are dwarfs by Gaia DR3 (Gaia Collaboration et al. 2023). I normalized each quarter, detrended with a 1-day biweight (Hippke et al. 2019), and ran box least squares (Kovács et al. 2002) over 0.5–40 d with thresholds registered in advance. The same search ran on every light curve inverted ($2 - f$): 448 stars passed on real flux and 97 on inverted flux. Two positive controls (KIC 5689121 and Kepler-138) passed before any pool result was read.

## 2. VETTING

Of these, 22 passed a second stage (masked re-detrending, per-transit depths, gap proximity, variability, duration). Some second-stage criteria and both pixel tests were defined after an early look at the results; every revision is dated in the public log, and I treat everything after the first screen as post hoc. For localization I differenced in- and out-of-transit images from the originating target-pixel files and fitted point sources at the Gaia positions of the target and of every neighbor within two pixels, plus a scaled stamp image, per quarter and per Kepler season (Figure 1). The test places KIC 5689121 on target and two known off-target signals on other stars. Five of the 22 localize on target. Applied to the 97 inverted signals, the second stage passes six and the pixel test places none on target. The search is not calibrated for completeness, and I infer no occurrence rate. Each signal then received an independent re-derivation with separate code and an adversarial review, and four of the five survived; for the three candidates the re-derivation reproduces the period within 10 s and the depth within 7%. Across all reviewed signals, rejections included post-downlink thermal ramps, single-cadence artifacts, a saturated neighbor, and an eclipsing binary $5.6'$ away on the same detector channel. No candidate matches the period (within 0.1%, harmonics 1:4–4:1) and epoch of any KOI, DR25 TCE (Thompson et al. 2018) or Kepler eclipsing binary (Kirk et al. 2016) in the Kepler field, and the brighter star on each stamp shows no dip ($\geq 29\sigma$).

## 3. FALSE-POSITIVE PROBABILITY

I ran TRICERATOPS (Giacalone et al. 2021) in Kepler mode, eight runs of $10^6$ draws each, removing a neighbor only when its own light curve excludes the eclipse it would need. With default priors the false-positive probabilities (FPP) are 2.4, 2.7 and 3.6%, almost entirely a planet orbiting an unresolved bound companion. The hosts have Gaia

RUWE 1.06–1.09. MOLUSC (Wood et al. 2021), using RUWE and Gaia imaging and a multiplicity fraction of 0.46, reduces the companion prior from 26–29% to 15–19%, giving FPP $= 0.63 \pm 1.03$, $0.71 \pm 0.46$ and $0.70 \pm 0.21$% (mean and scatter over runs) and zero nearby-star FPP. A fourth survivor, KIC 3963519, has FPP $\approx 39$% because of an unmodeled $G = 18$ neighbor and is not included. Transit fits (Kreidberg 2015; Foreman-Mackey et al. 2013) give stellar densities within $0.4\sigma$ of Gaia FLAME values (Creevey et al. 2023).

## 4. CANDIDATES

Ephemerides are $\mathrm{BJD_{TDB}}$; radii use Gaia FLAME stellar radii; S/N is the depth over the scatter of same-duration boxes at random phases, and SDE is the BLS peak significance (in parentheses, the strongest peak on inverted flux).

- KIC 6699887 (Gaia DR3 2077554781499350144, TIC 270864056; $T_{\rm eff} = 4,542$ K, 859 pc): $P = 15.11366^{+0.00006}_{-0.00005}$ d, $T_0 = 2455693.1849^{+0.0012}_{-0.0013}$, depth 1,023 ppm, $T_{14} = 3.78^{+0.18}_{-0.13}$ h, $R_p = 2.76^{+0.17}_{-0.14}\,R_\oplus$, S/N 20.0 over 86 transits, SDE 89 (14). A second signal ($P = 47.6901$ d, 951 ppm, 28 transits) passes the same vetting and the seasonal pixel test.
- KIC 3831969 (Gaia DR3 2100154070473374976, TIC 120425498; 5,275 K, 1,969 pc): $P = 15.31347^{+0.00013}_{-0.00012}$ d, $T_0 = 2455697.0076^{+0.0030}_{-0.0029}$, depth 997 ppm, $T_{14} = 4.04^{+0.26}_{-0.22}$ h, $R_p = 3.47^{+0.38}_{-0.32}\,R_\oplus$, S/N 23.8 over 86 transits, SDE 46 (13).
- KIC 11138423 (Gaia DR3 2129669944725625344, TIC 424864042; 4,258 K, 894 pc): $P = 4.89443^{+0.00002}_{-0.00002}$ d, $T_0 = 2455726.7654^{+0.0020}_{-0.0018}$, depth 1,316 ppm, $T_{14} = 1.91^{+0.17}_{-0.14}$ h, $R_p = 2.83^{+0.27}_{-0.24}\,R_\oplus$, S/N 15.2 over 253 transits, SDE 33 (15).

None appears in public candidate lists as of 2026 October 17. Statistical limits on companions within $0.5''$ are the main remaining uncertainty; speckle or adaptive-optics imaging of these $G = 16$–17 hosts would test them. TESS observed each host in 9–10 sectors, but at $T = 15.5$–16.5 even an undiluted, ideal-aperture estimate (Stassun et al. 2018) gives a combined transit S/N of at most 1.0, so TESS can neither confirm nor exclude these signals. I used Kepler Bonus light curves ( doi:10.17909/7jbr-w430) and Kepler long-cadence data, Q0–Q17 ( doi:10.17909/T9488N), from MAST, and the KOI cumulative, DR25 TCE, Kepler stellar and Planetary Systems tables of the NASA Exoplanet Archive ( doi:10.26133/NEA4, doi:10.26133/NEA15, doi:10.26133/NEA6, doi:10.26133/NEA12), accessed 2026 October 17. Code, light curves, run outputs and evidence packs are archived at Zenodo, doi:10.5281/zenodo.23442225.

## ACKNOWLEDGMENTS

This paper includes data collected by the Kepler mission and obtained from the MAST data archive at the Space Telescope Science Institute (STScI); funding for the Kepler mission is provided by the NASA Science Mission Directorate, and STScI is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5–26555. It uses data from the European Space Agency mission Gaia, processed by the Gaia Data Processing and Analysis Consortium, and has made use of the NASA Exoplanet Archive, operated by the California Institute of Technology under contract with NASA under the Exoplanet Exploration Program. I used Claude (Anthropic) to help write analysis code and draft the text; I reviewed all results and the text and take responsibility for them.

*Facilities:* Kepler, Gaia, Exoplanet Archive

## REFERENCES

Bienias, D., & Szabó, R. 2025, A&A, 694, A209

Creevey, O. L., et al. 2023, A&A, 674, A26

Foreman-Mackey, D., et al. 2013, PASP, 125, 306

Gaia Collaboration, Vallenari, A., et al. 2023, A&A, 674, A1

Giacalone, S., et al. 2021, AJ, 161, 24

Hippke, M., et al. 2019, AJ, 158, 143

Kirk, B., et al. 2016, AJ, 151, 68

Kovács, G., Zucker, S., & Mazeh, T. 2002, A&A, 391, 369

Kreidberg, L. 2015, PASP, 127, 1161

Mehta, K. 2026, Planethunt: archival evidence for the KIC 5689121 transit candidate, Zenodo, doi:10.5281/zenodo.23325261





**Figure 1.** Top: phase-folded Kepler Bonus photometry (gray), binned (black) with the transit model (red). Bottom: missing flux during transit (S/N) for one Kepler season; red circle, target Gaia position; blue crosses, other Gaia sources. $\Delta\chi^2$ values compare a source at the target with no signal and with the best alternative, summed over four seasons; the season shown is labeled.

Martínez-Palomera, J., Hedges, C., & Dotson, J. 2023, AJ, 166, 265

Stassun, K. G., et al. 2018, AJ, 156, 102

Thompson, S. E., et al. 2018, ApJS, 235, 38

Wood, M. L., Mann, A. W., & Kraus, A. L. 2021, AJ, 162, 128