



# Tomato

256 pesticide parameters screened across 4 samples — unwashed, mineral water, baking soda, and Estoqi Wash — from the same produce batch.



| Parameters                                          | <i>No Wash</i> | <i>Mineral Water</i> | <i>Baking Soda</i> | <i>Estoqi Wash</i>    |
|-----------------------------------------------------|----------------|----------------------|--------------------|-----------------------|
| <b>Profenofos</b><br>Organophosphate                | 7.936<br>mg/kg | 7.801<br>mg/kg       | 7.633<br>mg/kg     | <b>4.896</b><br>mg/kg |
| <b>Lambda Cyhalothrin</b><br>Pyrethroid insecticide | 2.786<br>mg/kg | 2.732<br>mg/kg       | 1.359<br>mg/kg     | <b>1.299</b><br>mg/kg |
| <b>Propargite</b><br>Acaricide                      | 0.900<br>mg/kg | 0.890<br>mg/kg       | 0.883<br>mg/kg     | <b>0.656</b><br>mg/kg |
| <b>Paclobutrazole</b><br>Fungicide · PGR            | 0.379<br>mg/kg | 0.373<br>mg/kg       | 0.328<br>mg/kg     | <b>0.234</b><br>mg/kg |
| <b>Tebuconazole</b><br>Triazole fungicide           | 0.067<br>mg/kg | 0.059<br>mg/kg       | 0.058<br>mg/kg     | <b>Eliminated</b>     |
| <b>Tolfenpyrad</b><br>Pyrazole insecticide          | 0.036<br>mg/kg | 0.033<br>mg/kg       | 0.028<br>mg/kg     | <b>Eliminated</b>     |
| <b>Thiamethoxam</b><br>Neonicotinoid                | 0.019<br>mg/kg | 0.014<br>mg/kg       | 0.013<br>mg/kg     | <b>Eliminated</b>     |
| <b>Imidacloprid</b><br>Neonicotinoid                | 0.033<br>mg/kg | 0.025<br>mg/kg       | 0.025<br>mg/kg     | <b>Eliminated</b>     |
| <b>Monocrotophos</b><br>Organophosphate             | 0.021<br>mg/kg | 0.012<br>mg/kg       | 0.012<br>mg/kg     | <b>Eliminated</b>     |
| <b>Copper compounds</b><br>Fungicide · inorganic    | 1.090<br>mg/kg | 0.910<br>mg/kg       | 0.700<br>mg/kg     | <b>0.680</b><br>mg/kg |

Estoqi outperforms baking soda — the most widely recommended home alternative. Systemic pesticides absorbed into plant tissue are not addressed by surface washing.

