



Tomato

A controlled 6-day observation study on tomatoes sourced from a leading quick commerce partner - comparing Estoqi-treated samples against untreated controls across firmness, texture, colour progression, and Brix stability.



12 Sample groups

3 Ripeness stages

2 storage conditions

+3 days

Marketable window extended Estoqi-treated tomatoes remained sellable up to Day 5-6 vs Day 3 cutoff for controls.

50%

Better firmness retention
Pulp pressure readings on Day 6 showed 35-50% better retention in Estoqi-treated samples.

Zero

Additional inputs required
No cold chain upgrades. No packaging changes. No chemical coatings. The Estoqi wash alone, pH 11.5 ionised water, within 3-4 minutes.

Observed Outcomes

Summary across all 12 sample groups
Days 1-6

Parameter

Without Estoqi

Estoqi

Skin Cracking

Moderate-High

Minimal

Shrivelling

Visible by Day 3-4

Delayed past Day 5

Colour Ripening

Faster, uneven

Gradual, uniform

Brix Stability

Early spike → drop

Controlled, stable





The difference a wash makes, visible over 10 days.

NORMAL WASH

Marketable
quality lost at
Day 10



Day 0



Day 3



Day 7



Day 10

ESTOQI WASH

Still viable at
Day 10



Day 0



Day 3



Day 7



Day 10

