AI1103-Assignment 9

Name: Ayush Jha Roll Number: CS20BTECH11006

Download all python codes from

https://github.com/ayushjha2612/AI11003/tree/main /Assignment9/Codes

and latex-tikz codes from

https://github.com/ayushjha2612/AI11003/tree/main /Assignment9

CSIR UGC NET EXAM (June 2017) Q. 103

Let $c \in \mathbb{R}$ be a constant. Let X, Y be random variables with joint probability density function

$$f(x, y) = \begin{cases} cxy & \text{, if } 0 < x < y < 1\\ 0 & \text{, otherwise} \end{cases}$$

Which of the following statements are correct?

- 1) $c = \frac{1}{8}$ 2) c = 8
- 3) X and Y are independent
- 4) Pr(X = Y) = 0

Answer

Option (2) c = 8 and option (4) Pr(X = Y) = 0.

SOLUTION

Solving all options:

- 2) X and Y are two random variables with joint pdf

$$f(x,y) = \begin{cases} cxy & \text{, if } 0 < x < y < 1 \\ 0 & \text{, otherwise} \end{cases}$$
 (0.0.1)

The marginal probability density functions are as follows:

$$f_X(x) = \int_{-\infty}^{\infty} f(x, y) \, dy \qquad (0.0.2)$$

$$f_Y(y) = \int_{-\infty}^{\infty} f(x, y) dx$$
 (0.0.3)

Calculating $f_X(x)$

$$f_X(x) = \int_{-\infty}^{\infty} f(x, y) \, dy \qquad (0.0.4)$$

$$= \int_{x}^{1} cxy \, dy \tag{0.0.5}$$

$$= cx \left(\frac{y^2}{2}\right)\Big|_{x}^{1} \tag{0.0.6}$$

$$=cx\left(\frac{1-x^2}{2}\right)\tag{0.0.7}$$

$$f_X(x) = \begin{cases} cx\left(\frac{1-x^2}{2}\right) & \text{, if } 0 < x < 1\\ 0 & \text{, otherwise} \end{cases}$$
 (0.0.8)

Calculating $f_{Y}(y)$

$$f_{y}(y) = \int_{-\infty}^{\infty} f(x, y) dx$$
 (0.0.9)

$$= \int_0^y cxy \, dx \tag{0.0.10}$$

$$= cy \left(\frac{x^2}{2}\right) \Big|_{0}^{y} \tag{0.0.11}$$

$$=\frac{cy^3}{2}$$
 (0.0.12)

$$f_Y(y) = \begin{cases} \frac{cy^3}{2} & \text{, if } 0 < y < 1\\ 0 & \text{, otherwise} \end{cases}$$
 (0.0.13)

Now by using property of pdf and equation (0.0.13) we have,

$$\int_{-\infty}^{\infty} f_Y(y) \, dy = 1 \tag{0.0.14}$$

$$\int_0^1 c \frac{y^3}{2} = 1 \tag{0.0.15}$$

$$\frac{c}{0} = 1$$
 (0.0.16)

$$c = 8$$
 (0.0.17)

Therefore option (2) is correct.

3) The pdf of X is

$$f_X(x) = \begin{cases} 4x(1-x^2) & \text{, if } 0 < x < 1\\ 0 & \text{, otherwise} \end{cases}$$
(0.0.18)

The pdf of Y is

$$f_Y(y) = \begin{cases} 4y^3 & \text{, if } 0 < y < 1\\ 0 & \text{, otherwise} \end{cases}$$
 (0.0.19)

To check whether X and Y are independent, we calculate $f_X(x) \times f_Y(y)$. From (0.0.18) and (0.0.19)

$$f_X(x) \times f_Y(y) = \begin{cases} 16xy^3 (1 - x^2) & \text{, if } 0 < x, y < 1 \\ 0 & \text{, otherwise} \end{cases}$$

$$(0.0.20)$$

$$\neq f(x, y) \tag{0.0.21}$$

Since f(x, y) and $f_X(x) \times f_Y(y)$ are different, random variables X and Y are not independent. Therefore option (3) is not correct.

4) To find Pr(X = Y), we use the conditional pdf,

$$f_{X|Y}(x|y) = \frac{f_{XY}(x,y)}{f_{Y}(y)}$$

$$= \lim_{\Delta x, \Delta y \to 0} \Pr(x < X \le x + \Delta x | y < Y \le y + \Delta y)$$
(0.0.23)

$$\approx \frac{\Pr(X = x, Y = y)}{\Pr(X = x)}$$
 (0.0.24)

Similarly we have,

$$f_{Y|X}(y|x) = \frac{f_{XY}(x,y)}{f_{X}(x)}$$

$$\approx \frac{\Pr(X = x, Y = y)}{\Pr(Y = y)}$$
(0.0.25)

$$f_{XY}(x, y) = \begin{cases} 8xy & \text{, if } 0 < x < y < 1\\ 0 & \text{, otherwise} \end{cases}$$
 (0.0.27)

Calculation of conditional PDF and using

(0.0.27),

$$f_{X|Y}(x|y) = \frac{f_{XY}(x,y)}{f_Y(y)}$$
 (0.0.28)

$$= \frac{0}{f_Y(y)} \text{ for } x = y \tag{0.0.29}$$

$$= 0 \text{ for } x = y$$
 (0.0.30)

Similarly we have,

$$f_{Y|X}(y|x) = 0 \text{ for } x = y$$
 (0.0.31)

Therefore for x = ywe have,

$$f_{X|Y}(x|y) = 0 (0.0.33)$$

$$f_{Y|X}(y|x) = 0 (0.0.34)$$

$$\implies \frac{\Pr(X = x, Y = y)}{\Pr(Y = y)} = 0 \qquad (0.0.35)$$

$$Pr(X = x, Y = y) = 0 \text{ for } x = y$$
 (0.0.36)

$$\implies \Pr(X = Y) = 0 \tag{0.0.37}$$

Therefore option (2) and (4) are correct.

The marginal PDF of X and Y are shown at figure 4 and figure 4.




Fig. 4: The marginal PDF of X




Fig. 4: The marginal PDF of Y