


PARAMETERS III: BIOLOGY AND NATURAL HISTORY

Steven M. Goodreau, Ph.D.
Samuel M. Jenness, Ph.D.
Eli S. Rosenberg, Ph.D.



SCHOOL OF PUBLIC HEALTH
UNIVERSITY AT ALBANY State University of New York

Outline for this session

- Circumcision
- CCR5- Δ 32 mutation
- HIV disease progression: VL and AIDS/mortality among ART naïve
- How these and behaviors come together for per-act tx probability
- Implementing in the model

Circumcision




Circumcision: concept

- Biomedical intervention that reduces HIV acquisition risk
- Trials in sub-Saharan African heterosexual males showed ~60% efficacy
- Among MSM, marginal efficacy is lower because of varied sex roles
 - No benefit to receptive partners
 - In insertive partners...
 - Similar efficacy assumed as to heterosexuals
 - Metanalysis found insufficient evidence of efficacy in insertive MSM (Millett et al, JAMA 2008)

Circumcision: assessment and model

47. Is your penis circumcised (cut) or uncircumcised (uncut)?
Please select one choice.

1 ☐ Circumcised (cut) 0 ☐ Uncircumcised (uncut) 9 ☐ Don't know

Circumcised (cut) **Uncircumcised (uncut)** **Don't know**

circumcised **circumcised.**

- 90% of both B and W men were circumcised
- Node attribute at entry. Conveys 60% protection to insertive HIV- men

CCR5- Δ 32


CCR5- Δ 32: Concept

- Protein receptor on CD4+ T-cells that facilitates viral entry
- Δ 32 mutation results in defective protein
 - Homozygote (2 copies) confers full immunity
 - Heterozygote: 70% reduced susceptibility in 1 MSM study
- Distribution of Δ 32 allele different by race
 - Whites: 3.4% homozygote, 17.6% heterozygote
 - Blacks: ~0% homozygote 2.1% heterozygote

HIV disease progression: ART naïve

HIV disease progression

- HIV infection natural history, per CD4 and VL



HIV disease progression in model

- Model currently tracks VL and AIDS status (but not CD4)
- Progression to death in about 10 years:

	Duration	Start VL value	End VL value
'Up' part of curve to peak of acute virema	45 days	0	$6.886 \log_{10}$ copies per mL
'Down' part to set-point VL	45 days	$6.886 \log_{10}$ copies per mL	$4.5 \log_{10}$ copies per mL
Chronic infection	3550 days	$4.5 \log_{10}$ copies per mL	$4.5 \log_{10}$ copies per mL
AIDS to death	728 days	$4.5 \log_{10}$ copies per mL	$7 \log_{10}$ copies per mL (Death)

- Segments of VL curve are linear
- Deterministic timing. VL values too?

**Bringing biology and behaviors
together for HIV transmission**

Bringing biology and behaviors together for HIV transmission

- Probability of HIV tx within act of serodiscordant dyad drawn as Bernoulli event with p determined by:

Predictor	Partner	Parameters	References
Sexual role (insertive or receptive)	HIV-	Receptive: 0.008938 base probability when HIV+ partner has 4.5 log ₁₀ viral load copies per mL	Vittinghoff ²⁹
		Insertive: 0.003379 base probability when HIV+ partner has 4.5 log ₁₀ viral load copies per mL	Vittinghoff ²⁹
HIV viral load (VL)	HIV+	Multiplier of $2.45^{(VL - 4.5)}$, where VL is expressed as log ₁₀ copies per mL	Wilson ³⁰
Acute stage	HIV+	Multiplier of 6	Leynaert, ¹⁸ Bellan ³¹
CCR5 status	HIV-	Δ32 homozygote: multiplier of 0	Marmor ¹⁴
		heterozygote: multiplier of 0.3	Marmor ¹⁴
Condom use	Both	Multiplier of 0.25	Varghese, ³² Weller ³³ , Smith ³⁴
Circumcision status	HIV-, insertive	Multiplier of 0.40	Gray ¹³

Implementing in the model

Circumcision, CCR5

- param.mard:

```
disc.inst.B.prob = 0.445, disc.inst.W.prob = 0.691, circ.B.prob = 0.874,  
circ.W.prob = 0.918, ccr5.B.prob = c(0, 0.034), ccr5.W.prob = c(0.021,  
0.176),
```

- EpiModelHIV:::setBirthAttr.mard:

```
dat$attr$circ[newIds[newB]] <- rbinom(nBirths.B, 1, dat$param$circ.B.prob)  
dat$attr$circ[newIds[newW]] <- rbinom(nBirths.W, 1, dat$param$circ.W.prob)  
dat$attr$role.class[newIds[newB]] <- sample(c("T" "R" "V")
```

```
ccr5.B.prob <- dat$param$ccr5.B.prob  
ccr5.W.prob <- dat$param$ccr5.W.prob  
dat$attr$ccr5[newIds[newB]] <- sample(c("WW", "DW", "DD"),  
nBirths.B, replace = TRUE, prob = c(1 - sum(ccr5.B.prob),  
ccr5.B.prob[2], ccr5.B.prob[1]))  
dat$attr$ccr5[newIds[newW]] <- sample(c("WW", "DW", "DD"),  
nBirths.W, replace = TRUE, prob = c(1 - sum(ccr5.W.prob),  
ccr5.W.prob[2], ccr5.W.prob[1]))
```


- And stay tuned for transmission....

VL and progression – ART-naive

- param.mard:

```
7, vl.acute.rise.int = 21, vl.acute.peak = 6.886, vl.acute.fall.int = 21,  
vl.set.point = 4.5, vl.aids.onset.int = 520 * 7, vl.aids.int = 52 *  
2 * 7, vl.fatal = 7, vl.full.supp = 1.5, vl.part.supp = 3.5,
```

- update.vl.mard (for viral load)
- progress.mard (for which stage)
- way too much code to show meaningfully. Summary:



- And stay tuned for transmission....

VL and progression – ART

- param.mard:

```
max.time.off.tx.full.int = 520 *  
7, max.time.on.tx.part.int = 52 * 15 * 7, max.time.off.tx.part.int = 520 *  
7, vl.acute.rise.int = 21, vl.acute.peak = 6.886, vl.acute.fall.int = 21,
```

```
2 * 7, vl.fatal = 7, vl.full.supp = 1.5, vl.part.supp = 3.5,  
full.supp.down.slope = 0.25, full.supp.up.slope = 0.25, part.supp.down.slope = 0.25,  
part.supp.up.slope = 0.25, b.B.rate = 0.001/7, b.W.rate = 0.001/7,  
birth.rate = 10, b.method = "fixed", up.tx.rate = 0.0002 *
```

- update.vl.mard: changes to viral load b/c of tx

- summary:

- tx makes VL go down (to different levels depending on whether full or partial suppression)
 - stopping treatment makes VL go up
 - Reinitiating tx makes VL go down again 😊

VL and progression – ART

- progress.mard: tx failure and AIDS initiation occurs when:

```
aids.tx.naive <- which(active == 1 & status == 1 & cum.time.on.tx ==  
  0 & (time.since.inf >= vl.aids.onset) & stage != "D")  
part.tx.score <- (cum.time.off.tx/max.time.off.tx.part) +  
  (cum.time.on.tx/max.time.on.tx.part)  
aids.part.escape <- which(active == 1 & cum.time.on.tx >  
  0 & tt.traj == "YP" & stage == "C" & part.tx.score >=  
  1 & stage != "D")  
aids.off.tx.full.escape <- which(active == 1 & tx.status ==  
  0 & tt.traj == "YF" & cum.time.on.tx > 0 & cum.time.off.tx >=  
  max.time.off.tx.full & stage != "D")  
isAIDS <- c(aids.tx.naive, aids.part.escape, aids.off.tx.full.escape)
```

- And stay tuned for transmission....

Transmission!

- param.mard:

```
birth.age = 18, b.method = "fixed", URAI.prob = 0.0082 *  
1.09, UIAI.prob = 0.0031 * 1.09, acute.rr = 4, circ.rr = 0.4,  
condom.rr = 0.25, disc.outset.main.B.prob = 0.685, disc.outset.main.W
```

- trans.mard:

```
trans.ip.prob <- URAI.prob * 2.45^(ip.v1 - 4.5)  
trans.ip.prob[ip.stage == "AR"] <- trans.ip.prob[ip.stage ==  
  "AR"] * acute.rr  
trans.ip.prob[ip.stage == "AF"] <- trans.ip.prob[ip.stage ==  
  "AF"] * (1 + (acute.rr - 1) * (v1.acute.fall.int - ip.stage.time[ip.stage ==  
  "AF"])/v1.acute.fall.int)  
trans.ip.prob[disc.ip$uai == 0] <- trans.ip.prob[disc.ip$uai ==  
  0] * condom.rr  
trans.ip.prob[ip.ccr5 == "DD"] <- trans.ip.prob[ip.ccr5 ==  
  "DD"] * 0  
trans.ip.prob[ip.ccr5 == "DW"] <- trans.ip.prob[ip.ccr5 ==  
  "DW"] * ccr5.heteroz.rr  
trans.ip.prob[which(ip.prep == 1 & ip.prepc1 == "1")] <- trans.ip.prob[which(ip.prep ==  
  1 & ip.prepc1 == "1")] * (1 - pcc[1])
```

- Etc.