

PARAMETERS IV: CLINICAL CARE AND TREATMENT

Steven M. Goodreau, Ph.D.
Samuel M. Jenness, Ph.D.
Eli S. Rosenberg, Ph.D.



SCHOOL OF PUBLIC HEALTH
UNIVERSITY AT ALBANY State University of New York

Outline for this session

- Care continuum as modeled
 - HIV testing
 - ART initiation (linkage to care with ART)
 - Dynamic engagement in care via ART on/off status
 - Dynamic suppression status
 - On ART, fully suppressed
 - On ART, partially suppressed
 - (Not suppressed because not on ART or undiagnosed)

HIV testing

HIV testing: concept

- HIV testing is critical prevention tool
 - Behavior changes post-diagnosis
 - Necessary for receipt of HIV care (and PrEP in PrEP models)
- Model conceptualized 2 types of testers
 - Never testers – 6.5% of sample didn't have HIV test by age 40
 - Regular testers: memoryless exponential process, per mean time between tests
 - Susie Cassells (UCSB) has R21 extending to more types
- Also incorporated 21-day window period until Ab detection
 - HIV+ positive persons thus test HIV-ne

HIV testing: assessment

62. Have you ever been tested for HIV?

- 1 ☐ Yes If yes, then #63, #64, #65, #66
0 ☐ No

HIVtest_ever

Binary_prefernot.

63. In what month and year did you have your most recent HIV test?

Month:

January	=1
February	=2
March	=3
April	=4
May	=5
June	=6
July	=7
August	=8
September	=9
October	=10
November	=11
December	=12

HIVtest_month

HIVtest_month.

64. Year: HIVtest_year

65. What was the result of your most recent HIV test?

- 0 ☐ Negative
1 ☐ Positive If positive, then hide #67, #70, #71, #72, #73. If positive, then show #68, #69
2 ☐ Indeterminant/Inconclusive
3 ☐ Didn't get the results of my last HIV test

HIVtest_result

HIVtest_result.

HIV testing: model metric

- Mean time since last test: (today – date of last test)
 - BMSM: 301 days
 - WMSM: 315 days
- Assuming testing is memoryless exponential with rate parameter equal to above means, Bernoulli daily (-> weekly) testing probability is the inverse of these.

ART initiation
(linkage to care with ART)

ART linkage/initiation

- Linkage to care defined in many ways, but typically about the duration from diagnosis until first HIV clinical visit
- The model has no direct representation of clinical visits
 - It does have ART, which is a primary purpose of HIV care
 - Linkage to care to data empirically available in some CDC reports
 - But not time to ART initiation. Assumed instant initiation of therapy once in care
- Following HIV diagnosis, MSM initiated treatment:
 - BMSM rate = 0.924/week (10.8 weeks on avg.)
 - WMSM rate = 1.271/week (7.9 weeks on avg.)

Dynamic engagement in care

Dynamic engagement in care: model concept



- Persons on ART can fall out of care
 - Viral load returns to $4.5 \log_{10}$ set-point linearly over 3 months
- Persons can re-engage in care and resume adherence process
- Rates based on cross-sectional CDC care continuum sources
 - Volumes can be said about the veracity of these estimates

Dynamic suppression status | on ART

- Once on ART, may or not be fully adherent
 - Like “care” is modeled as ART on/off status, adherence is modeled in terms of VL suppression levels
 - Fully suppressed = $1.5 \log_{10} = 50$ copies/mL = LLD
 - Partially suppressed = (Not suppressed can occur if on ART or undiagnosed)
 - (unsuppressed occurs if not diagnosed or not on ART)
 - Can cycle between full/partial states
 - Rate parameter back-calculation complicated and used CDC paper on:
 - Viral suppression at last test
 - “Durable” viral suppression at all test for 12 months

Dynamic suppression status | on ART

Parameter	Black MSM	White MSM
Proportion of those initiating ART who achieved full suppression	0.614	0.651
Per-time step probability of falling out of suppression	0.0102	0.0071
Per-time step probability of re-achieving suppression	0.00066	0.00291

Implementing in the model

Testing

- param.mard:

```
function (nwstats, last.neg.test.B.int = 301, mean.test.B.int = 301,  
  last.neg.test.W.int = 315, mean.test.W.int = 315, testing.pattern = "interval",  
  test.window.int = 21, tt.traj.B.prob = c(0.077, 0, 0.356,
```

- test.mard:

```
}  
if (testing.pattern == "interval") {  
  tsinceIntst <- at - dat$attr$last.neg.test  
  tsinceIntst[is.na(tsinceIntst)] <- at - dat$attr$arrival.time[is.na(tsinceIntst)]  
  tst.B <- which(active == 1 & race == "B" & tt.traj !=  
    "NN" & (diag.status == 0 | is.na(diag.status)) &  
    tsinceIntst >= mean.test.B.int)  
  tst.W <- which(active == 1 & race == "W" & tt.traj !=  
    "NN" & (diag.status == 0 | is.na(diag.status)) &  
    tsinceIntst >= mean.test.W.int)  
}  
tst.pos.B <- tst.B[status[tst.B] == 1 & inf.time[tst.B] <=  
  at - twind.int]  
tst.neg.B <- setdiff(tst.B, tst.pos.B)  
tst.pos.W <- tst.W[status[tst.W] == 1 & inf.time[tst.W] <=
```

Treatment initiation, halting, re-initiation

- param.mard:

```
test.window.int = 21, tt.traj.B.prob = c(0.077, 0, 0.356,  
    0.567), tt.traj.W.prob = c(0.052, 0, 0.331, 0.617), tx.init.B.prob = 0.092,  
tx.init.W.prob = 0.127, tx.halt.B.prob = 0.0102, tx.halt.W.prob = 0.0071,  
tx.reinit.B.prob = 0.00066, tx.reinit.W.prob = 0.00291,
```

- tx.mard: who initiates, halts, and re-starts treatment, e.g.:

```
tx.reinit.w.prob <- dat$param$tx.reinit.w.prob  
tx.init.elig.B <- which(active == 1 & race == "B" & status ==  
    1 & tx.status == 0 & diag.status == 1 & tt.traj %in%  
    c("YP", "YF") & cum.time.on.tx == 0 & stage != "D")  
tx.init.B <- tx.init.elig.B[rbinom(length(tx.init.elig.B),  
    1, tx.init.B.prob) == 1]  
tx.init.elig.W <- which(active == 1 & race == "W" & status ==
```