

Questions with Answer Keys

MathonGo

Q1. The value of x for which $\sin(\cot^{-1}(1+x)) = \cos(\tan^{-1}x)$ is

A. $\frac{1}{2}$

B. 1

C. 0

D. $-\frac{1}{2}$

Ans: $-\frac{1}{2}$

Solution: Let, $\cot^{-1}(x+1) = \theta$ and $\tan^{-1}x = \phi$

Now, given equation becomes $\sin \theta = \cos \phi$

$$\Rightarrow \frac{1}{\sqrt{1+(x+1)^2}} = \frac{1}{\sqrt{1+x^2}}$$

$$\Rightarrow (x+1)^2 + 1 = x^2 + 1 \Rightarrow x = -\frac{1}{2}$$

Q2. The value of $\cos\left(\frac{1}{2}\cos^{-1}\left(\cos\left(\sin^{-1}\frac{\sqrt{63}}{8}\right)\right)\right)$ is

A. $\frac{3}{16}$

B. $\frac{3}{8}$

C. $\frac{3}{4}$

D. $\frac{3}{2}$

Ans: $\frac{3}{4}$

Solution:

The given trigonometric ratio

Questions with Answer Keys

$$= \cos\left(\frac{1}{2} \cos^{-1}\left(\cos\left(\cos^{-1}\frac{1}{8}\right)\right)\right)$$

$$= \cos\left(\frac{1}{2} \cos^{-1}\frac{1}{8}\right)$$

$$= \sqrt{\frac{1 + \cos\left(\cos^{-1}\frac{1}{8}\right)}{2}} = \frac{3}{4}$$

Note: One may also proceed by writing the

ratio as $\cos\left(\frac{1}{2} \sin^{-1}\frac{\sqrt{63}}{8}\right)$.

Q3. The value of $\sin^{-1} \sin 17 + \cos^{-1} \cos 10$ is equal to

A. 27

B. -27

C. $17 - 5\pi$

D. $9\pi - 27$

Ans: $9\pi - 27$

Solution: $\sin^{-1} \sin 17 = \sin^{-1} \sin(17 - 5\pi + 5\pi)$

$$= 5\pi - 17$$

$$\cos^{-1}(\cos 10) = \cos^{-1} \cos(10 - 3\pi + 3\pi)$$

$$= \cos^{-1} \cos\{3\pi + (10 - 3\pi)\}$$

$$= \cos^{-1}\{-\cos(10 - 3\pi)\}$$

$$= \pi - \cos^{-1} \cos(10 - 3\pi)$$

$$= \pi - (10 - 3\pi) = 4\pi - 10$$

$$\text{Hence, } \sin^{-1} \sin 17 + \cos^{-1}(\cos 10) = 9\pi - 27$$

Q4. The value of $\sin\left\{\cot^{-1}\left[\cos\left(\cot^{-1}\left(\frac{1}{x}\right)\right)\right]\right\}$ is equal to ($x > 0$)

Questions with Answer Keys

MathonGo

A. $\sqrt{\frac{1+x^2}{2+x^2}}$

B. $\sqrt{\frac{1-x^2}{2+x^2}}$

C. $\sqrt{\frac{1+x^2}{2-x^2}}$

D. $\sqrt{\frac{2+x^2}{1+x^2}}$

Ans: $\sqrt{\frac{1+x^2}{2+x^2}}$

Solution: Let $\cot^{-1}\left(\frac{1}{x}\right) = \alpha \Rightarrow \tan \alpha = x$

So, $\cos \alpha = \frac{1}{\sqrt{1+x^2}}$

$\therefore \sin\left\{\cot^{-1}\left[\cos\left(\tan^{-1} x\right)\right]\right\}$

$= \sin\left(\cot^{-1} \frac{1}{\sqrt{1+x^2}}\right)$

Let $\cot^{-1} \frac{1}{\sqrt{1+x^2}} = \beta \Rightarrow \cot \beta = \frac{1}{\sqrt{1+x^2}}$

$\therefore \sin \beta = \frac{\sqrt{1+x^2}}{\sqrt{2+x^2}}$

$\therefore \sin \beta = \sin\left\{\cot^{-1}\left[\cos\left(\cot^{-1}\left(\frac{1}{x}\right)\right)\right]\right\} = \sqrt{\frac{1+x^2}{2+x^2}}$

Q5. If the value of the expression $\tan\left(\frac{1}{2}\cos^{-1}\frac{2}{\sqrt{5}}\right)$ is in the form of $a + \sqrt{b}$ where $a, b \in \mathbb{Z}$, then the value of $\frac{a+b}{b}$ is

Ans: 0.6

Solution: We know that,

$\tan\left(\frac{x}{2}\right) = \frac{1-\cos x}{\sin x}$

$$\therefore \tan\left(\frac{1}{2}\cos^{-1}\frac{2}{\sqrt{5}}\right) = \frac{1-\cos\left(\cos^{-1}\left(\frac{2}{\sqrt{5}}\right)\right)}{\sin\left(\cos^{-1}\left(\frac{2}{\sqrt{5}}\right)\right)}$$

$$= \frac{1-2/\sqrt{5}}{\sin\left(\sin^{-1}\left(1/\sqrt{5}\right)\right)}$$

Questions with Answer Keys

$$= \frac{1-2/\sqrt{5}}{1/\sqrt{5}} = \sqrt{5} - 2$$

$$\text{Now, } a + \sqrt{b} = -2 + \sqrt{5}$$

$$\Rightarrow a = -2 \text{ and } b = 5$$

$$\text{Hence, } \frac{a+b}{b} = \frac{3}{5} = 0.6$$

Q6. If $\sin^{-1} \frac{1}{3} + \sin^{-1} \frac{2}{3} = \sin^{-1} x$, then the value of x is

A. 0

B. $\frac{(\sqrt{5}-4\sqrt{2})}{9}$

C. $\frac{(\sqrt{5}+4\sqrt{2})}{9}$

D. $\frac{\pi}{2}$
Ans: $\frac{(\sqrt{5}+4\sqrt{2})}{9}$

Solution: Given that, $\sin^{-1} \frac{1}{3} + \sin^{-1} \frac{2}{3} = \sin^{-1} x$

Taking sine on both sides

$$\Rightarrow \left(\frac{1}{3} \sqrt{1 - \frac{4}{9}} + \frac{2}{3} \sqrt{1 - \frac{1}{9}} \right) = x$$

$$\Rightarrow \left(\frac{1}{3} \cdot \frac{\sqrt{5}}{3} + \frac{2}{3} \cdot \frac{\sqrt{8}}{3} \right) = x$$

$$\Rightarrow \left(\frac{\sqrt{5}+4\sqrt{2}}{9} \right) = x$$

$$\therefore x = \left(\frac{\sqrt{5} + 4\sqrt{2}}{9} \right)$$

Q7. The complete solution set of the inequality $\cos^{-1}(\cos 4) > 3x^2 - 4x$ is

A. $\left(0, \frac{2 + \sqrt{6\pi - 8}}{3} \right)$

B. $\left(\frac{2 - \sqrt{6\pi - 8}}{3}, 0 \right)$

C. $(-2, 2)$

D. $\left(\frac{2 - \sqrt{6\pi - 8}}{3}, \frac{2 + \sqrt{6\pi - 8}}{3} \right)$

Ans: $\left(\frac{2 - \sqrt{6\pi - 8}}{3}, \frac{2 + \sqrt{6\pi - 8}}{3} \right)$

Solution: As, $\cos^{-1}(\cos 4) = \cos^{-1}\{\cos(2\pi - 4)\} = 2\pi - 4$

$$\Rightarrow 2\pi - 4 > 3x^2 - 4x$$

$$\Rightarrow 3x^2 - 4x - (2\pi - 4) < 0$$

$$\Rightarrow \frac{2 - \sqrt{6\pi - 8}}{3} < x < \frac{2 + \sqrt{6\pi - 8}}{3}$$

Q8. If $x = \sin(2\tan^{-1}3)$ and $y = \sin\left(\frac{1}{2}\tan^{-1}\frac{4}{3}\right)$, then

A. $2x = 1 - y$

B. $x^2 = 1 - 2y$

C. $x^2 = 1 + y$

D. $y^2 = 2x - 1$

Ans: $y^2 = 2x - 1$

Solution:

Questions with Answer Keys

Let, $x = \sin 2\theta$ (where $\tan \theta = 3$)

$$\Rightarrow x = \frac{2 \tan \theta}{1 + \tan^2 \theta} = \frac{6}{1+9} = \frac{3}{5}$$

$$\text{If } \alpha = \tan^{-1} \frac{4}{3}$$

$$\Rightarrow y = \sin\left(\frac{\alpha}{2}\right) = \frac{1}{\sqrt{2}} \sqrt{1 - \cos \alpha}$$

$$= \frac{1}{\sqrt{2}} \sqrt{1 - \frac{3}{5}} = \frac{1}{\sqrt{5}}$$

$$\therefore y^2 = \frac{1}{5} \Rightarrow y^2 = 2x - 1$$

Q9. $\tan\left(2 \tan^{-1} \frac{1}{5} + \sec^{-1} \frac{\sqrt{5}}{2} + 2 \tan^{-1} \frac{1}{8}\right)$ is equal to:

A. 1

B. 2

C. $\frac{1}{4}$

D. $\frac{5}{4}$

Ans: 2

Solution:

Given,

$$\tan\left(2 \tan^{-1} \frac{1}{5} + \sec^{-1} \frac{\sqrt{5}}{2} + 2 \tan^{-1} \frac{1}{8}\right)$$

$$= \tan\left(2\left(\tan^{-1} \frac{1}{5} + \tan^{-1} \frac{1}{8}\right) + \tan^{-1}\left(\frac{1}{2}\right)\right)$$

$$= \tan\left(2\left(\tan^{-1} \frac{\frac{1}{5} + \frac{1}{8}}{1 - \frac{1}{40}}\right) + \tan^{-1}\left(\frac{1}{2}\right)\right)$$

$$= \tan\left[2 \tan^{-1}\left(\frac{1}{3}\right) + \tan^{-1}\left(\frac{1}{2}\right)\right]$$

$$= \tan\left[\tan^{-1}\left(\frac{2 \times \frac{1}{3}}{1 - \left(\frac{1}{3}\right)^2}\right) + \tan^{-1}\left(\frac{1}{2}\right)\right]$$

Questions with Answer Keys

$$= \tan \left[\tan^{-1} \left(\frac{3}{4} \right) + \tan^{-1} \left(\frac{1}{2} \right) \right]$$

$$= \tan \left[\tan^{-1} \frac{\frac{3}{4} + \frac{1}{2}}{1 - \frac{3}{8}} \right]$$

$$= \tan \tan^{-1} \frac{\frac{5}{4}}{\frac{5}{8}}$$

$$= \tan \tan^{-1} 2$$

$$= 2$$

Q10. The value of $2\sin^{-1} \frac{4}{5} + 2\sin^{-1} \frac{5}{13} + 2\sin^{-1} \frac{16}{65}$ is equal to

A. $\frac{3\pi}{2}$

B. $\frac{\pi}{2}$

C. π

D. 2π

Ans: π

Solution: $2 \left[\sin^{-1} \frac{4}{5} + \sin^{-1} \frac{5}{13} + \sin^{-1} \frac{16}{65} \right]$

$$= 2\sin^{-1} \left(\frac{4}{5} \sqrt{1 - \left(\frac{5}{13} \right)^2} + \frac{5}{13} \sqrt{1 - \left(\frac{4}{5} \right)^2} \right) + 2\sin^{-1} \frac{16}{65}$$

$$= 2\sin^{-1} \left(\frac{48}{65} + \frac{15}{65} \right) + 2\sin^{-1} \left(\frac{16}{65} \right)$$

$$= 2 \left[\sin^{-1} \left(\frac{63}{65} \right) + \sin^{-1} \left(\frac{16}{65} \right) \right]$$

$$= 2 \left[\cos^{-1} \left(\frac{16}{65} \right) + \sin^{-1} \left(\frac{16}{65} \right) \right] \left(\because \sin^{-1} x = \cos^{-1} \sqrt{1 - x^2} \right)$$

$$= \pi$$

Q11. The number of solution of the equation $2 \tan^{-1} x + \cot^{-1} x = \frac{7\pi}{6}$ is

Questions with Answer Keys

A. 0

B. 1

C. 2

D. 3

Ans: 0

Solution:

$$2 \tan^{-1} x + \cot^{-1} x = \frac{7\pi}{6}$$

$$\Rightarrow \tan^{-1} x + \frac{\pi}{2} = \frac{7\pi}{6} \Rightarrow \tan^{-1} x = \frac{2\pi}{3}$$

$$\Rightarrow \tan^{-1} x \in \left(-\frac{\pi}{2}, \frac{\pi}{2} \right) (x \in \phi)$$

Q12. The value of $\tan^{-1} \left[\frac{\sqrt{1-\sin x} + \sqrt{1+\sin x}}{\sqrt{1-\sin x} - \sqrt{1+\sin x}} \right] \left(\forall x \in \left[0, \frac{\pi}{2} \right] \right)$ is equal to

A. $\frac{x}{2} - \frac{\pi}{2}$ B. $\frac{x}{2} + \frac{\pi}{2}$ C. $\frac{x}{2} - \pi$ D. $\frac{\pi}{2} - \frac{x}{2}$ Ans: $\frac{x}{2} - \frac{\pi}{2}$

$$\text{Solution: } \tan^{-1} \left[\frac{\sqrt{1-\sin x} + \sqrt{1+\sin x}}{\sqrt{1-\sin x} - \sqrt{1+\sin x}} \right]$$

$$= \tan^{-1} \left[\frac{(\sqrt{1-\sin x} + \sqrt{1+\sin x})}{(\sqrt{1-\sin x} - \sqrt{1+\sin x})} \cdot \frac{(\sqrt{1-\sin x} + \sqrt{1+\sin x})}{(\sqrt{1-\sin x} + \sqrt{1+\sin x})} \right]$$

$$= \tan^{-1} \left[\frac{(1-\sin x) + (1+\sin x) + 2\sqrt{1-\sin^2 x}}{(1-\sin x) - (1+\sin x)} \right] = \tan^{-1} \left[\frac{2(1+\cos x)}{-2\sin x} \right]$$

$$= \tan^{-1} \left[\frac{-2\cos^2 \left(\frac{x}{2} \right)}{2\sin \left(\frac{x}{2} \right) \cos \left(\frac{x}{2} \right)} \right] = \tan^{-1} \left(-\cot \frac{x}{2} \right) = \tan^{-1} \left[\cot \left(\pi - \frac{x}{2} \right) \right]$$

$$= \frac{\pi}{2} - \cot^{-1} \left[\cot \left(\pi - \frac{x}{2} \right) \right] = \frac{\pi}{2} - \left(\pi - \frac{x}{2} \right) = \frac{x}{2} - \frac{\pi}{2}$$

Q13. If the equation $\sin^{-1}(4x^2 - 12x + 10) + \cos^{-1}(12x - 4x^2 - 10) + \lambda x = 0$ has a real solution, then λ is equal to

A. $\frac{\pi}{4}$

B. $-\pi$

C. $\frac{\pi}{2}$

D. $\frac{-\pi}{2}$

Ans: $-\pi$

Solution:

$$\sin^{-1}(1 + (2x - 3)^2) + \cos^{-1}(-1 - (2x - 3)^2) + \lambda x = 0$$

$$x = \frac{3}{2}$$

$$\Rightarrow \frac{\pi}{2} + \pi + \frac{3\lambda}{2} = 0 \Rightarrow \lambda = -\pi.$$

Q14. If $\cos^{-1} x + \cos^{-1} y + \cos^{-1} z = \pi$, then

A. $x^2 + y^2 + z^2 + xyz = 0$

B. $x^2 + y^2 + z^2 + 2xyz = 0$

C. $x^2 + y^2 + z^2 + xyz = 1$

D. $x^2 + y^2 + z^2 + 2xyz = 1$

Ans: $x^2 + y^2 + z^2 + 2xyz = 1$

Solution: Given that $\cos^{-1} x + \cos^{-1} y + \cos^{-1} z = \pi$

Questions with Answer Keys

MathonGo

$$\Rightarrow \cos^{-1}(x) + \cos^{-1}(y) + \cos^{-1}(z) = \cos^{-1}(-1)$$

$$\Rightarrow \cos^{-1}(x) + \cos^{-1}(y) = \cos^{-1}(-1) - \cos^{-1}(z)$$

$$\Rightarrow \cos^{-1}\left(xy - \sqrt{1-x^2}\sqrt{1-y^2}\right) = \cos^{-1}\{(-1)(z)\}$$

$$\Rightarrow xy - \sqrt{(1-x^2)(1-y^2)} = -2$$

$$\Rightarrow (xy + z) = \sqrt{(1-x^2)(1-y^2)}$$

Squaring both sides we get $x^2 + y^2 + z^2 + 2xyz = 1$ Trick: Put $x = y = z = \frac{1}{2}$, so that

$$\cos^{-1} \frac{1}{2} + \cos^{-1} \frac{1}{2} + \cos^{-1} \frac{1}{2} = \pi$$

Obviously (d) holds for these values of x, y, z .

Q15. If $(\cot^{-1} x)^2 - 7(\cot^{-1} x) + 10 > 0$, then the range of x will be

A. $(-\infty, \cot 2)$

B. $(-\infty, \cot 5)$

C. $(\cot 2, \cot 5)$

D. $(\cot 2, \infty)$

Ans: $(\cot 2, \infty)$

Solution: $(\cot^{-1} x - 5)(\cot^{-1} x - 2) > 0$

$$\Rightarrow \cot^{-1} x \in (-\infty, 2) \cup (5, \infty)$$

$$\Rightarrow \cot^{-1} x \in (0, 2) \quad (\text{Taking intersection with range of } \cot^{-1} x)$$

$$\Rightarrow x \in (\cot 2, \infty)$$

Q16. The value of a for which $ax^2 + \sin^{-1}(x^2 - 2x + 2) + \cos^{-1}(x^2 - 2x + 2) = 0$ has a real solution, is

Questions with Answer Keys

MathonGo

A. $-\frac{2}{\pi}$

B. $\frac{2}{\pi}$

C. $-\frac{\pi}{2}$

D. $\frac{\pi}{2}$

Ans: $-\frac{\pi}{2}$

Solution: Here, $x^2 - 2x + 2 = (x - 1)^2 + 1 \geq 1$

But, $-1 \leq (x^2 - 2x + 2) \leq 1$

Which is possible only when

$$x^2 - 2x + 2 = 1$$

$$\Rightarrow x = 1$$

Then, $a(1)^2 + \sin^{-1}(1) + \cos^{-1}(1) = 0$

$$\Rightarrow a + \frac{\pi}{2} + 0 = 0$$

$$\Rightarrow a = -\frac{\pi}{2}$$

Q17. Number of real roots of the equation $\sin^{-1} \sin x = \cos^{-1} \cos 4$ in $[0, 2\pi]$ is

A. 0

B. 1

Questions with Answer Keys

C. 2

D. more than 2

Ans: 0

Solution:

$$\cos^{-1} \cos 4 = 2\pi - 4 > \frac{\pi}{2} \text{ and } \sin^{-1} \sin x \leq \frac{\pi}{2}$$

 $\Rightarrow \sin^{-1} \sin x = \cos^{-1} \cos 4$ has no real root.

Q18. The greatest and the least value of $(\sin^{-1} x)^3 + (\cos^{-1} x)^3$ are

A. $-\frac{\pi}{2}, \frac{\pi}{2}$

B. $-\frac{\pi^3}{8}, \frac{\pi^3}{8}$

C. $\frac{7\pi^3}{8}, \frac{\pi^3}{32}$

D. None of these

Ans: $\frac{7\pi^3}{8}, \frac{\pi^3}{32}$

Solution: We have $(\sin^{-1} x)^3 + (\cos^{-1} x)^3$

$$= (\sin^{-1} x + \cos^{-1} x)^3 - 3 \sin^{-1} x \cos^{-1} x (\sin^{-1} x + \cos^{-1} x)$$

$$= \frac{\pi^3}{8} - 3 (\sin^{-1} x \cos^{-1} x) \frac{\pi}{2}$$

$$= \frac{\pi^3}{8} - \frac{3\pi}{2} \sin^{-1} x \left(\frac{\pi}{2} - \sin^{-1} x \right)$$

$$= \frac{\pi^3}{8} - \frac{3\pi^2}{4} \sin^{-1} x + \frac{3\pi}{2} (\sin^{-1} x)^2$$

$$= \frac{\pi^3}{8} + \frac{3\pi}{2} \left[(\sin^{-1} x)^2 - \frac{\pi}{2} \sin^{-1} x \right]$$

$$= \frac{\pi^3}{8} + \frac{3\pi}{2} \left[\left(\sin^{-1} x - \frac{\pi}{4} \right)^2 \right] - \frac{3\pi^3}{32}$$

$$= \frac{\pi^3}{32} + \frac{3\pi}{2} \left(\sin^{-1} x - \frac{\pi}{4} \right)^2$$

\therefore The least value is $\frac{\pi^3}{32}$

and since $\left(\sin^{-1} x - \frac{\pi}{4} \right)^2 \leq \left(\frac{3\pi}{4} \right)^2$

\therefore The greatest value is $\frac{\pi^3}{32} + \frac{9\pi^2}{16} \times \frac{3\pi}{2} = \frac{7\pi^3}{8}$.

Q19. The real solutions of the equation $\tan^{-1}\sqrt{x(x+1)} + \sin^{-1}\sqrt{x^2+x+1} = \frac{\pi}{2}$ are

A. $-1, 0$

B. $0, 1$

C. $-1, 1$

D. $-1, 2$

Ans: $-1, 0$

Solution: $\tan^{-1}\sqrt{x(x+1)} + \sin^{-1}\sqrt{x^2+x+1} = \frac{\pi}{2}$

$$\Rightarrow \tan^{-1}\sqrt{x^2+x} + \sin^{-1}\sqrt{x^2+x+1} = \frac{\pi}{2}$$

This equation holds true, if

$$x^2+x \geq 0 \text{ and } 0 \leq x^2+x+1 \leq 1$$

Now, $x^2+x \geq 0$ and $0 \leq x^2+x+1 \leq 1$

$$\Rightarrow x^2+x \geq 0 \text{ and } x^2+x+1 \leq 1 \quad [\because x^2+x+1 > 0 \text{ for all } x]$$

$$\Rightarrow x^2+x \geq 0 \text{ and } x^2+x \leq 0$$

$$\Rightarrow x^2+x = 0 \Rightarrow x = 0, -1.$$

Clearly, these two values satisfy the given equation. Hence, $x = -1, 0$ are the solutions of the given equation.

Q20. The number of integers for which the equation $\sin^{-1} x + \cos^{-1} x + \tan^{-1} x = n$ has real solution(s) is

A. 0

B. 1

C. 2

D. 3

Ans: 2

Solution:

Given equation is $\frac{\pi}{2} + \tan^{-1} x = n, \forall x \in [-1, 1]$

Now LHS $\in \left[\frac{\pi}{4}, \frac{3\pi}{4}\right]$

Integers in this interval are 1 & 2

Hence, there are 2 integers for which the equation has real solutions

Q21. Let $a = (\sin^{-1} x)^{\sin^{-1} x}$, $b = (\sin^{-1} x)^{\cos^{-1} x}$, $c = (\cos^{-1} x)^{\sin^{-1} x}$, $d = (\cos^{-1} x)^{\cos^{-1} x}$ and if $x \in (0, 1)$, then

A. $a > b > d > c$

B. $d > c > a > b$

C. $b > a > d > c$

D. $a < b < d < c$

Ans: $d > c > a > b$

Solution:

It is given that $a = (\sin^{-1} x)^{\sin^{-1} x}$,

Questions with Answer Keys

$$b = (\sin^{-1} x)^{\cos^{-1} x}, c = (\cos^{-1} x)^{\sin^{-1} x}$$

$$d = (\cos^{-1} x)^{\cos^{-1} x}$$

Also, here $x \in (0, 1)$

$$\Rightarrow \cos^{-1} x < \sin^{-1} x$$

Also, $\cos^{-1} x > 1$

and $\sin^{-1} x < 1$

$\therefore (\cos^{-1} x)^{\cos^{-1} x}$ is greatest and $(\sin^{-1} x)^{\cos^{-1} x}$ is least.

$$\Rightarrow (\sin^{-1} x)^{\sin^{-1} x} < (\cos^{-1} x)^{\sin^{-1} x}$$

$$\Rightarrow d > c > a > b$$

Q22.

The value of $\tan^{-1}\left(\frac{9}{19}\right) + \tan^{-1}\left(\frac{9}{49}\right) + \tan^{-1}\left(\frac{9}{97}\right) + \tan^{-1}\left(\frac{9}{163}\right) + \dots \infty$ equals

A. $\tan^{-1}(3)$

B. $\tan^{-1}\left(\frac{1}{3}\right)$

C. $\tan^{-1}\left(\frac{2}{3}\right)$

D. $\tan^{-1}\left(\frac{3}{2}\right)$

Ans: $\tan^{-1}\left(\frac{3}{2}\right)$

Solution:

$$S = \lim_{m \rightarrow \infty} \sum_{n=1}^m \tan^{-1}\left(\frac{9}{9n^2+3n+7}\right) = \lim_{m \rightarrow \infty} \sum_{n=1}^m \tan^{-1}\left(\frac{1}{1+n^2+\frac{n}{3}-\frac{2}{9}}\right)$$

Questions with Answer Keys

$$= \lim_{m \rightarrow \infty} \sum_{n=1}^m \tan^{-1} \left(\frac{\left(n + \frac{2}{3}\right) - \left(n - \frac{1}{3}\right)}{1 + \left(n + \frac{2}{3}\right) \left(n - \frac{1}{3}\right)} \right) = \lim_{m \rightarrow \infty} \sum_{n=1}^m \left[\tan^{-1} \left(n + \frac{2}{3} \right) - \tan^{-1} \left(n - \frac{1}{3} \right) \right]$$

Q23. If $y = \tan^{-1} \frac{1}{1+x+x^2} + \tan^{-1} \frac{1}{x^2+3x+3} + \tan^{-1} \frac{1}{x^2+5x+7} + \dots +$ upto $2n$ terms ($\forall x \geq 0$), then $y(0)$ is

- A. $\tan^{-1}(n)$
- B. $\tan^{-1}(2n)$
- C. $2\tan^{-1}(n)$

D. 0

Ans: $\tan^{-1}(2n)$

Solution: $y = \tan^{-1} \frac{1}{1+x+x^2} + \tan^{-1} \frac{1}{x^2+3x+3} + \dots + 2n$ terms

$$= \tan^{-1} \frac{(x+1) - x}{1+x(1+x)} + \tan^{-1} \frac{(x+2) - (x+1)}{1+(x+1)(x+2)} + \dots + (2n \text{ terms})$$

$$= \tan^{-1}(x+1) - \tan^{-1}x + \tan^{-1}(x+2) - \tan^{-1}(x+1) + \dots + \tan^{-1}(x+2n) - \tan^{-1}(x+(2n-1))$$

$$= \tan^{-1}(x+2n) - \tan^{-1}x$$

$$y(0) = \tan^{-1}(2n)$$

Q24. The value of the expression $\cot^{-1} \frac{1}{2} + \cot^{-1} \frac{9}{2} + \cot^{-1} \frac{25}{2} + \cot^{-1} \frac{49}{2} + \dots$ upto n terms is

- A. $\tan^{-1} 2n$
- B. $\tan^{-1}(2n-1)$
- C. $\tan^{-1} n$
- D. $\tan^{-1} 2n - \tan^{-1} 1$

Ans: $\tan^{-1} 2n$

Solution:

Questions with Answer Keys

Given expression = $\tan^{-1} 2 + \tan^{-1} \frac{2}{9} + \tan^{-1} \frac{2}{25} + \tan^{-1} \frac{2}{49} + \dots$

$$\text{General term} = \frac{2}{(2n-1)^2} = \frac{2}{4n^2-4n+1} = \frac{2}{1+4n(n-1)} = \frac{2n-(2n-2)}{1+2n(2n-2)}$$

$$T_n = \tan^{-1} 2n - \tan^{-1}(2n-2)$$

\therefore Sum of the series

$$= \tan^{-1} 2 - \tan^{-1} 0 + \tan^{-1} 4 - \tan^{-1} 2 + \tan^{-1} 6 - \tan^{-1} 4 + \dots + \tan^{-1} 2n - \tan^{-1}(2n-2)$$

$$= \tan^{-1} 2n - \tan^{-1} 0 = \tan^{-1} 2n$$

Q25. The value(s) of x satisfying the equation $\sin^{-1}(1-x) - 2\sin^{-1} x = \frac{\pi}{2}$ is/are

A. 0

B. $\frac{1}{2}$

C. $0, \frac{1}{2}$

D. $-\frac{1}{2}$

Ans: 0

Solution: We have,

$$\sin^{-1}(1-x) - 2\sin^{-1} x = \frac{\pi}{2}$$

$$\Rightarrow \sin^{-1}(1-x) = \frac{\pi}{2} + 2\sin^{-1} x$$

$$\Rightarrow 1-x = \sin\left(\frac{\pi}{2} + 2\sin^{-1} x\right)$$

$$\Rightarrow 1-x = \cos(2\sin^{-1} x)$$

$$\Rightarrow 1-x = \cos\{\cos^{-1}(1-2x^2)\} [\because 2\sin^{-1} x = \cos^{-1}(1-2x^2)]$$

$$\Rightarrow 1-x = (1-2x^2)$$

$$\Rightarrow x = 2x^2 \Rightarrow x(2x-1) = 0 \Rightarrow x = 0, \frac{1}{2}$$

For, $x = \frac{1}{2}$, we have

Questions with Answer Keys

$$\text{LHS} = \sin^{-1}(1-x) - 2\sin^{-1}x$$

$$= \sin^{-1}\frac{1}{2} - 2\sin^{-1}\frac{1}{2} = -\sin^{-1}\frac{1}{2} = \frac{-\pi}{6} \neq \text{R. H. S.}$$

So, $x = \frac{1}{2}$ is not a root of the given equation.

Clearly, $x = 0$ satisfies the equation

Here, $x = 0$ is the root of the given equation.

Q26. Solution set of $[\sin^{-1}x] > [\cos^{-1}x]$, where $[.]$ denotes the greatest integer function, is

A. $\left[\frac{1}{\sqrt{2}}, 1\right]$

B. $(\cos 1, \sin 1)$

C. $[\sin 1, 1]$

D. None of these

Ans: $[\sin 1, 1]$

Solution: $\because [\sin^{-1}x] > [\cos^{-1}x]$

$$\Rightarrow x > 0$$

Here, $[\cos^{-1}x] = \begin{cases} 0, & x \in (\cos 1, 1] \\ 1, & x \in (0, \cos 1] \end{cases}$

and $[\sin^{-1}x] = \begin{cases} 0, & x \in (0, \sin 1) \\ 1, & x \in [\sin 1, 1] \end{cases}$

$$\therefore x \in [\sin 1, 1]$$

Q27. If $\cot^{-1}(\alpha) = \cot^{-1}2 + \cot^{-1}8 + \cot^{-1}18 + \cot^{-1}32 + \dots$ upto 100 terms, then α is:

Questions with Answer Keys

MathonGo

A. 1.01

B. 1.00

C. 1.02

D. 1.03

Ans: 1.01

Solution:

$$\text{Given } \cot^{-1}(\alpha) = \cot^{-1}(2) + \cot^{-1}(8) + \cot^{-1}(18) + \dots$$

$$= \sum_{n=1}^{100} \tan^{-1}\left(\frac{2}{4n^2}\right)$$

$$= \sum_{n=1}^{100} \tan^{-1}\left(\frac{(2n+1)-(2n-1)}{1+(2n+1)(2n-1)}\right)$$

$$= \sum_{n=1}^{100} \tan^{-1}(2n+1) - \tan^{-1}(2n-1)$$

$$= \tan^{-1} 201 - \tan^{-1} 1$$

$$= \tan^{-1}\left(\frac{200}{202}\right)$$

$$\therefore \cot^{-1}(\alpha) = \cot^{-1}\left(\frac{202}{200}\right)$$

$$\alpha = 1.01$$

Q28. For $k \in \mathbb{R}$, let the solutions of the equation

$$\cos(\sin^{-1}(x \cot(\tan^{-1}(\cos(\sin^{-1} x)))) = k, 0 < |x| < \frac{1}{\sqrt{2}} \text{ be } \alpha \text{ and } \beta, \text{ where the inverse}$$

trigonometric functions take only principal values. If the solutions of the equation

$$x^2 - bx - 5 = 0 \text{ are } \frac{1}{\alpha^2} + \frac{1}{\beta^2} \text{ and } \frac{\alpha}{\beta}, \text{ then } \frac{b}{k^2} \text{ is equal to } \underline{\hspace{2cm}}.$$

Ans: 12

Solution:

Questions with Answer Keys

Given,

$$\cos(\sin^{-1}(x \cot(\tan^{-1}(\cos(\sin^{-1} x)))))) = k$$

Now simplifying $\cos(\sin^{-1} x) = \cos(\cos^{-1} \sqrt{1-x^2}) = \sqrt{1-x^2}$

So, $\cos(\sin^{-1}(x \cot(\tan^{-1}(\cos(\sin^{-1} x)))))) = k$

becomes $\cos(\sin^{-1}(x \cot(\tan^{-1} \sqrt{1-x^2}))) = k$

And now solving $\cot(\tan^{-1} \sqrt{1-x^2}) = \cot \cot^{-1} \left(\sqrt{\frac{1}{1-x^2}} \right) = \frac{1}{\sqrt{1-x^2}}$

So, $\cos(\sin^{-1}(x \cot(\tan^{-1} \sqrt{1-x^2}))) = k$ becomes

$$\cos\left(\sin^{-1}\left(\frac{x}{\sqrt{1-x^2}}\right)\right) = k$$

Now solving $\cos\left(\sin^{-1}\left(\frac{x}{\sqrt{1-x^2}}\right)\right) = \frac{\sqrt{1-2x^2}}{\sqrt{1-x^2}}$

So, $\frac{\sqrt{1-2x^2}}{\sqrt{1-x^2}} = k$

$$\Rightarrow 1 - 2x^2 = k^2(1 - x^2)$$

$$\Rightarrow (k^2 - 2)x^2 = k^2 - 1$$

$$\Rightarrow x^2 = \frac{k^2 - 1}{k^2 - 2}$$

So, roots are $\alpha = \sqrt{\frac{k^2 - 1}{k^2 - 2}} \Rightarrow \alpha^2 = \frac{k^2 - 1}{k^2 - 2}$

And $\beta = \sqrt{\frac{k^2 - 1}{k^2 - 2}} \Rightarrow \beta^2 = \frac{k^2 - 1}{k^2 - 2}$

Now finding $\frac{1}{\alpha^2} + \frac{1}{\beta^2} = 2\left(\frac{k^2 - 2}{k^2 - 1}\right)$ and $\frac{\alpha}{\beta} = -1$

So, sum of roots of $x^2 - bx - 5 = 0$ will be $= \frac{1}{\alpha^2} + \frac{1}{\beta^2} + \frac{\alpha}{\beta} = b$

Questions with Answer Keys

$$\Rightarrow \frac{2(k^2-2)}{k^2-1} - 1 = b \quad \dots (1)$$

Product of roots of $x^2 - bx - 5 = 0$ will be $= \left(\frac{1}{\alpha^2} + \frac{1}{\beta^2} \right) \frac{\alpha}{\beta} = -5$

$$\Rightarrow \frac{2(k^2-2)}{k^2-1} (-1) = -5$$

$$\Rightarrow 2k^2 - 4 = 5k^2 - 5$$

$$\Rightarrow 3k^2 = 1 \Rightarrow k^2 = \frac{1}{3} \quad \dots \text{Put in (1)}$$

$$\Rightarrow b = \frac{2(k^2-2)}{k^2-1} - 1 = 5 - 1 = 4$$

$$\frac{b}{k^2} = \frac{4}{\frac{1}{3}} = 12$$

Q29. The number of solutions of the equation $\sin^{-1}x = (\sin x)^{-1}$ is/are

A. one

B. two


C. three

D. zero

Ans: two

Solution: $\therefore (\sin x)^{-1} = \frac{1}{\sin x} = \operatorname{cosec} x$

Now, from graphs of $\sin^{-1}x$, $\operatorname{cosec} x$



Questions with Answer Keys

Clearly, both graph intersects at two points

\therefore two solutions

Q30. The number of real solutions (x, y) where $|y| = \sin x, y = \cos^{-1}(\cos x), -2\pi \leq x \leq 2\pi$, is

A. 2

B. 1

C. 3

D. 4

E. 0

Ans: 2

Solution:

$$\text{In } [0, \pi], |y| = \sin x, y = \cos^{-1}(\cos x) = x$$

$$\text{In } [\pi, 2\pi], |y| = \sin x, y = \cos^{-1}\{\cos(2\pi - x)\} = 2\pi - x$$


$$\text{In } [-\pi, 0], |y| = \sin x, y = \cos^{-1}\{\cos(-x)\} = -x$$

$$\text{In } [-2\pi, -\pi], |y| = \sin x, y = \cos^{-1}\{\cos(2\pi + x)\} = 2\pi + x$$

Plotting the graphs, we have

Questions with Answer Keys

MathonGo



There are 2 solutions, i.e., $(0, 0)$ and $(-2\pi, 0)$.

MathonGo

Q1 (4)	Q2 (3)	Q3 (4)	Q4 (1)
Q5 (0.6)	Q6 (3)	Q7 (4)	Q8 (4)
Q9 (2)	Q10 (3)	Q11 (1)	Q12 (1)
Q13 (2)	Q14 (4)	Q15 (4)	Q16 (3)
Q17 (1)	Q18 (3)	Q19 (1)	Q20 (3)
Q21 (2)	Q22 (4)	Q23 (2)	Q24 (1)
Q25 (1)	Q26 (3)	Q27 (1)	Q28 (12)
Q29 (2)	Q30 (1)		