
Miglitol

Helps lower your blood sugar

What does it do?

- Miglitol helps lower your blood sugar by slowing down the digestion and absorption of carbohydrates (sugars and starches) from food.
- It helps prevent a sudden rise in blood sugar, especially after meals.



How should I take it?

- Take Miglitol with the first bite of your meal.
- Take it exactly as prescribed by your doctor.
- It is usually taken with each main meal.
- Take it at the same time each day.
- Do not skip meals without medical advice.


Take with the first bite of your meal


Take at the same time every day

Common side effects

- Gas, bloating or increased stomach discomfort.
- Loose stools or diarrhoea.
- Stomach pain or abdominal discomfort.
- Nausea or feeling sick.
- These stomach-related side effects may become less troublesome as your body gets used to the medicine.


Gas, bloating or stomach discomfort


Loose stools or diarrhoea


Stomach pain or abdominal discomfort


Nausea or feeling sick

If blood sugar becomes low

Miglitol alone usually does not cause low blood sugar. However, low blood sugar may occur when it is taken with insulin or other diabetes medicines.

If you feel shaky, sweaty, dizzy, hungry or weak, your blood sugar may be low. Take 15 g of fast-acting sugar (e.g. glucose tablets or glucose solution), wait 15 minutes and check again. If it is still low, repeat the same steps.


1. Take 15 grams of fast-acting sugar (e.g., glucose tablets or glucose solution)


2. Wait 15 minutes and check your blood sugar again (if possible).



3. If still low, repeat the same steps.

Important warning

- Tell your doctor if you develop severe or persistent stomach pain, severe diarrhoea, or unusual abdominal discomfort.
- Do not rely on ordinary table sugar because Miglitol slows its digestion. Use glucose/dextrose for low blood sugar.




Severe or persistent stomach pain or diarrhoea - tell your doctor

Do not stop Miglitol without consulting your doctor or pharmacist.