
Voglibose

Helps lower your blood sugar

What does it do?

- Voglibose helps lower your blood sugar by slowing down the digestion and absorption of carbohydrates (sugars and starches) from food.
- It helps prevent a sudden rise in blood sugar, especially after meals.



How should I take it?

- Take Voglibose just before or with the first bite of your meal.
- Take it exactly as prescribed by your doctor.
- It is usually taken with each main meal (breakfast, lunch and dinner).
- Take the medicine at the same time each day.
- Do not skip meals without medical advice.





Common side effects

- Gas, bloating or increased stomach discomfort.
- Loose stools or diarrhoea.
- Stomach pain or abdominal discomfort.
- Nausea or feeling sick.
- These stomach-related side effects may become less troublesome as your body gets used to the medicine.









If blood sugar becomes low

Voglibose alone usually does not cause low blood sugar. However, low blood sugar may occur when it is taken with insulin or other diabetes medicines.

If you feel shaky, sweaty, dizzy, hungry or weak, your blood sugar may be low. Take 15 grams of fast-acting sugar, wait 15 minutes, and check again. If it is still low, repeat the same steps.





If you become unconscious or cannot swallow, do not give anything by mouth and **seek emergency medical help**.



Important warning

Tell your doctor if you develop severe or persistent stomach pain, severe diarrhoea, constipation, or unusual abdominal discomfort.



Do not stop Voglibose without consulting your doctor or pharmacist.