Saxagliptin

Helps lower your blood sugar

What does it do?

- Helps your body use insulin better.
- Increases the amount of insulin released when needed.
- Reduces the extra sugar made by your liver.
- Helps control blood sugar levels.



How should I take it?

- Take it once daily.
- Can be taken with or without food.
- Take it at the same time every day.
- Swallow the tablet whole with water.
- Do not stop the medicine on your own.



Common side effects

- Stomach upset (nausea, bloating, or stomach pain).
- Runny nose or cold symptoms.
- Sore throat.
- Headache.
- Diarrhoea (loose motions).

These side effects are usually mild and may improve with time.

Symptoms of low blood sugar (hypoglycemia)

You may feel:

- Sweating
- Shaking
- Dizziness
- Hunger
- Weakness
- Confusion
- Irritability
- Fast heartbeat
- Blurred vision
- Difficulty in concentrating

15–15 Rule for Low Blood Sugar

If you feel symptoms of low blood sugar (such as shakiness, sweating, dizziness, hunger or confusion):

1. **Take 15 grams of fast-acting sugar** (e.g., glucose tablets, ½ cup fruit juice or 1 tablespoon sugar).
2. **Wait 15 minutes** and check your blood sugar again (if possible).
3. **If still low, repeat 15 grams** and recheck after 15 minutes.

Once your blood sugar is above 70 mg/dL, have a small snack or meal to prevent it from dropping again.

Seek medical attention immediately if you develop:

- Severe stomach pain, nausea, vomiting or diarrhoea (which does not get better).
- Severe sore throat, fever or signs of infection.
- Allergic reaction (face or throat swelling, difficulty breathing).
- Unusual or persistent symptoms.



Do not stop saxagliptin without consulting your doctor or pharmacist.