Vildagliptin

Helps lower your blood sugar

What does it do?

- Helps your body release insulin when blood sugar is high.
- Reduces a hormone that raises blood sugar.
- Helps control blood sugar levels.



How should I take it?

- Take exactly as prescribed by your doctor (often once or twice daily).
- Take with or without food.
- Take at the same times each day.
- Do not stop or change the dose on your own.
- If you miss a dose, take it when remembered unless it is nearly time for the next dose. Do not double the dose.



Common side effects

- Headache
- Dizziness
- Nausea
- Stomach discomfort
- Joint pain
- Cold-like symptoms (runny nose or sore throat)

Side effects may vary from person to person.



Symptoms of low blood sugar (hypoglycaemia)

- Shaking or trembling
- Sweating
- Sudden hunger
- Dizziness
- Fast heartbeat
- Headache
- Confusion or difficulty concentrating
- Weakness or irritability

Vildagliptin alone rarely causes low blood sugar, but risk is higher with insulin or sulfonylureas.



15–15 Rule for Low Blood Sugar



1. **If you are awake and can** swallow, take 15 g of fast-acting sugar (for example, glucose tablets according to the label or 150 mL regular juice/sugary drink).



2. **Wait 15 minutes,** then check blood sugar if possible.



3. If still below **70 mg/dL,** take another 15 g and recheck after 15 minutes.

Once blood sugar is above 70 mg/dL and you feel better, eat a snack or meal if your next meal is not soon.

Seek medical attention immediately if you develop



- Severe, persistent stomach pain, especially spreading to the back, with or without vomiting.
- Yellow skin or eyes, dark urine, pale stools, or unusual tiredness.
- Severe allergic reaction: swelling of the face, lips, tongue or throat, or difficulty breathing.
- Severe skin rash, blistering or peeling.
- Unusual bleeding or bruising.
- Repeated or severe low blood sugar, fainting or seizures.

Do not stop vildagliptin without consulting your doctor or pharmacist.