
Repaglinide



Helps control your blood sugar after meals

What does it do?

- Stimulates your pancreas to release insulin.
- Helps lower blood sugar, especially after meals.
- Works best with a healthy diet and regular physical activity.
- Used for type 2 diabetes. It is not used for type 1 diabetes or diabetic ketoacidosis.



How should I take it?

- Take repaglinide **within 30 minutes before a meal.**
- **If you skip a meal, skip that dose.**
- If you have an additional meal, take the prescribed dose for that meal.
- Take it exactly as prescribed by your doctor.
- Do not take an extra dose to make up for a missed dose.
- Continue taking it even if you feel well, unless your doctor tells you to stop.



Common side effects

- Low blood sugar (hypoglycemia)
- Headache
- Dizziness
- Tiredness or weakness
- Nausea
- Increased sweating
- Stomach discomfort

Low blood sugar is the most important side effect to watch for.



Symptoms of low blood sugar (hypoglycemia)

You may feel:

- Shakiness
- Sweating
- Hunger
- Dizziness
- Weakness
- Irritability or nervousness
- Headache
- Confusion
- Difficulty in concentrating
- Fast heartbeat



15-15 Rule for Low Blood Sugar

If you have symptoms of low blood sugar (such as shakiness, sweating, dizziness, hunger or confusion):



1. **Take 15 grams of fast-acting sugar** (e.g., 3–4 glucose tablets, ½ cup fruit juice or 1 tablespoon sugar/honey).



2. **Wait 15 minutes** and check your blood sugar again (if possible).



3. **If still low,** repeat the 15 grams of sugar and wait another 15 minutes.

Once your blood sugar is above 70 mg/dL, have a meal or snack if your next meal is not soon.

Seek medical attention immediately if you develop:

- Severe confusion or unusual behavior
- Loss of consciousness
- Seizures
- Severe weakness
- Inability to swallow or treat low blood sugar yourself

Severe hypoglycemia requires urgent treatment.





Do not stop repaglinide without consulting your doctor or pharmacist.