Semaglutide

Helps lower your blood sugar

What does it do?

- Helps your body release insulin.
- Reduces excess sugar production by your liver.
- Helps control hunger and may support weight management.



How should I take it?

- Take it exactly as prescribed by your doctor.
- Swallow the tablet whole with a glass of water.
- Take it on an empty stomach (usually 30 minutes before food) or as directed by your doctor.
- Do not crush, chew or break the tablet.
- Do not stop the medicine on your own.



Common side effects

- Nausea
- Vomiting
- Diarrhoea
- Constipation (less common)
- Abdominal pain
- Decreased appetite
- Headache
- Dizziness
- Fatigue

These side effects may be more noticeable when treatment is started and may improve with continued use.



Symptoms of low blood sugar (hypoglycaemia)

You may feel:

- Shakiness
- Sweating
- Dizziness
- Hunger
- Weakness
- Confusion
- Irritability
- Difficulty in concentrating



15-15 Rule for Low Blood Sugar

1. **Take 15 grams of fast-acting sugar** (e.g., 3–4 glucose tablets, ½ cup fruit juice, or 1 tablespoon sugar/honey).
2. **Wait 15 minutes** and check your blood sugar again (if possible).
3. **If still low,** repeat the 15 grams of sugar and wait another 15 minutes.

Once above 70 mg/dL, have a small snack or meal if appropriate.



Seek medical attention immediately if you develop:

- **Severe allergic reaction** (itching or hives, swelling in your face or hands, swelling or tingling in your mouth or throat, chest tightness, trouble breathing).
- Blurred vision or any change in vision.
- Decreased urination, lower back or side pain, blood in your urine.
- Shaking, trembling, sweating, fast or pounding heartbeat, lightheadedness, hunger, confusion.
- Sudden and severe stomach pain, diarrhoea, nausea, vomiting, fever, yellow eyes or skin.
- Constipation and loss of appetite (less common).

Do not stop semaglutide without consulting your doctor or pharmacist.