Canagliflozin

Helps lower your blood sugar and may protect your heart and kidneys

What does it do?

- Helps your kidneys remove extra sugar from your body through urine.
- Works best with a healthy diet and regular physical activity.



Extra sugar comes out in urine

How should I take it?

- Take exactly as prescribed by your doctor, usually once daily.
- Can be taken with or without food.
- Take at the same time every day.
- Drink enough water unless your doctor has told you to restrict fluids.
- Do not stop the medicine on your own.



Common side effects

- You may pass urine more often or in larger amounts.
- Burning or pain while passing urine.
- Frequent urges to urinate.
- Lower abdominal discomfort.
- Female genital fungal infection – itching, redness, irritation, or unusual discharge.
- Male genital fungal infection – redness, itching, swelling, or irritation.

Symptoms of low blood sugar (hypoglycemia)

You may feel:

- Sweating
- Shaking
- Dizziness
- Hunger
- Weakness
- Confusion
- Irritability
- Blurred vision
- Difficulty in concentrating

Fast heartbeat

15–15 Rule for Low Blood Sugar

If you feel symptoms of low blood sugar (such as shakiness, sweating, dizziness, hunger or confusion):

1. **Take 15 grams of fast-acting sugar** (e.g., glucose tablets, ½ cup fruit juice or 1 tablespoon sugar).
2. **Wait 15 minutes** and check your blood sugar again (if possible).
3. **If still low, repeat 15 grams** and recheck after 15 minutes.

Once your blood sugar is above 70 mg/dL, have a small snack or meal to prevent it from dropping again.

Seek medical attention immediately if you develop:

- Severe or repeated urinary symptoms with fever (possible kidney infection).
- Severe dizziness, weakness or signs of dehydration (even if blood sugar is normal).
- Very low blood sugar symptoms that do not improve after taking sugar.
- Severe pain, redness or swelling around genital or anal area (possible serious infection).
- Severe nausea, vomiting, abdominal pain, unusual tiredness, or difficulty breathing (possible diabetic ketoacidosis, even if blood sugar is not very high).
- Any new foot problems, unusual wounds, or signs of infection.



Do not stop Canagliflozin without consulting your doctor or pharmacist.