
Dapagliflozin

Helps lower your blood sugar and may protect your heart and kidneys

What does it do?

- Helps your kidneys remove extra sugar from your body through urine.
- Also helps your body remove some extra salt and water, which may protect your heart and kidneys.



How should I take it?

- Take exactly as prescribed by your doctor, usually once daily.
- Can be taken with or without food.
- Take at the same time every day.
- Drink enough water unless your doctor has told you to restrict fluids.
- Do not stop the medicine on your own.



Common side effects

- You may pass urine more often.
- Burning or pain while passing urine.
- Frequent urges to urinate.
- Lower abdominal discomfort.
- Excessive thirst, dry mouth, dizziness, or weakness due to loss of extra fluid.
- Female genital infection – itching, redness, irritation, or unusual discharge.

Symptoms of low blood sugar (hypoglycemia)

You may feel:

- Sweating
- Shaking
- Dizziness
- Hunger
- Weakness
- Confusion
- Irritability
- Blurred vision
- Difficulty in concentrating



15–15 Rule for Low Blood Sugar

If you feel symptoms of low blood sugar (such as shakiness, sweating, dizziness, hunger or confusion):

1. **Take 15 grams of fast-acting sugar** (e.g., glucose tablets, ½ cup fruit juice or 1 tablespoon sugar).
2. **Wait 15 minutes** and check your blood sugar again (if possible).
3. **If still low, repeat 15 grams** and recheck after 15 minutes.

Once your blood sugar is above 70 mg/dL, have a small snack or meal to prevent it from dropping again.

Seek medical attention immediately if you develop:

- Severe or repeated urinary symptoms with fever or back pain (possible kidney infection).
- Severe dizziness, weakness or signs of dehydration (even if blood sugar is normal).
- Very low blood sugar symptoms that do not improve after taking sugar.
- Severe nausea, vomiting, abdominal pain, unusual tiredness, or difficulty breathing (possible diabetic ketoacidosis, even if blood sugar is not very high).



Do not stop Dapagliflozin without consulting your doctor or pharmacist.