# Empagliflozin

**Helps lower your blood sugar and may protect your heart and kidneys**

## What does it do?

- Helps your kidneys remove extra sugar from your blood through urine.
- Also helps your body remove some extra salt and water, which can reduce the workload on your heart and kidneys.
- Works best with a healthy diet and regular physical activity.



**May help protect your heart and kidneys.**

## How should I take it?

- Take exactly as prescribed by your doctor, usually once daily.
- Can be taken with or without food.
- Take at the same time every day.
- Drink enough water unless your doctor has told you to restrict fluids.
- Do not stop the medicine on your own.



## Common side effects

- You may need to pass urine more often than usual.
- Burning or pain while urinating.
- Frequent urges to urinate.
- Lower abdominal discomfort.
- These effects occur because the medicine removes extra sugar and some water through urine.

## Symptoms of low blood sugar (hypoglycemia)

You may feel:

- Sweating
- Shaking
- Dizziness
- Hunger
- Weakness
- Confusion
- Irritability
- Fast heartbeat
- Blurred vision
- Difficulty in concentrating

## 15–15 Rule for Low Blood Sugar

If you feel symptoms of low blood sugar (such as shakiness, sweating, dizziness, hunger or confusion):

1. **Take 15 grams of fast-acting sugar** (e.g., glucose tablets, ½ cup fruit juice or 1 tablespoon sugar).
2. **Wait 15 minutes** and check your blood sugar again (if possible).
3. **If still low, repeat 15 grams** and recheck after 15 minutes.

Once your blood sugar is above 70 mg/dL, have a small snack or meal to prevent it from dropping again.

## Seek medical attention immediately if you develop:

- Severe or repeated urinary symptoms (burning, lower abdominal pain, or fever).
- Severe dizziness, weakness or signs of dehydration (even if blood sugar is normal).
- Very low blood sugar symptoms that do not improve after taking sugar.



**Do not stop Empagliflozin without consulting your doctor or pharmacist.**