Glibenclamide (Glyburide)

Helps lower your blood sugar

What does it do?

- Glibenclamide helps lower your blood sugar by stimulating your pancreas to release more insulin.
- This helps your body use the sugar in your blood and lowers your blood sugar.



How should I take it?

- Take Glibenclamide with meals or as directed by your doctor.
- Take it at the same time each day.
- Follow the prescribed dose carefully.
- Do not skip meals after taking the medicine, as this can increase the risk of low blood sugar.



Common side effects

- Low blood sugar — sweating, shaking, hunger, dizziness, weakness, fast heartbeat or confusion.
- Weight gain.
- Nausea or stomach discomfort.
- Headache or dizziness.
- Rarely, skin rash or allergic reactions.



If blood sugar becomes low

If you feel shaky, sweaty, dizzy, hungry, weak or confused, your blood sugar may be low. Take 15 grams of fast-acting sugar, wait 15 minutes, and check again. If it is still low, repeat the same steps.





If you become unconscious or cannot swallow, do not give anything by mouth and **seek emergency medical help**.



Important warning

Tell your doctor if you have frequent episodes of low blood sugar, severe weakness, unusual skin reactions or yellowing of the skin or eyes.



Do not stop glibenclamide (glyburide) without consulting your doctor or pharmacist.