
Gliclazide

Helps lower your blood sugar


What does it do?

- Gliclazide helps lower your blood sugar by stimulating your pancreas to release more insulin.
- This helps your body use the sugar in your blood and keeps your blood sugar under control.



How should I take it?

- Take Gliclazide with your meals.
- Take it exactly as prescribed by your doctor.
- If you are taking a modified-release tablet, it is usually taken once daily with breakfast.
- Take it at the same time every day.
- Do not skip meals, as missing a meal can increase the risk of low blood sugar.



Common side effects

- Low blood sugar — sweating, shaking, hunger, dizziness, weakness, fast heartbeat or confusion.
- Weight gain.
- Headache or dizziness.
- Nausea or stomach discomfort.
- Rarely, skin rash or allergic reactions.



If blood sugar becomes low

If you feel shaky, sweaty, dizzy, hungry, weak or confused, your blood sugar may be low. Take 15 grams of fast-acting sugar, wait 15 minutes, and check again. If it is still low, repeat the same steps.





If you become unconscious or cannot swallow, do not give anything by mouth and **seek emergency medical help.**



Important warning

Tell your doctor if you have frequent episodes of low blood sugar, severe weakness, unusual skin reactions or yellowing of the skin or eyes.



Do not stop gliclazide without consulting your doctor or pharmacist.