
Glimepiride

Helps lower your blood sugar


What does it do?

- Glimepiride helps lower your blood sugar by making your pancreas release more insulin.
- Insulin helps your body use sugar from your blood for energy.



How should I take it?

- Usually taken once daily with breakfast or the first main meal of the day.
- Take it at approximately the same time every day.
- Take the dose exactly as prescribed by your doctor.
- Do not skip or delay your meal after taking Glimepiride, as this may increase the risk of low blood sugar.



Common side effects

- Low blood sugar — sweating, shaking, hunger, dizziness, weakness, fast heartbeat or confusion.
- Weight gain.
- Headache or dizziness.
- Nausea or stomach discomfort.
- Rarely, skin rash or yellowing of the skin/eyes.



If blood sugar becomes low

If you feel shaky, sweaty, dizzy, hungry, weak or confused, your blood sugar may be low. Take 15 grams of fast-acting sugar, wait 15 minutes, and check again. If it is still low, repeat the same steps.



1. Take 15 grams of fast-acting sugar (e.g., ½ glass of fruit juice or 3 teaspoons of sugar dissolved in water)

2. Wait 15 minutes and check your blood sugar again (if possible).

3. If still low, repeat the same steps.



If you become unconscious or cannot swallow, do not give anything by mouth and **seek emergency medical help**.



Important warning

Tell your doctor if you have frequent episodes of low blood sugar, unusual bleeding or bruising, yellowing of the skin/eyes, dark urine or severe weakness.



Do not stop glimepiride without consulting your doctor or pharmacist.