Glipizide

Helps lower your blood sugar

What does it do?

- Glipizide helps lower your blood sugar by stimulating your pancreas to release more insulin.
- This helps your body use the sugar in your blood and keeps your blood sugar under control.



How should I take it?

- Take Glipizide with your meals.
- The regular tablet is usually taken about 30 minutes before breakfast or a meal.
- The extended-release tablet is usually taken once daily with breakfast.
- Swallow extended-release tablets whole. Do not crush, chew or divide them.
- Take the medicine at the same time every day.
- Take the dose exactly as prescribed by your doctor.



Regular tablet: Take about 30 minutes before breakfast or a meal.



Extended-release tablet: Take once daily with breakfast. Swallow whole.



Take at the same time every day.



Take with meals.

Common side effects

- Low blood sugar — sweating, shaking, hunger, dizziness, weakness, fast heartbeat or confusion.
- Weight gain.
- Headache or dizziness.
- Nausea or stomach discomfort.
- Rarely, skin rash or other allergic reactions.



Low blood sugar (symptoms as above)



Weight gain



Headache or dizziness



Nausea or stomach discomfort



Skin rash or allergic reactions (rarely)

If blood sugar becomes low

If you feel shaky, sweaty, dizzy, hungry, weak or confused, your blood sugar may be low. Take 15 grams of fast-acting sugar, wait 15 minutes, and check again. If it is still low, repeat the same steps.



1. Take 15 grams of fast-acting sugar (e.g., ½ glass of fruit juice or 3 teaspoons of sugar dissolved in water)



2. Wait 15 minutes and check your blood sugar again (if possible).



3. If still low, repeat the same steps.



If you become unconscious or cannot swallow, do not give anything by mouth and **seek emergency medical help**.



Important warning

Tell your doctor if you have frequent episodes of low blood sugar, severe weakness, unusual skin reactions or yellowing of the skin or eyes.



Do not stop glipizide without consulting your doctor or pharmacist.