

MA 214 - Introduction to Numerical Analysis

Vishal Neeli

January 17, 2021

1 Interpolation Theory

1.1 Introduction

- Given finite set of points, reconstructing the original curve is interpolation.
- There will be obviously infinitely many curve.
- Interpolation problem

Given $n+1$ real distinct points: x_0, x_1, \dots, x_n and real numbers: y_0, y_1, \dots, y_n

Find a function $f : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$f(x_i) = y_i \quad \text{for } i = 0, 1, \dots, n$$

Such a function is called **interpolant** and points x_i are called **interpolation points**.

We attempt to rebuild original function using polynomial functions. This is called polynomial interpolation and function is polynomial interpolant.

- A polynomial is function of the form

$$p(x) = a_0 + a_1x + \dots + a_nx^n$$

- \mathbb{P}_n is the set of polynomials consisting of all polynomials of degree $\leq n$

1.2 Polynomial Interpolation

Theorem (Joseph-Louis Lagrange Theorem). *Given $n+1$ data points with unique x_i s, then there exists a unique polynomial $p_n \in \mathbb{P}_n$ such that*

$$p(x_i) = y_i \quad \text{for } i = 0, 1, \dots, n$$

Proof. (1) This can be shown by linear algebra. In a n degree polynomial, we substitute the points and get $n+1$ equations in $n+1$ variables (coeff) and all the rows are unique (since x_0, x_1, \dots, x_n are unique), hence in $AX = b$, $|A| \neq 0$. □

Proof. (2) Part 1: Uniqueness : If there is an interpolant, then the interpolant is unique
Let there be 2 interpolants, p_n and q_n and let $r(x) = p(x) - q(x)$,

$$r(x) = 0 \quad \text{for } i = 0, 1, \dots, n$$

This contradicts the fundamental theorem of Algebra. (A polynomial of degree n can have at most n real roots). Therefore

$$\begin{aligned} r(x) &= 0 \quad \forall x \in \mathbb{R} \\ p(x) &= q(x) \quad \forall x \in \mathbb{R} \end{aligned}$$

Part 2: Existence (construction):

Given $n+1$ data points, build $n+1$ Lagrange polynomials

$$L_k^n(x_i) = \begin{cases} 0 & \text{for } i \neq k \\ 1 & \text{for } i = k \end{cases}$$

$$L_k^n(x) = \frac{(x - x_0) \dots (x - x_{k-1})(x - x_{k+1}) \dots (x - x_n)}{(x_k - x_0) \dots (x_k - x_{k-1})(x_k - x_{k+1}) \dots (x_k - x_n)}$$

$$p(x) = \sum_{k=0}^n y_k L_k^n(x)$$

□

1.3 Closeness between functions

Given two continuous functions $f, g : [a, b] \rightarrow \mathbb{R}$, to evaluate how close the functions are consider the following

$$\max_{x \in [a, b]} |f(x) - g(x)|$$

1.4 Set of continuous Functions

$C[a, b]$ is the set of all continuous functions on $[a, b]$

$C[a, b]$ is a infinite dimensional vector space

$$f, g \in C \implies f + g \in C \text{ and } \lambda f \in C$$

We define norm on $C[a, b]$ as

$$\|f\| = \max_{x \in [a, b]} |f(x)|$$

$C^k[a, b]$ denotes the set of all functions which are continuously k-times differentiable

1.5 Polynomial Approximation and Error

Theorem (Weierstrass approximation Theorem). *Given a function $f \in C[a, b]$ and given $\epsilon > 0$, there exists a polynomial $p(x)$ such that,*

$$\|f(x) - p\| < \epsilon$$

Using Langrange's recipe to approximate

Take $n + 1$ interpolation points in the $[a, b]$ and collect the function values at all the points. We have $n + 1$ data points. Using Lagrange polynomials, find the interpolant

Theorem (Error equation). *Let $f \in C^k[a, b]$, $x_0, x_1, \dots, x_n \in [a, b]$ and $p \in \mathbb{P}_n$ be the interpolant using these points, then for all x , there exists a $\zeta = \zeta(x) \in (a, b)$ such that*

$$f(x) - p(x) = \frac{1}{(n+1)!} f^{(n+1)}(\zeta) \prod_{k=0}^n (x - x_k)$$

Note: Here ζ is dependent on the x , i.e, for every x you choose, ζ generally changes.

Proof. Consider the function,

$$\psi(t) = (f(t) - p(t)) \prod_{k=0}^n (t - x_k) - (f(x) - p(x)) \prod_{k=0}^n (x - x_k)$$

This $n + 2$ roots ($n+1$ data points and x), applying Rolle theorem's gives us that $\psi^{(1)}(t)$ has at least $n+1$ roots. Applying like this repeatedly on its derivatives, we get that $\psi^{(n+1)}$ has at least 1 root in $[a, b]$. Assuming the root to be ζ . We have,

$$f(x) - p(x) = \frac{1}{(n+1)!} f^{(n+1)}(\zeta) \prod_{k=0}^n (x - x_k)$$

□

Approximating the error:

Taking norm on both the sides of error equation, we have,

$$\max_{x \in [a, b]} |f(x) - p(x)| = \frac{1}{(n+1)!} \|f^{(n+1)}(\zeta)\| \prod_{k=0}^n (x - x_k) \quad (1)$$

$$\max_{x \in [a, b]} |f(x) - p(x)| \leq \frac{1}{(n+1)!} \|f^{(n+1)}\| \max_{x \in [a, b]} \prod_{k=0}^n (x - x_k) \quad (2)$$

Chebyshev interpolation points:

$$x_k = \frac{a+b}{2} + \frac{b-a}{2} \cos\left(\frac{j\pi}{n}\right)$$

These points minimise $\max_{x \in [a, b]} \prod_{k=0}^n (x - x_k)$ and therefore preferred over equally spaced points on real line. These points can be visualised as projections of equally spaced points on the arc of the semicircle with $\frac{a+b}{2}$ as center and $\frac{b-a}{2}$ as radius.

1.6 Another method for calculating interpolant

This is similar to the linear algebra method (given as proof(1) to Joseph-Louis Lagrange Theorem) for finding the interpolant.

Consider the polynomial

$$p(x) = a_0 + a_1(x - x_0) + a_2(x - x_0)(x - x_1) + \dots + a_n(x - x_0)\dots(x - x_{n-1})$$

Find the coefficients a_0, a_1, \dots, a_n by substituting the data points. On substituting x_0 , we get a_0 , again on substituting x_1 and using a_0 , we get a_1 and so on.

1.7 Divided difference - recursion relation

- **Divided difference** : It is the coefficient of x_n in the interpolant $p \in \mathbb{P}_n$ and denoted by $f[x_0, x_1, \dots, x_n]$.

Using Lagrange polynomials, we have

$$p(x) = \sum_{k=0}^n f(x_k) \prod_{j=0, j \neq k}^n \frac{x - x_j}{x_k - x_j}$$

So the divided difference is

$$f[x_0, x_1, \dots, x_n] = \sum_{k=0}^n f(x_k) \prod_{j=0, j \neq k}^n \frac{1}{x_k - x_j}$$

- interpolant $p(x)$ of x_0, x_1, \dots, x_n can be calculated using interpolants $a(x)$ and $b(x)$ of x_1, x_2, \dots, x_n and x_0, x_1, \dots, x_{n-1} respectively as

$$p(x) = \frac{(x - x_0)a(x) - (x - x_n)b(x)}{x_n - x_0}$$

Note: If replace the data point x_n in $a(x)$ with x_0 , we get $b(x)$

Theorem (Divided difference recursion theorem).

$$f[x_0, x_1, \dots, x_{m+1}] = \frac{[f[x_1, x_2, \dots, x_{m+1}] - f[x_0, x_1, \dots, x_m]]}{x_{m+1} - x_0}$$

Proof. Let $p(x)$ be the interpolant for x_0, x_1, \dots, x_m and $q(x)$ be the interpolant for x_1, x_2, \dots, x_{m+1} . Then,

$$L(x) = \frac{(x - x_0)q(x) + (x_{m+1} - x)p(x)}{x_{m+1} - x_0}$$

is an interpolant. Since, interpolant is unique, considering coeff of x_m we have,

$$f[x_0, x_1, \dots, x_m] = \frac{f[x_1, x_2, \dots, x_{m+1}] - f[x_0, x_1, \dots, x_m]}{x_{m+1} - x_0}$$

□

Theorem (Interpolant using divided differences). Suppose x_0, x_1, \dots, x_n be the data points. Then interpolant $p \in \mathbb{P}_n$ is

$$p(x) = f[x_0] + f[x_0, x_1](x - x_0) + \dots + f[x_0, x_1, \dots, x_n] \prod_{j=0}^{n-1} (x - x_j)$$

Proof. We prove this by induction. Base case $n = 0$ is trivially satisfied. Assume that this is satisfied for p_k ,

$$p_k(x) = f[x_0] + f[x_0, x_1](x - x_0) + \dots + f[x_0, x_1, \dots, x_k] \prod_{j=0}^{k-1} (x - x_j)$$

Consider the polynomial $p_{k+1}(x) - p_k(x) \in \mathbb{P}_{k+1}$ which has x_0, x_1, \dots, x_k as roots. Hence,

$$p_{k+1}(x) - p_k(x) = c \prod_{j=0}^k (x - x_j)$$

Comparing leading coefficient on both sides, we have $c = f[x_0, x_1, \dots, x_k]$. Hence,

$$p_{k+1}(x) = f[x_0] + f[x_0, x_1](x - x_0) + \dots + f[x_0, x_1, \dots, x_{k+1}] \prod_{j=0}^k (x - x_j)$$

By PMI,

$$p(x) = f[x_0] + f[x_0, x_1](x - x_0) + \dots + f[x_0, x_1, \dots, x_n] \prod_{j=0}^{n-1} (x - x_j)$$

□