



Respiration in Plants

12 Food, Oxidation and the ATP Currency

All living organisms need energy for carrying out daily life activities, be it **absorption, transport, movement, reproduction** or even **breathing**. All the energy required for 'life' processes is obtained by **oxidation of some macromolecules that we call 'food'**.

- **Green plants and cyanobacteria** can **prepare their own food**: by **photosynthesis** they **trap light energy** and convert it into **chemical energy** stored in the **bonds of carbohydrates** like **glucose, sucrose and starch**.
- In green plants too, **not all cells, tissues and organs photosynthesise** - **only cells containing chloroplasts**, that are **most often located in the superficial layers**, carry out photosynthesis.
- Even in green plants **all other organs, tissues and cells that are non-green need food for oxidation**. Hence, **food has to be translocated to all non-green parts**.
- **Animals are heterotrophic**, i.e., they obtain food from plants **directly (herbivores)** or **indirectly (carnivores)**.
- **Saprophytes like fungi** are dependent on **dead and decaying matter**.

So ultimately all the food that is respired for life processes comes from photosynthesis. Photosynthesis takes place within the chloroplasts (in the eukaryotes), whereas the **breakdown of complex molecules to yield energy** takes place in the cytoplasm and in the mitochondria (also only in eukaryotes).

- **Respiration**: the **breaking of the C-C bonds of complex compounds through oxidation within the cells**, leading to **release of considerable amount of energy**.
- **Respiratory substrates**: the **compounds that are oxidised** during this process. Usually carbohydrates are **oxidised** to release energy, but **proteins, fats and even organic acids can be used** as respiratory substances in some plants, under certain conditions.

During oxidation within a cell, **all the energy contained in respiratory substrates is not released free into the cell, or in a single step**. It is released in a **series of slow step-wise reactions controlled by enzymes**, and it is **trapped as chemical energy in the form of ATP**. The energy released by oxidation in respiration is **not (or rather cannot be) used directly** but is used to **synthesise ATP**, which is **broken down whenever (and wherever) energy needs to be utilised**. Hence **ATP acts as the energy currency of the cell**.

① **[NOTE]** Respiration does not only pay the energy bill. The **carbon skeleton produced during respiration is used as precursors for biosynthesis of other molecules in the cell** - the point SS12.6 returns to as the **amphibolic nature of the pathway**.

12.1 Do Plants Breathe?

The answer to this question is **not quite so direct**. Yes, plants require O_2 for respiration to occur and they **also give out CO_2** . But **plants, unlike animals, have no specialised organs for gaseous exchange** - they have **stomata and lenticels** for this purpose.

12.1a Why No Respiratory System Is Needed

Reason	What NCERT states
Each part is on its own	Each plant part takes care of its own gas-exchange needs. There is very little transport of gases from one plant part to another.
The demand is low	Plants do not present great demands for gas exchange. Roots, stems and leaves respire at rates far lower than animals do.
Peak demand is self-served	Only during photosynthesis are large volumes of gases exchanged, and each leaf is well adapted to take care of its own needs during these periods. When cells photosynthesise, availability of O_2 is not a problem in these cells since O_2 is released within the cell.
Diffusion distances are short	The distance that gases must diffuse even in large, bulky plants is not great. Each living cell in a plant is located quite close to the surface of the plant. In stems, the 'living' cells are organised in thin layers inside and beneath the bark; they also have openings called lenticels. The cells in the interior are dead and provide only mechanical support.

Reason	What NCERT states
Air reaches the cells	Most cells of a plant have at least a part of their surface in contact with air. This is also facilitated by the loose packing of parenchyma cells in leaves, stems and roots, which provide an interconnected network of air spaces.

The complete combustion of glucose, which produces CO_2 and H_2O as end products, yields energy most of which is given out as heat:



The key is to oxidise glucose not in one step but in several small steps, enabling some steps to be just large enough such that the energy released can be coupled to ATP synthesis. During the process of respiration, oxygen is utilised, and carbon dioxide, water and energy are released as products.

12.1b The Anaerobic Legacy

There are sufficient reasons to believe that the first cells on this planet lived in an atmosphere that lacked oxygen. Some of these organisms are facultative anaerobes, while in others the requirement for anaerobic condition is obligate. In any case, all living organisms retain the enzymatic machinery to partially oxidise glucose without the help of oxygen.

- **Glycolysis:** this breakdown of glucose to pyruvic acid.

12.2 Glycolysis

The term **glycolysis** has originated from the Greek words, **glycos** for sugar, and **lysis** for splitting. The scheme of glycolysis was given by Gustav Embden, Otto Meyerhof, and J. Parnas, and is often referred to as the EMP pathway.

- In anaerobic organisms, it is the only process in respiration.
- Glycolysis occurs in the cytoplasm of the cell and is present in all living organisms.
- In this process, glucose undergoes partial oxidation to form two molecules of pyruvic acid.

In plants, this glucose is derived from sucrose, which is the end product of photosynthesis, or from storage carbohydrates. Sucrose is converted into glucose and fructose by the enzyme, invertase, and these two monosaccharides readily enter the glycolytic pathway. Glucose and fructose are phosphorylated to give rise to glucose-6-phosphate by the activity of the enzyme hexokinase. This phosphorylated form of glucose then isomerises to produce fructose-6-phosphate. Subsequent steps of metabolism of glucose and fructose are same.

12.2a The Ten Reactions, Step by Step

In glycolysis, a chain of ten reactions, under the control of different enzymes, takes place to produce pyruvate from glucose.

- 1 Glucose (6C) is phosphorylated to glucose-6-phosphate (6C) by hexokinase - ATP is utilised here (first of the two ATP-spending steps) and ADP is released.
- 2 Glucose-6-phosphate isomerises to fructose-6-phosphate (6C).
- 3 Fructose 6-phosphate to fructose 1,6-bisphosphate (6C) - the second step at which ATP is utilised.
- 4 The fructose 1,6-bisphosphate is split into the two triose phosphates: dihydroxyacetone phosphate (3C) and 3-phosphoglyceraldehyde, PGAL - also written glyceraldehyde-3-phosphate (3C).
- 5 PGAL to 1,3-bisphosphoglycerate (BPGA): this is the one step where $\text{NADH} + \text{H}^+$ is formed from NAD^+ . Two redox-equivalents are removed (in the form of two hydrogen atoms) from PGAL and transferred to a molecule of NAD^+ ; PGAL is oxidised and with inorganic phosphate to get converted into BPGA. As a triose bisphosphate, BPGA is the 1,3-bisphosphoglyceric acid (3C) of the chart.
- 6 BPGA to 3-phosphoglyceric acid (PGA) - the triose phosphate 3-phosphoglyceric acid (3C). This is also an energy yielding process; this energy is trapped by the formation of ATP.
- 7 PGA to 2-phosphoglycerate.

8

2-phosphoglycerate loses H_2O to give phosphoenolpyruvate (PEP).

9

Another ATP is synthesised during the conversion of PEP to pyruvic acid.

10

Pyruvic acid (3C) is then the key product of glycolysis.

What is the metabolic fate of pyruvate? This depends on the cellular need. There are three major ways in which different cells handle pyruvic acid produced by glycolysis.

Fate of pyruvic acid	Conditions	Outcome
Lactic acid fermentation	Anaerobic	Incomplete oxidation; pyruvic acid is reduced to lactic acid
Alcoholic fermentation	Anaerobic	Incomplete oxidation; pyruvic acid is converted to CO_2 and ethanol
Aerobic respiration	Requires O_2 supply	Complete oxidation of glucose to CO_2 and H_2O through Krebs' cycle

Fermentation takes place under anaerobic conditions in many prokaryotes and unicellular eukaryotes. For the complete oxidation of glucose to CO_2 and H_2O , however, organisms adopt Krebs' cycle which is also called as aerobic respiration. This requires O_2 supply.




Fig. 12.1 - Steps of glycolysis. The chart runs from **Glucose (6C)** down through **Glucose-6-phosphate (6C)** and **Fructose-6-phosphate (6C)** - each of the first two steps spending one ATP and releasing ADP - to **Fructose 1,6-bisphosphate (6C)**, which splits into the **Triose phosphate dihydroxyacetone phosphate (3C)** and the **Triose phosphate glyceraldehyde-3-phosphate (3C)**. The latter, with NAD^+ reduced to $\text{NADH} + \text{H}^+$, becomes the **Triose bisphosphate 1,3-bisphosphoglyceric acid (3C)**, then the **Triose phosphate 3-phosphoglyceric acid (3C)**, then **2-phosphoglycerate**, which loses H_2O to give **phosphoenolpyruvate** and finally **Pyruvic acid (3C)**.




Fig. 12.2 - Major pathways of anaerobic respiration. **Glucose** passes through **Glyceraldehyde 3-phosphate**, **3-phosphoglyceric acid** and **phosphoenol pyruvic acid** to **Pyruvic acid**, which then branches either to **Lactic acid** or to **Ethanol plus CO₂**. **NAD⁺** and **NADH + H⁺** are shown being interconverted along the pathway, since the reducing power taken out of the sugar is handed back at the branch.

12.3 Fermentation

In fermentation, say by yeast, the **incomplete oxidation of glucose** is achieved under **anaerobic conditions** by sets of reactions where **pyruvic acid** is converted to **CO₂** and **ethanol**. The enzymes, **pyruvic acid decarboxylase** and **alcohol dehydrogenase** catalyse these reactions.

- Other organisms like some bacteria produce lactic acid from pyruvic acid.
- In animal cells also, like muscles during exercise, when oxygen is inadequate for cellular respiration pyruvic acid is reduced to lactic acid by lactate dehydrogenase.
- The reducing agent is **NADH + H⁺**, which is **reoxidised to NAD⁺** in both the processes.

12.3a Why Fermentation Is a Poor Deal

- Less than seven per cent of the energy in glucose is released, and not all of it is trapped as high energy bonds of ATP.
- Also, the processes are hazardous - either acid or alcohol is produced.
- Yeasts poison themselves to death when the concentration of alcohol reaches about 13 per cent.

① [NOTE] Where fermentation happens: under anaerobic conditions in many prokaryotes, unicellular eukaryotes and in germinating seeds.

12.3b The Aerobic Alternative

- **Aerobic respiration:** the process that leads to a **complete oxidation of organic substances** in the presence of oxygen, and releases **CO₂**, water and a large amount of energy present in the substrate.

This type of respiration is most common in higher organisms. In eukaryotes these steps take place within the mitochondria and this requires **O₂**.

12.4 Aerobic Respiration

For aerobic respiration to take place **within the mitochondria**, the **final product of glycolysis, pyruvate** is transported from the cytoplasm into the mitochondria. Two crucial events then follow:



The complete oxidation of pyruvate by the stepwise removal of all the hydrogen atoms, leaving three molecules of **CO₂**. This first process takes place in the matrix of the mitochondria.

2

The passing on of the electrons removed as part of the hydrogen atoms to molecular O_2 with simultaneous synthesis of ATP. This second process is located on the inner membrane of the mitochondria.

Pyruvate, which is formed by the glycolytic catabolism of carbohydrates in the cytosol, after it enters mitochondrial matrix undergoes oxidative decarboxylation by a complex set of reactions catalysed by pyruvic dehydrogenase. These reactions require the participation of several coenzymes, including NAD^+ and Coenzyme A:

Pyruvic acid + CoA + NAD^+ (with Mg^{2+} and pyruvate dehydrogenase) yields Acetyl CoA + CO_2 + $NADH + H^+$

During this process, two molecules of $NADH$ are produced from the metabolism of two molecules of pyruvic acid (produced from one glucose molecule during glycolysis). The acetyl CoA then enters a cyclic pathway, tricarboxylic acid cycle, more commonly called as Krebs' cycle after the scientist Hans Krebs who first elucidated it.

12.4.1

Tricarboxylic Acid Cycle

▲

(cycle - last step feeds back to step 1)

1

The TCA cycle starts with the condensation of acetyl group with oxaloacetic acid (OAA) and water to yield citric acid. The reaction is catalysed by the enzyme citrate synthase and a molecule of CoA is released.

2

Citrate is then isomerised to isocitrate.

3

It is followed by two successive steps of decarboxylation, leading to the formation of α -ketoglutaric acid and then succinyl-CoA.

4

During the conversion of succinyl-CoA to succinic acid a molecule of GTP is synthesised. This is a substrate level phosphorylation. In a coupled reaction GTP is converted to GDP with the simultaneous synthesis of ATP from ADP.

5

In the remaining steps of citric acid cycle, succinyl-CoA is oxidised to OAA allowing the cycle to continue.

- There are three points in the cycle where NAD^+ is reduced to $NADH + H^+$ and one point where FAD^+ is reduced to $FADH_2$.
- The continued oxidation of acetyl CoA via the TCA cycle requires the continued replenishment of oxaloacetic acid, the first member of the cycle.
- In addition it also requires regeneration of NAD^+ and FAD^+ from $NADH$ and $FADH_2$ respectively.

The summary equation for this stage of respiration, in the mitochondrial matrix, may thus be written as:

Pyruvic acid + $4NAD^+$ + FAD^+ + $2H_2O$ + $ADP + P_i$ yields $3CO_2 + 4NADH + 4H^+ + FADH_2 + ATP$

By the end of the cycle, glucose has been broken down to release CO_2 and eight molecules of $NADH + H^+$ and two of $FADH_2$ have been synthesised, besides just two molecules of ATP in TCA cycle.




Fig. 12.3 - The Citric acid cycle. **Pyruvate (3C)** loses CO₂ and, with **CoA** and **NAD⁺** reduced to **NADH + H⁺**, gives **Acetyl coenzyme A (2C)**, which condenses with **Oxaloacetic acid (4C)** to form **Citric acid (6C)**. Around the ring, decarboxylations give **alpha-ketoglutaric acid (5C)** and then, via **Succinic acid (4C)** - the step in which **GDP** becomes **GTP** - and **Malic acid (4C)**, the cycle returns to oxaloacetic acid. **FAD⁺** to **FADH₂** marks the single flavin-linked oxidation.

12.4.2 Electron Transport System (ETS) and Oxidative Phosphorylation

The following steps in the respiratory process are to release and utilise the energy stored in **NADH + H⁺** and **FADH₂**. This is accomplished when they are oxidised through the electron transport system and the electrons are passed on to O₂ resulting in the formation of H₂O.

● **Electron transport system (ETS):** the metabolic pathway through which the electron passes from one carrier to another. It is present in the inner mitochondrial membrane.

- 1 **Electrons from NADH produced in the mitochondrial matrix during citric acid cycle are oxidised by an NADH dehydrogenase (complex I), and electrons are then transferred to ubiquinone located within the inner membrane.**
- 2 **Ubiquinone also receives reducing equivalents via FADH₂ (complex II) that is generated during oxidation of succinate in the citric acid cycle.**
- 3 **The reduced ubiquinone (ubiquinol) is then oxidised with the transfer of electrons to cytochrome c via cytochrome bc₁ complex (complex III).**
- 4 **Cytochrome c is a small protein attached to the outer surface of the inner membrane and acts as a mobile carrier for transfer of electrons between complex III and IV.**
- 5 **Complex IV refers to cytochrome c oxidase complex containing cytochromes a and a₃, and two copper centres.**
- 6 **When the electrons pass from one carrier to another via complex I to IV in the electron transport chain, they are coupled to ATP synthase (complex V) for the production of ATP from ADP and inorganic phosphate.**

The number of ATP molecules synthesised depends on the nature of the electron donor.

Electron donor oxidised	ATP produced
One molecule of NADH	3 molecules of ATP
One molecule of FADH ₂	2 molecules of ATP

Although the **aerobic process of respiration** takes place only in the presence of **oxygen**, the **role of oxygen** is limited to the terminal stage of the process. Yet the **presence of oxygen** is vital, since it drives the whole

process by removing hydrogen from the system: oxygen acts as the final hydrogen acceptor.

- **Oxidative phosphorylation:** unlike photophosphorylation where it is the light energy that is utilised for the production of proton gradient required for phosphorylation, in respiration it is the energy of oxidation-reduction utilised for the same process. It is for this reason that the process is called oxidative phosphorylation.

12.4.2a ATP Synthase - the F_1/F_0 Machine

The energy released during the electron transport system is utilised in synthesising ATP with the help of ATP synthase (complex V). This complex consists of two major components, F_1 and F_0 .

- The F_1 headpiece is a peripheral membrane protein complex and contains the site for synthesis of ATP from ADP and inorganic phosphate.
- F_0 is an integral membrane protein complex that forms the channel through which protons cross the inner membrane.
- The passage of protons through the channel is coupled to the catalytic site of the F_1 component for the production of ATP.
- For each ATP produced, $4H^+$ pass through F_0 from the intermembrane space to the matrix down the electrochemical proton gradient.




Fig. 12.4 - Electron Transport System (ETS). The carriers are drawn in the **inner mitochondrial membrane** with the **matrix** below. **NADH** feeds **complex I** and **succinate** feeds **FADH₂** at **complex II**; both hand electrons to **ubiquinone**, which passes them through **complex III** to **cytochrome c** and on to **complex IV**, where **O₂** is reduced to **H₂O**. **Complex V** is the **ATP synthase** that makes ATP from **ADP** and **inorganic phosphate**.




Fig. 12.5 - Diagrammatic presentation of ATP synthesis in mitochondria. The F_1 headpiece projects into the **matrix** and carries the site where **ADP** and **inorganic phosphate** are joined into **ATP**; F_0 is the channel through the **inner mitochondrial membrane** through which $4H^+$ return from the **intermembrane space** down the **electrochemical proton gradient** for every ATP made.

It is possible to make calculations of the net gain of ATP for every glucose molecule oxidised; but in reality this can remain only a theoretical exercise. The calculations can be made only on certain assumptions:

- There is a sequential, orderly pathway functioning, with one substrate forming the next and with glycolysis, TCA cycle and ETS pathway following one after another.
- The NADH synthesised in glycolysis is transferred into the mitochondria and undergoes oxidative phosphorylation.
- None of the intermediates in the pathway are utilised to synthesise any other compound.
- Only glucose is being respired - no other alternative substrates are entering in the pathway at any of the intermediary stages.

① [NOTE] None of these assumptions is strictly valid in a living system: all pathways work simultaneously and do not take place one after another; substrates enter the pathways and are withdrawn from it as and when necessary; ATP is utilised as and when needed; enzymatic rates are controlled by multiple means.

With the assumptions above, there can be a net gain of 38 ATP molecules during aerobic respiration of one molecule of glucose.

Point of difference	Fermentation	Aerobic respiration
Extent of breakdown	Accounts for only a partial breakdown of glucose	Glucose is completely degraded to CO ₂ and H ₂ O
Net ATP gain	Net gain of only two molecules of ATP for each molecule of glucose degraded to pyruvic acid	Many more molecules of ATP are generated under aerobic conditions
Oxidation of NADH to NAD ⁺	Rather slow	Very vigorous

Glucose is the favoured substrate for respiration. All carbohydrates are usually first converted into glucose before they are used for respiration. Other substrates can also be respired, as has been mentioned earlier, but then they do not enter the respiratory pathway at the first step.

Substrate	Broken down first into	Enters the respiratory pathway as
Fats	Glycerol and fatty acids	Fatty acids would first be degraded to acetyl CoA and enter the pathway; glycerol would enter the pathway after being converted to PGAL
Proteins	Individual amino acids, by proteases	The amino acids, after deamination, depending on their structure would enter the pathway at some stage within the Krebs' cycle or even as pyruvate or acetyl CoA

Since respiration involves breakdown of substrates, the respiratory process has traditionally been considered a catabolic process and the respiratory pathway as a catabolic pathway. But the same intermediates are also drawn off for synthesis: when the organism needs to synthesise fatty acids, acetyl CoA would be withdrawn from the respiratory pathway for it.

- Catabolism and anabolism: breaking down processes within the living organism is catabolism, and synthesis is anabolism.
- Amphibolic pathway: because the respiratory pathway is involved in both anabolism and catabolism, it would hence be better to consider the respiratory pathway as an amphibolic pathway rather than as a catabolic one.




Fig. 12.6 - Interrelationship among metabolic pathways showing respiration mediated breakdown of different organic molecules to CO_2 and H_2O . **Fats**, **Carbohydrates** and **Proteins** enter from the top as **fatty acids and glycerol**, **simple sugars e.g. glucose**, and **amino acids**. The sugar arm runs **glucose 6-phosphate** to **fructose 1,6 bisphosphate** to the trioses **dihydroxy acetone phosphate** and **glyceraldehyde 3-phosphate** and on to **pyruvic acid**; every arm converges on **acetyl CoA** and the **Krebs cycle**, with arrows drawn both ways to show that intermediates are withdrawn for synthesis as well as oxidised.

12.7

Respiratory Quotient

Let us now look at another aspect of respiration.

- **Respiratory quotient (RQ) or respiratory ratio:** the ratio of the volume of CO_2 evolved to the volume of O_2 consumed in respiration.

$$\text{RQ} = \text{volume of } \text{CO}_2 \text{ evolved} / \text{volume of } \text{O}_2 \text{ consumed}$$

The respiratory quotient depends upon the type of respiratory substrate used during respiration.

Respiratory substrate	RQ	Why
Carbohydrates (completely oxidised)	1	Equal amounts of CO_2 and O_2 are evolved and consumed, respectively: $\text{RQ} = 6\text{CO}_2 / 6\text{O}_2 = 1.0$
Fats	Less than 1	For tripalmitin: $2(\text{C}_{51}\text{H}_{98}\text{O}_6) + 145\text{O}_2$ yields $102\text{CO}_2 + 98\text{H}_2\text{O} + \text{energy}$, so $\text{RQ} = 102\text{CO}_2 / 145\text{O}_2 = 0.7$
Proteins	About 0.9	When proteins are respiratory substrates the ratio would be about 0.9.

① **[NOTE]** These clean values are *textbook cases*: in living organisms, respiratory substrates are often more than one; pure proteins or fats are never used as respiratory substrates.

☆ **[MEMORY AID - not in NCERT]** RQ falls in the alphabetical order of the substrates - Carbohydrate 1.0, Protein 0.9, Fat 0.7 is not alphabetical, but $\text{C} > \text{P} > \text{F}$ is: the more oxygen a substrate still needs, the lower its RQ, and fat is the most hydrogen-rich of the three.

Recap

Quick Recap

- Plants unlike animals have no special systems for breathing or gaseous exchange. Stomata and lenticels allow gaseous exchange by diffusion, and almost all living cells in a plant have their surfaces exposed to air.
- The breaking of C-C bonds of complex organic molecules by oxidation leading to the release of a lot of energy is called cellular respiration. Glucose is the favoured substrate for respiration; fats and proteins

can also be broken down to yield energy.

- The initial stage of cellular respiration takes place in the cytoplasm. Each glucose molecule is broken through a series of enzyme catalysed reactions into two molecules of pyruvic acid; this process is called glycolysis.
- The fate of the pyruvate depends on the availability of oxygen and the organism. Under anaerobic conditions either lactic acid fermentation or alcohol fermentation occurs, and fermentation takes place in many prokaryotes, unicellular eukaryotes and in germinating seeds - releasing less than seven per cent of the energy in glucose for a net gain of two ATP.
- In eukaryotic organisms aerobic respiration occurs in the presence of oxygen. Pyruvic acid is transported into the mitochondria where it is converted into acetyl CoA with the release of CO_2 . Acetyl CoA then enters the tricarboxylic acid pathway or Krebs' cycle operating in the matrix of the mitochondria.
- $\text{NADH} + \text{H}^+$ and FADH_2 are generated in the Krebs' cycle. The energy in these molecules as well as that in the $\text{NADH} + \text{H}^+$ synthesised during glycolysis are used to synthesise ATP.
- This is accomplished through a system of electron carriers called electron transport system (ETS) located on the inner membrane of the mitochondria. The electrons, as they move through the system, release enough energy that are trapped to synthesise ATP; this is called oxidative phosphorylation. In this process O_2 is the ultimate acceptor of electrons and it gets reduced to water. Net theoretical yield: 38 ATP per glucose.
- The respiratory pathway is an amphibolic pathway as it involves both anabolism and catabolism. The respiratory quotient depends upon the type of respiratory substance used during respiration - 1 for carbohydrates, 0.7 for fats (tripalmitin) and about 0.9 for proteins.