

Gradient Descent for Multiple Variables

LR with Multiple Variables

- Linear regression with multiple variables is also known as "multivariate linear regression".
- We now introduce notation for equations where we can have any number of input variables.

Multiple features (variables).

Size (feet ²)	Price (\$1000)
x	y
2104	460
1416	232
1534	315
852	178
...	...

$$h_{\theta}(x) = \theta_0 + \theta_1 x$$

Windows'u Etkinleştir
Windows'u etkinleştirmek için Ayarlar'a gidin.

Multiple features (variables).

Size (feet ²)	Number of bedrooms	Number of floors	Age of home (years)	Price (\$1000)
2104	5	1	45	460
1416	3	2	40	232
1534	3	2	30	315
852	2	1	36	178
...

Windows'u Etkinleştir
Windows'u etkinleştirmek için Ayarlar'a gidin.

Multiple features (variables).

Size (feet ²) x_1	Number of bedrooms x_2	Number of floors x_3	Age of home (years) x_4	Price (\$1000) y
2104	5	1	45	460
1416	3	2	40	232
1534	3	2	30	315
852	2	1	36	178
...

Windows'u Etkinleştir
Windows'u etkinleştirmek için Ayarlar'a gidin.

Multiple features (variables).

Size (feet ²)	Number of bedrooms	Number of floors	Age of home (years)	Price (\$1000)
x_1	x_2	x_3	x_4	y
2104	5	1	45	460
1416	3	2	40	232
1534	3	2	30	315
852	2	1	36	178
...

Notation:

→ n = number of features $n = 4$

$x^{(i)}$ = input (features) of i^{th} training example.

$x_j^{(i)}$ = value of feature j in i^{th} training example.

Windows'u Etkinleştir
Windows'u etkinleştirmek için Ayarlar'a gidin.

Multiple features (variables).

Size (feet ²)	Number of bedrooms	Number of floors	Age of home (years)	Price (\$1000)	
x_1	x_2	x_3	x_4	y	
2104	5	1	45	460	} $m = 47$
1416	3	2	40	232	
1534	3	2	30	315	
852	2	1	36	178	
...	

Notation:

→ n = number of features $n = 4$

$x^{(i)}$ = input (features) of i^{th} training example.

$x_j^{(i)}$ = value of feature j in i^{th} training example.

Windows'u Etkinleştir
Windows'u etkinleştirmek için Ayarlar'a gidin.

Multiple features (variables).

Size (feet ²)	Number of bedrooms	Number of floors	Age of home (years)	Price (\$1000)
x_1	x_2	x_3	x_4	y
2104	5	1	45	460
1416	3	2	40	232
1534	3	2	30	315
852	2	1	36	178
...

$$m = 47$$

$$x^{(2)} = \begin{bmatrix} 1416 \\ 3 \\ 2 \\ 40 \end{bmatrix}$$

Notation:

→ n = number of features

$$n = 4$$

→ $x^{(i)}$ = input (features) of i^{th} training example.

$x_j^{(i)}$ = value of feature j in i^{th} training example.

Windows'u Etkinleştir
Windows'u etkinleştirmek için Ayarlar'a gidin.

Multiple features (variables).

Size (feet ²)	Number of bedrooms	Number of floors	Age of home (years)	Price (\$1000)
x_1	x_2	x_3	x_4	y
2104	5	1	45	460
1416	3	2	40	232
1534	3	2	30	315
852	2	1	36	178
...

Notation:

- n = number of features
- $x^{(i)}$ = input (features) of i^{th} training example.
- $x_j^{(i)}$ = value of feature j in i^{th} training example.

$n = 4$

$m = 47$

$$x^{(2)} = \begin{bmatrix} 1416 \\ 3 \\ 2 \\ 40 \end{bmatrix}$$

$$x_3^{(2)} = 2$$

Windows'u Etkinleştir
Windows'u etkinleştirmek için Ayarlar'a gidin.


Hypothesis:

Previously: $h_{\theta}(x) = \theta_0 + \theta_1 x$

Windows'u Etkinleştir
Windows'u etkinleştirmek için Ayarlar'a gidin.


Hypothesis:

Previously: $h_{\theta}(x) = \theta_0 + \theta_1 x$

A handwritten red 'X' is drawn over the entire equation $h_{\theta}(x) = \theta_0 + \theta_1 x$. Below the equation, a blue horizontal line is drawn, ending with a blue arrow pointing to the right.

Windows'u Etkinleştir
Windows'u etkinleştirmek için Ayarlar'a gidin.


Hypothesis:

Previously: $\underline{h_{\theta}(x) = \theta_0 + \theta_1 x}$ 

$$h_{\Theta}(x) = \Theta_0 + \Theta_1 x_1 + \Theta_2 x_2 + \Theta_3 x_3 + \Theta_4 x_4$$

Windows'u Etkinleştir
Windows'u etkinleştirmek için Ayarlar'a gidin.

Hypothesis:

Previously: $\underline{h_{\theta}(x) = \theta_0 + \theta_1 x}$ 

$$h_{\theta}(x) = \theta_0 + \theta_1 x_1 + \theta_2 x_2 + \theta_3 x_3 + \theta_4 x_4$$

$$\text{e.g. } h_{\theta}(x) = 80 + 0.1x_1 + 0.01x_2 + 3x_3 - 2x_4$$

$$\rightarrow h_{\theta}(x) = \theta_0 + \theta_1 x_1 + \theta_2 x_2 + \cdots + \theta_n x_n$$

Windows'u Etkinleştir
Windows'u etkinleştirmek için Ayarlar'a gidin.

$$\rightarrow h_{\theta}(x) = \theta_0 + \theta_1 x_1 + \theta_2 x_2 + \cdots + \theta_n x_n$$

For convenience of notation, define $x_0 = 1$.

$$\rightarrow h_{\theta}(x) = \theta_0 + \theta_1 \underline{x_1} + \theta_2 \underline{x_2} + \dots + \theta_n \underline{x_n}$$

For convenience of notation, define $x_0 = 1$. $(x_0^{(i)} = 1)$

$$x = \begin{bmatrix} x_0 \\ x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \in \mathbb{R}^{n+1}$$

$$\rightarrow h_{\theta}(x) = \theta_0 + \theta_1 x_1 + \theta_2 x_2 + \dots + \theta_n x_n$$

For convenience of notation, define $x_0 = 1$.

$$(x_0^{(i)} = 1)$$

$$x = \begin{bmatrix} x_0 \\ x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \in \mathbb{R}^{n+1}$$

$$\theta = \begin{bmatrix} \theta_0 \\ \theta_1 \\ \theta_2 \\ \vdots \\ \theta_n \end{bmatrix}$$

$$\rightarrow h_{\theta}(x) = \underline{\theta_0} + \underline{\theta_1} \underline{x_1} + \underline{\theta_2} \underline{x_2} + \dots + \underline{\theta_n} \underline{x_n}$$

For convenience of notation, define $x_0 = 1$.

$$(x_0^{(i)} = 1)$$

$$x = \begin{bmatrix} x_0 \\ x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \in \mathbb{R}^{n+1}$$

$$\theta = \begin{bmatrix} \theta_0 \\ \theta_1 \\ \theta_2 \\ \vdots \\ \theta_n \end{bmatrix} \in \mathbb{R}^{n+1}$$

$$h_{\theta}(x) = \theta_0 x_0 + \theta_1 x_1 + \dots + \theta_n x_n$$

$$\rightarrow h_{\theta}(x) = \theta_0 + \theta_1 x_1 + \theta_2 x_2 + \dots + \theta_n x_n$$

For convenience of notation, define $x_0 = 1$. $(x_0^{(i)} = 1)$

$$x = \begin{bmatrix} x_0 \\ x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \in \mathbb{R}^{n+1}$$

$$\theta = \begin{bmatrix} \theta_0 \\ \theta_1 \\ \theta_2 \\ \vdots \\ \theta_n \end{bmatrix} \in \mathbb{R}^{n+1} !$$

$$h_{\theta}(x) = \theta_0 x_0 + \theta_1 x_1 + \dots + \theta_n x_n$$

$\downarrow = 1$

$$= \theta^T x.$$

$$\rightarrow h_{\theta}(x) = \theta_0 + \theta_1 x_1 + \theta_2 x_2 + \dots + \theta_n x_n$$

For convenience of notation, define $x_0 = 1$. ($x_0^{(i)} = 1$)

$$x = \begin{bmatrix} x_0 \\ x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \in \mathbb{R}^{n+1}$$

$$\theta = \begin{bmatrix} \theta_0 \\ \theta_1 \\ \theta_2 \\ \vdots \\ \theta_n \end{bmatrix} \in \mathbb{R}^{n+1}$$

$$h_{\theta}(x) = \theta_0 x_0 + \theta_1 x_1 + \dots + \theta_n x_n$$

$\downarrow = 1$

$$= \boxed{\theta^T x}$$

$$\underbrace{[\theta_0 \ \theta_1 \ \dots \ \theta_n]}_{\theta^T} \begin{bmatrix} x_0 \\ x_1 \\ \vdots \\ x_n \end{bmatrix}_x$$

$(n+1) \times 1$ matrix

$\theta^T x$

Multivariate linear regression. \leftarrow

- This is a vectorization of our hypothesis function for one training example

$$h_{\theta}(x) = [\theta_0 \quad \theta_1 \quad \dots \quad \theta_n] \begin{bmatrix} x_0 \\ x_1 \\ \vdots \\ x_n \end{bmatrix} = \theta^T x$$

Gradient Descent for MLR

Hypothesis: $h_{\theta}(x) = \theta^T x = \theta_0 x_0 + \theta_1 x_1 + \theta_2 x_2 + \dots + \theta_n x_n$

Parameters: $\theta_0, \theta_1, \dots, \theta_n$

Cost function:

$$J(\theta_0, \theta_1, \dots, \theta_n) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$

Hypothesis: $\underbrace{h_{\theta}(x) = \theta^T x = \theta_0 x_0 + \theta_1 x_1 + \theta_2 x_2 + \dots + \theta_n x_n}_{\text{green underline}}$ $\nearrow x_0 = 1$

Parameters: $\theta_0, \theta_1, \dots, \theta_n$ ⊙ $n+1$ -dimensional vector.

Cost function:

$$J(\theta_0, \theta_1, \dots, \theta_n) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$

Hypothesis: $h_{\theta}(x) = \theta^T x = \theta_0 x_0 + \theta_1 x_1 + \theta_2 x_2 + \dots + \theta_n x_n$

Parameters: $\theta_0, \theta_1, \dots, \theta_n$ \bigcirc $n+1$ -dimensional vector

Cost function:

$$J(\theta_0, \theta_1, \dots, \theta_n) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$

Hypothesis: $\underbrace{h_{\theta}(x) = \theta^T x = \theta_0 x_0 + \theta_1 x_1 + \theta_2 x_2 + \dots + \theta_n x_n}_{\text{green underline}}$ $\rightarrow x_0 = 1$

Parameters: $\theta_0, \theta_1, \dots, \theta_n$ $\underbrace{\quad}_{\text{blue underline}}$ $n+1$ -dimensional vector

Cost function:

$$\underbrace{J(\theta_0, \theta_1, \dots, \theta_n)}_{J(\theta)} = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$

Windows'u Etkinleştir
Windows'u etkinleştirmek için Ayarlar'a gidin.

1:18

Hypothesis: $\underbrace{h_{\theta}(x) = \theta^T x = \theta_0 x_0 + \theta_1 x_1 + \theta_2 x_2 + \dots + \theta_n x_n}_{\text{green underline}}$ $\rightarrow x_0 = 1$

Parameters: $\theta_0, \theta_1, \dots, \theta_n$ \bigcirc $n+1$ -dimensional vector

Cost function:

$$\underbrace{J(\theta_0, \theta_1, \dots, \theta_n)}_{J(\theta)} = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$

Gradient descent:

Repeat {

$$\theta_j := \theta_j - \alpha \frac{\partial}{\partial \theta_j} J(\theta_0, \dots, \theta_n)$$

}

(simultaneously update for every $j = 0, \dots, n$)

When there are n features, we define the cost function as

$$J(\theta) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2.$$

For linear regression, which of the following are also equivalent and correct definitions of $J(\theta)$?

☐ $J(\theta) = \frac{1}{2m} \sum_{i=1}^m (\theta^T x^{(i)} - y^{(i)})^2$

☐ $J(\theta) = \frac{1}{2m} \sum_{i=1}^m \left(\left(\sum_{j=0}^n \theta_j x_j^{(i)} \right) - y^{(i)} \right)^2$ (Inner sum starts at 0)

☐ $J(\theta) = \frac{1}{2m} \sum_{i=1}^m \left(\left(\sum_{j=1}^n \theta_j x_j^{(i)} \right) - y^{(i)} \right)^2$ (Inner sum starts at 1)

☐ $J(\theta) = \frac{1}{2m} \sum_{i=1}^m \left(\left(\sum_{j=0}^n \theta_j x_j^{(i)} \right) - \left(\sum_{j=0}^n y_j^{(i)} \right) \right)^2$

Hypothesis: $\underline{h_{\theta}(x) = \theta^T x = \theta_0 x_0 + \theta_1 x_1 + \theta_2 x_2 + \dots + \theta_n x_n}$

Parameters: $\underline{\theta_0, \theta_1, \dots, \theta_n}$ $\underline{\mathcal{O}}$ $n+1$ -dimensional vector

Cost function:

$$\underline{J(\theta_0, \theta_1, \dots, \theta_n)} = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$

$\underline{J(\theta)}$

Gradient descent:

Repeat {

$$\rightarrow \theta_j := \theta_j - \alpha \left[\frac{\partial}{\partial \theta_j} \underline{J(\theta_0, \dots, \theta_n)} \right] \underline{J(\theta)}$$

}

(simultaneously update for every $j = 0, \dots, n$)

Gradient Descent

Previously (n=1):

Repeat {

$$\rightarrow \theta_0 := \theta_0 - \alpha \underbrace{\frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})}_{\frac{\partial}{\partial \theta_0} J(\theta)}$$

$$\rightarrow \theta_1 := \theta_1 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x^{(i)}$$

(simultaneously update θ_0, θ_1)

• }

Windows'u Etkinleştir
Windows'u etkinleştirmek için Ayarlar'a gidin.

Gradient Descent

Previously (n=1):

Repeat {

→ $\theta_0 := \theta_0 - \alpha \underbrace{\frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})}_{\frac{\partial}{\partial \theta_0} J(\theta)}$

→ $\theta_1 := \theta_1 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) \underline{x^{(i)}}$
(simultaneously update θ_0, θ_1)

}

Gradient Descent

Previously (n=1):

Repeat {

$$\rightarrow \theta_0 := \theta_0 - \alpha \underbrace{\frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})}_{\frac{\partial}{\partial \theta_0} J(\theta)}$$

$$\rightarrow \theta_1 := \theta_1 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) \underbrace{x^{(i)}}_{x_1^{(i)}}$$

(simultaneously update θ_0, θ_1)

}

New algorithm (n ≥ 1):

Repeat {

$$\rightarrow \theta_j := \theta_j - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_j^{(i)}$$

(simultaneously update θ_j for $j = 0, \dots, n$)

}

Handwritten note: $\downarrow \frac{2}{2\theta_j} J(\theta)$

$$h_{\theta}(x) = [\theta_0 \quad \theta_1 \quad \dots \quad \theta_n] \begin{bmatrix} x_0 \\ x_1 \\ \vdots \\ x_n \end{bmatrix} = \theta^T x$$

Windows'u Etkinleştir
Windows'u etkinleştirmek için Ayarlar'a gidin.

Gradient Descent

Previously (n=1):

Repeat {

$$\rightarrow \theta_0 := \theta_0 - \alpha \underbrace{\frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})}_{\frac{\partial}{\partial \theta_0} J(\theta)}$$

$$\rightarrow \theta_1 := \theta_1 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_1^{(i)}$$

(simultaneously update θ_0, θ_1)

}

New algorithm (n ≥ 1):

Repeat {

$$\rightarrow \theta_j := \theta_j - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_j^{(i)}$$

(simultaneously update θ_j for $j = 0, \dots, n$)

} $\downarrow \frac{2}{2\theta_j} J(\theta)$

$$\theta_0 := \theta_0 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_0^{(i)}$$

$$\theta_1 := \theta_1 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_1^{(i)}$$

$$\theta_2 := \theta_2 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_2^{(i)}$$

...

Gradient Descent

Previously (n=1):

Repeat {

→ $\theta_0 := \theta_0 - \alpha \underbrace{\frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})}_{\frac{\partial}{\partial \theta_0} J(\theta)}$

→ $\theta_1 := \theta_1 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) \underbrace{x_1^{(i)}}_{x_1^{(i)}}$
(simultaneously update θ_0, θ_1)

}

New algorithm (n ≥ 1):

Repeat {

→ $\theta_j := \theta_j - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_j^{(i)}$
(simultaneously update θ_j for $j = 0, \dots, n$)

} $\downarrow \frac{2}{2\theta_j} J(\theta)$

→ $\theta_0 := \theta_0 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_0^{(i)}$

→ $\theta_1 := \theta_1 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_1^{(i)}$

→ $\theta_2 := \theta_2 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_2^{(i)}$

...

Gradient Descent

Previously (n=1):

Repeat {

→ $\theta_0 := \theta_0 - \alpha \underbrace{\frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})}_{\frac{\partial}{\partial \theta_0} J(\theta)}$

→ $\theta_1 := \theta_1 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) \underbrace{x_1^{(i)}}_{x_1^{(i)}}$
(simultaneously update θ_0, θ_1)

}

New algorithm (n ≥ 1):

Repeat {

→ $\theta_j := \theta_j - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_j^{(i)}$
(simultaneously update θ_j for $j = 0, \dots, n$)

} x₀⁽ⁱ⁾ = 1

→ $\theta_0 := \theta_0 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_0^{(i)}$
→ $\theta_1 := \theta_1 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_1^{(i)}$
→ $\theta_2 := \theta_2 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_2^{(i)}$

...

Gradient Descent

Previously (n=1):

Repeat {

→ $\theta_0 := \theta_0 - \alpha \underbrace{\frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})}_{\frac{\partial}{\partial \theta_0} J(\theta)}$

→ $\theta_1 := \theta_1 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) \underbrace{x_1^{(i)}}_{x_1^{(i)}}$
(simultaneously update θ_0, θ_1)

}

New algorithm (n ≥ 1):

Repeat {

→ $\theta_j := \theta_j - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_j^{(i)}$
(simultaneously update θ_j for $j = 0, \dots, n$)

→ $\theta_0 := \theta_0 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_0^{(i)}$

→ $\theta_1 := \theta_1 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_1^{(i)}$

→ $\theta_2 := \theta_2 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_2^{(i)}$

...

$\frac{\partial}{\partial \theta_j} J(\theta)$

$x_0^{(i)} = 1$

Windows'u etkinleştirmek için Ayarlar'a gidin.

Gradient Descent

Previously (n=1):

Repeat {

→ $\theta_0 := \theta_0 - \alpha \underbrace{\frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})}_{\frac{\partial}{\partial \theta_0} J(\theta)}$

→ $\theta_1 := \theta_1 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) \underbrace{x_1^{(i)}}_{x_1^{(i)}}$
(simultaneously update θ_0, θ_1)

}

New algorithm (n ≥ 1):

Repeat {

→ $\theta_j := \theta_j - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_j^{(i)}$
(simultaneously update θ_j for $j = 0, \dots, n$)

} x₀⁽ⁱ⁾ = 1

→ $\theta_0 := \theta_0 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_0^{(i)}$

→ $\theta_1 := \theta_1 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_1^{(i)}$

→ $\theta_2 := \theta_2 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_2^{(i)}$

...

