Strength Mean (Signed) (O->B units) — baseline versus post-test

Task	Group	n	Baseline ± SEM	Post-test ± SEM	Change ± SEM	Paired t p	Wilcoxon p
both	both	10	0.0897 ± 0.118	0.0542 ± 0.103	-0.0355 ± 0.102	0.737	0.625
both	strength	10	0.208 ± 0.185	0.0227 ± 0.045	-0.185 ± 0.162	0.281	0.375
both	angle	10	0.00237 ± 0.158	0.233 ± 0.105	0.231 ± 0.216	0.313	0.625
both	ALL	30	0.1 ± 0.0884	0.103 ± 0.0524	0.00325 ± 0.0981	0.974	0.584
strength	both	10	0.124 ± 0.0999	0.0412 ± 0.0354	-0.0824 ± 0.095	0.408	0.275
strength	strength	10	0.191 ± 0.158	0.0243 ± 0.0533	-0.167 ± 0.137	0.253	0.322
strength	angle	10	-0.0909 ± 0.0649	0.0875 ± 0.0457	0.178 ± 0.0816	0.0566	0.0645
strength	ALL	30	0.0746 ± 0.0674	0.051 ± 0.0258	-0.0237 ± 0.0656	0.721	0.685

SEM describes between-participant variation. Paired tests are exploratory and unadjusted; they do not propagate stage-fit uncertainty. ALL pools training groups descriptively.

Strength Variance ((O->B units)^2) — baseline versus post-test

Task	Group	n	Baseline ± SEM	Post-test ± SEM	Change ± SEM	Paired t p	Wilcoxon p
both	both	10	0.528 ± 0.22	0.156 ± 0.0257	-0.372 ± 0.202	0.0984	0.0137
both	strength	10	0.587 ± 0.14	0.171 ± 0.0337	-0.416 ± 0.152	0.0231	0.0195
both	angle	10	0.32 ± 0.0692	0.321 ± 0.0715	0.00136 ± 0.113	0.991	1
both	ALL	30	0.478 ± 0.0895	0.216 ± 0.0301	-0.262 ± 0.0956	0.0103	0.00403
strength	both	10	0.304 ± 0.102	0.188 ± 0.0573	-0.116 ± 0.113	0.331	0.131
strength	strength	10	0.466 ± 0.153	0.182 ± 0.0384	-0.284 ± 0.149	0.0892	0.0371
strength	angle	10	0.258 ± 0.0629	0.197 ± 0.0324	-0.0615 ± 0.0682	0.391	0.375
strength	ALL	30	0.343 ± 0.0647	0.189 ± 0.0245	-0.154 ± 0.0665	0.0278	0.00993

SEM describes between-participant variation. Paired tests are exploratory and unadjusted; they do not propagate stage-fit uncertainty. ALL pools training groups descriptively.

Angle Mean (Signed) (deg) — baseline versus post-test

Task	Group	n	Baseline ± SEM	Post-test ± SEM	Change ± SEM	Paired t p	Wilcoxon p
both	both	10	0.296 ± 0.542	0.406 ± 0.222	0.11 ± 0.571	0.852	0.695
both	strength	10	-1.06 ± 0.382	-0.171 ± 0.242	0.884 ± 0.333	0.0261	0.0137
both	angle	10	-0.928 ± 0.377	0.0107 ± 0.188	0.938 ± 0.412	0.0488	0.0488
both	ALL	30	-0.562 ± 0.27	0.0818 ± 0.13	0.644 ± 0.26	0.0194	0.00501
angle	both	10	0.337 ± 0.369	0.0607 ± 0.191	-0.277 ± 0.253	0.303	0.322
angle	strength	10	-0.624 ± 0.591	-0.316 ± 0.289	0.307 ± 0.591	0.615	0.77
angle	angle	10	-0.967 ± 0.935	0.112 ± 0.215	1.08 ± 0.789	0.205	0.0488
angle	ALL	30	-0.418 ± 0.389	-0.0479 ± 0.136	0.37 ± 0.343	0.29	0.416

SEM describes between-participant variation. Paired tests are exploratory and unadjusted; they do not propagate stage-fit uncertainty. ALL pools training groups descriptively.

Angle Variance (deg^2) — baseline versus post-test

Task	Group	n	Baseline ± SEM	Post-test ± SEM	Change ± SEM	Paired t p	Wilcoxon p
both	both	10	6.13 ± 4.07	2.74 ± 1.88	-3.39 ± 4.16	0.436	0.131
both	strength	10	10.6 ± 4.69	2.98 ± 1.66	-7.65 ± 4.45	0.12	0.0195
both	angle	10	7.34 ± 3.04	1.32 ± 0.413	-6.01 ± 2.7	0.0527	0.0488
both	ALL	30	8.03 ± 2.25	2.35 ± 0.829	-5.68 ± 2.17	0.0138	0.000505
angle	both	10	8.1 ± 4.44	1.94 ± 0.852	-6.16 ± 3.67	0.128	0.131
angle	strength	10	20 ± 8.52	6.6 ± 3.55	-13.4 ± 6.45	0.0674	0.232
angle	angle	10	11.8 ± 8.15	2.28 ± 0.931	-9.51 ± 8.05	0.268	0.275
angle	ALL	30	13.3 ± 4.16	3.61 ± 1.27	-9.7 ± 3.56	0.0109	0.0221

SEM describes between-participant variation. Paired tests are exploratory and unadjusted; they do not propagate stage-fit uncertainty. ALL pools training groups descriptively.

Mean Abs. Strength Error (O->B units) — baseline versus post-test

Task	Group	n	Baseline ± SEM	Post-test ± SEM	Change ± SEM	Paired t p	Wilcoxon p
both	both	10	0.566 ± 0.0951	0.367 ± 0.0659	-0.198 ± 0.111	0.107	0.084
both	strength	10	0.689 ± 0.104	0.31 ± 0.0292	-0.379 ± 0.0985	0.00394	0.00195
both	angle	10	0.581 ± 0.0538	0.474 ± 0.0849	-0.107 ± 0.0725	0.176	0.193
both	ALL	30	0.612 ± 0.0495	0.384 ± 0.038	-0.228 ± 0.0571	0.00041	0.000153
strength	both	10	0.441 ± 0.0736	0.3 ± 0.0442	-0.141 ± 0.0629	0.0515	0.0645
strength	strength	10	0.593 ± 0.103	0.319 ± 0.0369	-0.274 ± 0.0854	0.0107	0.00391
strength	angle	10	0.418 ± 0.0363	0.337 ± 0.0312	-0.0809 ± 0.0409	0.0793	0.0488
strength	ALL	30	0.484 ± 0.0449	0.319 ± 0.0212	-0.165 ± 0.0395	0.000242	1.6e-05

SEM describes between-participant variation. Paired tests are exploratory and unadjusted; they do not propagate stage-fit uncertainty. ALL pools training groups descriptively.

Mean Abs. Angle Error (deg) — baseline versus post-test

Task	Group	n	Baseline ± SEM	Post-test ± SEM	Change ± SEM	Paired t p	Wilcoxon p
both	both	10	1.66 ± 0.518	0.996 ± 0.297	-0.663 ± 0.463	0.185	0.16
both	strength	10	2.15 ± 0.511	1.13 ± 0.295	-1.02 ± 0.428	0.0409	0.0137
both	angle	10	1.79 ± 0.475	0.96 ± 0.153	-0.835 ± 0.374	0.0524	0.131
both	ALL	30	1.87 ± 0.282	1.03 ± 0.144	-0.839 ± 0.237	0.00137	0.00073
angle	both	10	1.89 ± 0.664	0.999 ± 0.284	-0.887 ± 0.413	0.0604	0.084
angle	strength	10	3.14 ± 0.799	1.65 ± 0.434	-1.49 ± 0.58	0.03	0.0488
angle	angle	10	2.15 ± 0.84	1.36 ± 0.318	-0.784 ± 0.831	0.37	0.695
angle	ALL	30	2.39 ± 0.441	1.34 ± 0.202	-1.05 ± 0.357	0.00615	0.0137

SEM describes between-participant variation. Paired tests are exploratory and unadjusted; they do not propagate stage-fit uncertainty. ALL pools training groups descriptively.