

3-Manifolds

Alec Zabel-Mena

February 19, 2023

Contents

1	Manifolds	5
1.1	Topological Manifolds	5
1.2	Smooth Manifolds	8

Chapter 1

Manifolds

1.1 Topological Manifolds

Definition. A **topological n -manifold** is a second countable Hausdorff space M , together with a collection $\{(M_\alpha, \phi_\alpha)\}$ for which

- (1) $\{M_\alpha\}$ is a collection of open sets of M covering M ; that is, $M_\alpha \subseteq M$ is open and $M = \bigcup M_\alpha$.
- (2) ϕ_α is a homeomorphism of M_α onto an open subset U of \mathbb{R}^n .



We call the pairs (M_α, ϕ_α) **charts** of M , and we call the collection of all such charts of M an **atlas** of M . We define the **dimension** of M to be $\dim M = n$.

Example 1.1. (1) Every subset of \mathbb{R}^N is second countable and Hausdorff, so that a subset \mathbb{R}^N is an n -manifold if every point of M has a neighborhood homeomorphic to \mathbb{R}^n , for $n \leq N$. In particular, \mathbb{R}^n is an n -manifold.

- (2) The n -sphere $S^n = \{x \in \mathbb{R}^{n+1} : \|x\| = 1\}$ (see figure 1.1) is an n -manifold. It is a second countable Hausdorff space, since it is a subspace of \mathbb{R}^n . Moreover, the stereographic projection $h : S^n \setminus (0, \dots, 0, 1) \rightarrow \mathbb{R}^n$ is a homeomorphism. So for $x \in S^n$, $x \neq (0, \dots, 0, 1)$, x has a neighborhood homeomorphic to \mathbb{R}^n . Now, if we take the composition of $\mathbb{R}^n \times \{0\}$ with h to obtain the map $h' : S^n \setminus (0, \dots, 0, -1) \rightarrow \mathbb{R}^n$, then we get that $S^n \setminus (0, \dots, 0, -1)$ is a neighborhood of $(0, \dots, 0, 1)$ homeomorphic to \mathbb{R}^n .


- (3) The n -torus $T = \underbrace{S^1 \times \dots \times S^1}_{n \text{ times}}$ (see figure 1.2) is the quotient space obtained from \mathbb{R}^n by identifying two points $x, y \in \mathbb{R}^n$ if, and only if there is some $g \in G$ for which $g(x) = y$, where G is the group generated by all translations by distance 1 along the coordinate axes. Let $x \in T^n$ and $U = \partial B(x, \frac{1}{4})$ the sphere centered about x of radius $\frac{1}{4}$ and let $q : \mathbb{R}^n \rightarrow T^n$ the quotient map of the quotient space of T^n . Then $q^{-1}|_{q(U)}$ is a homeomorphism. This makes T an n -manifold, with atlas $\{(U, q^{-1}|_{q(U)})\}$.

- (4) Identify the antipodal points of S^n , then the resulting quotient space is an n -manifold called **n -dimensional real projective space** which we denote by $\mathbb{P}\mathbb{R}^n$. Let $x \in$

Figure 1.1: The 2-Sphere of \mathbb{R}^3 is a 2-manifold.Figure 1.2: The 2-torus is a 2 manifold of \mathbb{R}^2 .

$\mathbb{P}\mathbb{R}^n$, since S^n is an n -manifold, there is a neighborhood U of x and a homeomorphism $h : U \rightarrow \mathbb{R}^n$. Let $-U = a(U)$, where $a : S^n \rightarrow S^n$ is the antipodal map. Then $-U$ is a neighborhood of $-x$, and $-h = h \circ a$ is a homeomorphism of U onto \mathbb{R}^n . Then the collection $\{(U, h)\}$ is an atlas for $\mathbb{P}\mathbb{R}^n$.

- (5) Consider the unit circle S^1 and the following “trefoil knot” K_3 pictured below in figure 1.3 Both of these are 1-manifolds, and are homeomorphic to each other. Consider the

Figure 1.3: The unit circle S^1 and the trefoil knot.

neighborhoods on either space (colored in red). Notice that each of these neighborhoods is homeomorphic to an interval in \mathbb{R}_1 , and hence homeomorphic to each other.

Definition. Let M be an n -dimensional manifold. A **p -dimensional submanifold** of M is a closed subset L of M for which there exists an atlas $\{(M_\alpha, \phi_\alpha)\}$ of M such that for all $x \in L$, there exists a chart (M_α, ϕ_α) in which $x \in M_\alpha$ and $\phi_\alpha(L \cap M_\alpha) = \{0\} \times \mathbb{R}^p$.

Lemma 1.1.1. *Submanifolds of manifolds are manifolds.*

Lemma 1.1.2. *Let M be an m -manifold, and N an n -manifold. Then the product $M \times N$ is an $(n + m)$ -manifold.*

Proof. We have that both M and N are Hausdorff, which makes $M \times N$ Hausdorff. Moreover, since M and N are second countable, they have countable bases \mathcal{B}_M and \mathcal{B}_N . Then the product $\mathcal{B}_M \times \mathcal{B}_N$ serves as a countable basis for $M \times N$.

Now, let $\{(M_\alpha, \phi_\alpha)\}$ and $\{(N_\beta, \psi_\beta)\}$ be atlases for M and N respectively. Then since each M_α is open in M , and each N_β is open in N , $M_\alpha \times N_\beta$ is open in $M \times N$. Moreover we also have that $M = \bigcup M_\alpha$, $N = \bigcup N_\beta$ so that

$$M \times N = \left(\bigcup M_\alpha\right) \times \left(\bigcup N_\beta\right) = \bigcup M_\alpha \times N_\beta$$

Now, we also have that ϕ_α is a homeomorphism of M_α onto an open subset of \mathbb{R}^m , and ψ_β is a homeomorphism of N_β onto an open subset of \mathbb{R}^n . Since ϕ_α and ψ_β are homeomorphisms, they are continuous with continuous inverses ϕ_α^{-1} and ψ_β^{-1} . This makes the map $\phi_\alpha \times \psi_\beta$ continuous with continuous inverse $(\phi_\alpha \times \psi_\beta)^{-1}$, which makes $\phi_\alpha \times \psi_\beta$ a homeomorphism of $M_\alpha \times N_\beta$ onto a subset of $\mathbb{R}^m \times \mathbb{R}^n \simeq \mathbb{R}^{m+n}$. Therefore $M \times N$ is an $(m + n)$ -manifold. ■

Example 1.2. The equator, S^1 of S^2 is a submanifold of S^2 (see figure 1.1).

Definition. We define the **boundry** of an n -manifold M to be the set ∂M , of all points of M for which there is a neighborhood homeomorphic to a neighborhood of H^n , but no neighborhood homeomorphic to a neighborhood of \mathbb{R}^n ; where $H^n = \{x \in \mathbb{R}^n : x_1 \geq 0\}$.

Definition. Let $H^n = \{(x_1, \dots, x_n) \in \mathbb{R}^n : x_1 \geq 0\}$. We define an **n -manifold with boundry** to be a second countable Hausdorff space M with atlas $\{(M_\alpha, \phi_\alpha)\}$ such that ϕ_α is a homeomorphism from M_α to an open subset of \mathbb{R}^n , or H^n .

Example 1.3. (1) The unit ball $B^n = \{x \in \mathbb{R}^n : \|x\| \leq 1\}$ is an n -dimensional manifold with boundry $\partial B^n = S^{n-1}$. For interior points of B^n , this is clear. For points in S^{n-1} , extending the stereographic projection gives the required homeomorphism.

(2) The **pair of pants** (see figure 1.4) Is a 2-manifold with boundry.




Figure 1.4:

(3) The 1-holed torus is a 2-manifold with boundry.

Definition. A p -dimensional submanifold with boundary of an n -dimensional manifold M is a closed subset L of M for which there is an atlas $\{(M_\alpha, \phi_\alpha)\}$ of M and $0 \leq p \leq n$, such that for every $x \in L$ in the interior of M , there is a chart (M_α, ϕ_α) such that $x \in M_\alpha$, and $\phi_\alpha(L \cap M_\alpha) = \{0\} \times \mathbb{R}^p$, and for every $x \in L$ in the boundary of M , there is a chart (M_α, ϕ_α) such that $x \in M_\alpha$, and with $\phi_\alpha(L \cap M_\alpha) = \{0\} \times \mathbb{R}^p$, and for which $\phi_\alpha(x) \in \{0\} \times \partial H^p$.

Lemma 1.1.3. *The boundary of an n -manifold is an $(n - 1)$ -submanifold with boundary.*

Example 1.4. The diameter of the ball B^2 is a submanifold with boundary.

Definition. We call an n -manifold M **closed** if M is compact with nonempty boundary ∂M .

Example 1.5. The n -sphere and n -torus are closed manifolds. Additionally, the projection map $\pi_y : T^2 \rightarrow S^1$ to $T^2 = S^1 \times S^1$ onto the second factor is a continuous map between manifolds.

Lemma 1.1.4. *If M is an n -manifold with boundary, then its boundary ∂M is an $(n - 1)$ -manifold without boundary.*

Proof. We first notice by our definition of boundary that $\partial M \subseteq M$. Since M is second countable and Hausdorff, ∂M inherits these properties as a subspace of M .

Now, consider a point $x \in \partial M$. Then by definition, there is a neighborhood U of x for which U is homeomorphic to a neighborhood V of H^n . Let $\phi : U \rightarrow V$ be the given homeomorphism. Notice, that $\partial H^n = \{x \in \mathbb{R}^n : x_1 = 0\}$, and that $\phi(x) \in V$ implies $\phi(x) \in V \cap \partial H^n$. Since V is open in H^n , $V \cap \partial H^n$ is open in ∂H^n as a subspace of H^n . So that $\phi(U) \simeq V \cap \partial H^n$ and U is homeomorphic to an open set in ∂H^n .

Now, take the projection map $\pi_y : \partial H^n \rightarrow \mathbb{R}^{n-1}$ onto the second factor, defined by $\pi_y : (0, (x_2, \dots, x_n)) \rightarrow (x_2, \dots, x_n) = (y_1, \dots, y_{n-1})$. Then π_y defines a homeomorphism of ∂H^n onto \mathbb{R}^{n-1} , so that $\pi_y \circ \phi$ is a homeomorphism of U onto an open set of \mathbb{R}^{n-1} . This makes ∂M an $(n - 1)$ -manifold. Moreover, since $x \in \partial M$ was arbitrary, there is no $x \in \partial M$ with neighborhood homeomorphic to an open set in H^{n-1} , so that $\partial(\partial M) = \emptyset$; i.e. ∂M is without boundary. ■

1.2 Smooth Manifolds

Definition. We call a map $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ q -**smooth**, or C^q , if it has continuous partial derivatives of order q . We call f **smooth**, or C^∞ , if it has continuous partial derivatives of all orders.

Definition. A C -**smooth manifold**, or C^q -**manifold** with $q > 0$ is a topological manifold with an atlas that is C^q . That is, for any charts (M_α, ϕ_α) and (M_β, ϕ_β) , $\phi_\beta \circ \phi_\alpha^{-1}$ is C^q wherever it is defined. We call C^∞ -manifolds **smooth manifolds**, or **differentiable manifolds**.

Example 1.6. (1) \mathbb{R}^n is a smooth manifold, as are all its open subsets.

- (2) Consider the n -manifold S^n with charts

$$(S^n \setminus (0, \dots, 0, 1), h) \qquad (S^n \setminus (0, \dots, 0, -1), h')$$

where

$$h(x_1, \dots, x_{n+1}) = \frac{1}{1 - x_{n+1}}(x_1, \dots, x_n) \text{ and } h'(x_1, \dots, x_{n+1}) = \frac{1}{1 + x_{n+1}}(x_1, \dots, x_n)$$

The map $h' \circ h^{-1}$ is smooth. Notice that

$$h^{-1}(y_1, \dots, y_n) = \left(\frac{2y_1}{1 + y_1^2 + \dots + y_n^2}, \dots, \frac{2y_n}{1 + y_1^2 + \dots + y_n^2} \right)$$

So that

$$h' \circ h^{-1} = \frac{1}{y_1^2 + \dots + y_n^2}(y_1, \dots, y_n)$$

Moreover, for all $q > 0$, $\partial^q h' \circ h^{-1}$ exists, which makes $h' \circ h^{-1}$ smooth. This makes S^n a smooth manifold.

- (3) The product of smooth manifolds are smooth manifolds. In particular, the torus $T^2 = S^1 \times S^1$ is a smooth manifold.

Definition. Let M and N manifolds with atlases $\{(M_\alpha, \phi_\alpha)\}$ and $\{(N_\beta, \psi_\beta)\}$. We call a map $f : M \rightarrow N$ **q -smooth**, or C^q if $\psi_\beta \circ \phi_\alpha^{-1} \circ f$ is C^q wherever it is defined. We call C^q -maps between manifolds **C^q -diffeomorphisms**. We call C^∞ -diffeomorphisms **diffeomorphisms**. We call any two C^q -manifolds **diffeomorphic** if there exists a C^q -diffeomorphism between them.

Example 1.7. (1) The map $f : \mathbb{R} \rightarrow \mathbb{R}$ given by $f(x) = x^3$ is a smooth map, but it is not a diffeomorphism, since $f'(x) = 3x^2$ has a singular point at 0 (elaborate?). It is not even a C^1 -diffeomorphism.

- (2) The projection map of $T^2 = S^1 \times S^1$ onto the second factor is a smooth map between manifolds.

Definition. Let M a C^q -manifold, for some $q \geq 1$, and let $x \in M$ and (M_α, ϕ_α) a chart containing x . We call x a **critical point** of a map $f : M \rightarrow \mathbb{R}$ if it is a critical point of $f \circ \phi_\alpha^{-1}$. If $g : \mathbb{R}^n \rightarrow \mathbb{R}$ is a map, we call x a **nondegenerate critical point** if the Hessian of g is nonsingular at x , and we call x a **nondegenerate critical point** of f if it is a nondegenerate critical point of $f \circ \phi_\alpha^{-1}$.

Definition. We define a **Morse function** on a manifold M to be a smooth map $f : M \rightarrow \mathbb{R}$ such that

- (1) f has nondegenerate critical points.
- (2) Distinct critical points map to distinct values.

Example 1.8. The projection map of the Torus $T^2 \subseteq \mathbb{R}^3$ on to the third coordinate is a map with critical points. It has 1 maximum value, 2 minimum values, and 2 saddle points. Moreover these critical points are nondegenerate, so that the projection is a Morse function. Notice that these critical points are nondegenerate, since they consist of minima, maxima, and saddle points. Moreover, we see that the sum of the minima and maxima, minus the saddle points is $1 + 1 - 2 = 0$.




Figure 1.5: A Morse function on the torus T^2 .

Definition. A q -smooth submanifold, or C^q -submanifold of a C^q -manifold M is a p -dimensional submanifold of M for which the transition maps restricted to L are C^q .

Definition. A q -smooth manifold, or C^q -manifold with boundary M is a topological manifold with boundary such that for any charts (M_α, ϕ_α) and (M_β, ϕ_β) in the atlas of M , the transition map $\phi_\beta \circ \phi_\alpha^{-1}$ is C^q wherever it is defined. We call C^∞ manifolds with boundary smooth manifolds, or differentiable manifolds with boundary.

Bibliography

- [1] D. Dummit, *Abstract algebra*. Hoboken, NJ: John Wiley & Sons, Inc, 2004.
- [2] I. N. Herstein, *Topics in algebra*. New York: Wiley, 1975.