

Ring Theory.

Alec Zabel-Mena

November 10, 2022

Contents

1	Rings.	5
1.1	Definitions and Examples.	5
1.2	Polynomial Rings, Matrix Rings, and Group Rings.	9
1.3	Ring Homomorphisms and Factor Rings.	11
1.4	Ideals.	16
1.5	Rings of Fractions.	19
1.6	Sun Tzu's Theorem.	22
2	Domains.	25
2.1	Euclidean Domains.	25

Chapter 1

Rings.

1.1 Definitions and Examples.

Definition. A **ring** R is a set together with two binary operations $+: (a, b) \rightarrow a + b$ and $\cdot: (a, b) \rightarrow ab$ called **addition** and **multiplication** such that:

- (1) R is an Abelian group over $+$, where we denote the identity element as 0 and the inverse of each $a \in R$ as $-a$.
- (2) R is closed under \cdot and \cdot is associative. That is, $ab \in R$ whenever $a, b \in R$ and $a(bc) = (ab)c$.
- (3) $a(b + c) = ab + ac$ and $(a + b)c = ac + bc$.

If $ab = ba$ for all $a, b \in R$, then we call R **commutative**. If there exists an element $1 \in R$ such that $a_1 = 1a = a$, then we call R a ring with **identity**.

Definition. A ring R with identity $1 \neq 0$ is called a **division ring** if for all $a \in R$, where $a \neq 0$, there exists a $b \in R$ such that $ab = ba = 1$. We call a commutative division ring a **field**.

Example 1.1. Let R be an abelian group under an operation $+$, define the operation \cdot by $(a, b) \rightarrow ab = 0$ for all $a, b \in R$. Then R is a ring under $+$ and \cdot , called the **trivial ring**. If $R = \langle e \rangle$, the trivial group, then we call R the **zero ring**.

- (2) The integers \mathbb{Z} form a ring under the usual addition and multiplication.
- (3) The sets of rational numbers \mathbb{Q} and the set of real numbers \mathbb{R} are rings under their usual addition and multiplication; in fact, they are fields. The complex numbers \mathbb{C} also form a field under complex addition and complex multiplication, where

$$\begin{aligned} + : (a + ib, c + id) &\rightarrow (a + c) + i(b + d) \\ \cdot : (a + ib, c + id) &\rightarrow (ac - bd) + i(ad + bc) \end{aligned}$$

- (4) The factor group of integers modulo n , $\mathbb{Z}/n\mathbb{Z}$ is a commutative ring under addition modulo n , and multiplication modulo n , $\mathbb{Z}/n\mathbb{Z}$ has identity $1 \pmod n$. $\mathbb{Z}/n\mathbb{Z}$ forms a field if, and only if $n = p^r$, where p is a prime.
- (5) We define the **real quaternions** to be the set $\mathbb{H} = \{a + ib_jc_kd : a, b, c, d \in \mathbb{R}, i^2 = j^2 = k^2 = -1 \text{ and } ij = k, jk = i, \text{ and } ki = j\}$. \mathbb{H} is a ring under addition and multiplication are defined for all $x = a + ib + jc + kd$ and $y = e + if + jg + kh$ to be:

$$\begin{aligned} +(x, y) : \rightarrow x + y &= (a + e) + i(b + f) + j(c + g) + k(d + h) \\ \cdot(x, y) : \rightarrow xy &= (a + ib + jc + kd)(e + if + jg + kh) \end{aligned}$$

- (6) Let A be a ring and R the set of all maps $f : X \rightarrow A$. Then R forms a ring under function addition $f + g(x) = f(x) + g(x)$ and function multiplication $fg(x) = f(x)g(x)$. Notice that R is commutative if, and only if A is, moreover, R has identity if, and only if A has identity.
- (7) We say a realvalued function $f : \mathbb{R} \rightarrow \mathbb{R}$ has **compact support** if there exist $a, b \in \mathbb{R}$ such that $f(x) = 0$ for all $x \notin [a, b]$. The set of all functions with compact support forms a ring without identity under function addition and function multiplication.
- (8) Let $X, Y \subseteq \mathbb{R}$. We denote the set of all continuous functions $f : X \rightarrow Y$ by $C(X, Y)$. Then $C(X, Y)$ forms a commutative ring with identity under function addition and function multiplication.

Lemma 1.1.1. *Let R be a ring. Then the following are true for all $a, b \in R$.*

- (1) $0a = a0 = 0$.
- (2) $(-a)b = a(-b) = -(ab)$.
- (3) $(-a)(-b) = ab$
- (4) *If R has identity $1 \neq 0$, then 1 is unique and $-a = (-1)a$.*

Proof. (1) Notice $0a = (0 + 0)a = 0a + 0a$, so that $0a = 0$. Likewise, $a0 = 0$ by the same reasoning.

- (2) Notice that $b - b = 0$, so $a(b - b) = ab + a(-b) = 0$, so that $a(-b) = -(ab)$. The same argument with $(a - a)b$ gives $(-a)b = -(ab)$.
- (3) By the inverse laws of addition in R , we have $-(a(-b)) = -(-(ab))$, so that $(-a)(-b) = ab$.
- (4) Suppose R has identity $1 \neq 0$, and suppose there is an element $2 \in R$ for which $2a = a2 = a$ for all $a \in R$. Then we have that $1 \cdot 2 = 1$ and $1 \cdot 2 = 2$, making $1 = 2$; so 1 is unique. Now, we have that $a + (-a) = 0$, so that $1(a + (-a)) = 1a + 1(-a) = 1a + (-a) = 0$ So $(-a) = -(1a) = (-1)a$ by (2). ■

Definition. Let R be a ring. We call an element $a \in R$ a **zero divisor** if $a \neq 0$ and there exists an element $b \neq 0$ such that $ab = 0$. Similarly, we call $a \in R$ a **unit** if there is a $b \in R$ for which $ab = ba = 1$.

Example 1.2. Notice if R is a ring with identity 1, then 1 is a unit of R by definition.

Definition. Let R be a ring. We call the set of all units in R the **group of units** and denote it R^* .

Lemma 1.1.2. *Let R be a ring with identity $1 \neq 0$. Then the group of units R^* forms a group under multiplication.*

Proof. Let $a, b \in R$ be units in R . Then there are $c, d \in R$ for which $ac = ca = 1$ and $bd = db = 1$. Consider then ab . Then $ab(dc) = a(bd)c = ac = 1$ and $(dc)ab = d(ca)b = db = 1$ so that ab is also a unit in R . Moreover R^* inherits the associativity of \cdot and 1 serves as the identity element of R^* . Lastly, if $a \in R^*$ is a unit there is a $b \in R$ for which $ab = ba = 1$. This also makes b a unit in R , and the inverse of a . ■

Corollary. *a is a zero divisor if, and only if it is not a unit.*

Proof. Suppose that $a \neq 0$ is a zero divisor. Then there is a $b \in R$ such that $b \neq 0$ and $ab = 0$. Then for any $v \in R$, $v(ab) = (va)b = 0$ so that a cannot be a unit. On the other hand let a be a unit, and $ab = 0$ for some $b \neq 0$. Then there is a $v \in R$ for which $v(ab) = (va)b = 1b = b = 0$. Then $b = 0$ which is a contradiction. ■

Corollary. *If R is a field, then it has no zero divisors.*

Proof. Notice by definition of a field, every element is a unit, except for 0. ■

Example 1.3. (1) \mathbb{Z} has no zero divisors, and has as units the elements -1 and 1 .

(2) For any $n \in \mathbb{Z}^+$, the units of $\mathbb{Z}/n\mathbb{Z}$ are all elements $a \bmod n$ such that $(a, n) = 1$. That is $(\mathbb{Z}/n\mathbb{Z})^* = U(\mathbb{Z}/n\mathbb{Z})$; recall that $U(\mathbb{Z}/n\mathbb{Z})$ is called the unit group, or group of units of $\mathbb{Z}/n\mathbb{Z}$.

(3) Let $D \in \mathbb{Q}$ be squarefree. Define $\mathbb{Q}(\sqrt{D}) = \{a + b\sqrt{D} : a, b \in \mathbb{Q}\}$. Then $\mathbb{Q}(\sqrt{D})$ is a field called the **quadratic field** under the operations

$$\begin{aligned} + : (a + b\sqrt{D}, c + d\sqrt{D}) &\rightarrow (a + c) + (b + d)\sqrt{D} \\ \cdot((a + b\sqrt{D}, c + d\sqrt{D})) &\rightarrow (ac - bdD) + (ad - bc)\sqrt{D} \end{aligned}$$

Since $\mathbb{Q}(\sqrt{D})$ is a field, every element is a unit.

Definition. A commutative ring with identity $1 \neq 0$ is called an **integral domain** if it has no zero divisors.

Lemma 1.1.3. *Let R be a ring, and a not a zero divisor. Then if $ab = ac$, then either $a = 0$, or $b = c$.*

Proof. Notice that $ab = ac$ implies $ab - ac = a(b - c) = 0$. Since a is not a zero divisor, either $a = 0$ or $b - c = 0$. ■

Corollary. *Any finite integral domain is a field.*

Proof. Let R be a finite integral domain and consider the map on R , by $x \rightarrow ax$. By above, this map is 1-1, moreover since R is finite, it is also onto. So there is a $b \in R$ for which $ab = 1$, making a a unit. Since a is arbitrarily chosen, this makes R a field. ■

Corollary. *If R is a field it is a (not necessarily finite) integral domain.*

Example 1.4. We have that fields are integral domains, and finite integral domains are fields. However, notice that not every integral domain need be a field. \mathbb{Z} is an integral domain that is not a field. Moreover, so are the real quaternions \mathbb{H} .

Definition. A **subring** of a ring R is a subgroup of R closed under multiplication.

Example 1.5. (1) We have the following sequence of subrings $\mathbb{Z} \subseteq \mathbb{Q} \subseteq \mathbb{R} \subseteq \mathbb{C}$.

- (2) The factor group $\mathbb{Z}/n\mathbb{Z}$ is not a subring of \mathbb{Z} , well the multiplication and addition of \mathbb{Z} is different from that of $\mathbb{Z}/n\mathbb{Z}$.
- (3) The set $\mathbb{Z} + i\mathbb{Z} + j\mathbb{Z} + k\mathbb{Z} \subseteq \mathbb{H}$ is a subring of \mathbb{H} .
- (4) If F is a field, then any subring of F is also an integral domain by inheretence.
- (5) The set $\mathbb{Z}[\sqrt{D}] = \{a + b\sqrt{D} : a, b \in \mathbb{Z}\}$ is a subring of the quadratic field $\mathbb{Q}(\sqrt{D})$. Moreover if $D \equiv 1 \pmod{4}$, then the set

$$\mathbb{Z}\left[\frac{1+\sqrt{D}}{2}\right] = \left\{a + b\frac{1+\sqrt{D}}{2} : a, b \in \mathbb{Z}\right\}$$

is also a subring of $\mathbb{Q}(\sqrt{D})$. We call the subring $\mathbb{Z}[\omega]$, where

$$\omega = \begin{cases} \sqrt{D}, & \text{if } D \not\equiv 1 \pmod{4} \\ \frac{1+\sqrt{D}}{2}, & \text{if } D \equiv 1 \pmod{4} \end{cases}$$

the **ring of integers** in the quadratic field. When $D = -1$, we get the ring $\mathbb{Z}[i]$, with $i^2 = -1$ and call it the **Gaussian integers**. Notice then that $\mathbb{Z}[i]$ is a subring of \mathbb{C} ; in fact, it is field in \mathbb{C} .

- (6) Consider $\mathbb{Q}(\sqrt{D})$ where D is squarefree. We define the **field norm** $N : \mathbb{Q}(\sqrt{D}) \rightarrow \mathbb{Q}$ by taking $(a + b\sqrt{D}) \rightarrow (a + b\sqrt{D})(a - b\sqrt{D}) = a^2 - Db^2$. If $D = i^2 = -1$, then $N : a + ib \rightarrow a^2 + b^2$ which is the modulus of complex number restricted to \mathbb{Q} .

Notice that if $z = a + b\sqrt{D}$, $w = c + d\sqrt{D}$, then $N(zw) = N(z)N(w)$ moreover,

$$N(a + \omega b) = \begin{cases} a^2 - Db^2, & \text{if } D \equiv 2, 3 \pmod{4} \\ a^2 + ab + \frac{1-D}{4}b^2, & \text{if } D \equiv 1 \pmod{4} \end{cases}$$

where

$$\omega = \begin{cases} \sqrt{D}, & \text{if } D \not\equiv 1 \pmod{4} \\ \frac{1+\sqrt{D}}{2}, & \text{if } D \equiv 1 \pmod{4} \end{cases}$$

In either case, $N : \mathbb{Z}[\omega] \rightarrow \mathbb{Z}$.

Lemma 1.1.4. *Let $\omega = \begin{cases} \sqrt{D}, & \text{if } D \not\equiv 1 \pmod{4} \\ \frac{1+\sqrt{D}}{2}, & \text{if } D \equiv 1 \pmod{4} \end{cases}$ where $D \in \mathbb{Z}^+$ is squarefree. Then an element of $z \in \mathbb{Z}[\omega]$ is a unit if, and only if $N(z) = \pm 1$*

Proof. Let $z = a + \omega b$ such that $N(z) = \pm 1$. Then we have

$$z^{-1} = \pm(a + \bar{\omega}b) \in \mathbb{Z}[\omega]$$

making it a unit. On the other hand, if $N(zw) = N(z)N(w) = \pm 1$, then since $N(z), N(w) \in \mathbb{Z}$, we must have that both $N(z) = \pm 1$ and $N(w) = \pm 1$. ■

1.2 Polynomail Rings, Matrix Rings, and Group Rings.

Theorem 1.2.1. *Let R be a commutative ring with identity, and define $R[x] = \{f(x) = a_0 + a_1x + \cdots + a_nx^n : a_0, \dots, a_n \in R\}$. Define the operations $+$ and \cdot on $R[x]$ for $f(x) = a_0 + a_1x + \cdots + a_nx^n$ and $g(x) = b_0 + b_1x + \cdots + b_nx^n$ by:*

$$f + g = (a_0 + b_0) + (a_1 + b_1)x + \cdots + (a_n + b_n)x^n$$

$$fg = c_0 + c_1x + \cdots + c_kx^k \text{ where } c_j = \sum_{i=0}^j a_i b_{j-i} \text{ and } k = n + m$$

Then $R[x]$ is a commutative ring with identity.

Definition. Let R be a commutative ring with identity. We call the ring $R[x]$ the **ring of polynomials** in x with **coefficients** in R whose elements of the form

$$f(x) = a_0 + a_1x + \cdots + a_nx^n$$

where $n \geq 0$ are called **polynomails**. If $a_n \neq 0$, then the **degree** of f is denoted $\deg f = n$, and f is called **monic** if $a_n = 1$. We call $+$ and \cdot the **addition** and **multiplication** of polynomials.

Example 1.6. (1) Take R any commutative ring with identity and form $R[x]$. One can verify that the polynomial $0(x) = 0 + 0x + \cdots + 0x^n + \cdots = 0$, in this case we call 0 the **zero polynomial**. Similarly, the additive inverse of $f(x) = a_0 + a_1x + \cdots + a_nx^n$ is the polynomial $-f(x) = -a_0 - a_1x - \cdots - a_nx^n$. Now, since $R[x]$ has identity, the **identity** polynomial is $1(x) = 1 + 0x + \cdots = 1$, that is, it is the identity in R . Lastly, we call a polynomial f with $\deg f = 0$ a **constant polynomial**. Notice that 0 and 1 are constant polynomials.

- (2) $\mathbb{Z}[x]$, $\mathbb{Q}[x]$, $\mathbb{R}[x]$ and $\mathbb{C}[x]$ are the polynomial rings in x with coefficients in \mathbb{Z} , \mathbb{Q} , \mathbb{R} , and \mathbb{C} respectively.
- (3) Notice that the rings $\mathbb{Z}[\omega]$ and $\mathbb{Z}[i]$ are polynomial rings in ω and i , respectively, with coefficients in \mathbb{Z} , and where $\omega = \sqrt{D}$ if $D \not\equiv 1 \pmod{4}$ or $\omega = \frac{1+\sqrt{D}}{2}$ otherwise, and $i^2 = -1$. Notice that the highest degree a polynomial in $\mathbb{Z}[i]$ can achieve is $\deg = 1$; however, one may be able to form polynomial rings in other variables with coefficients in $\mathbb{Z}[i]$, i.e. take $Z[x]$, where $Z = \mathbb{Z}[i]$.
- (4) $\mathbb{Z}/3\mathbb{Z}[x]$ is the polynomial ring with coefficients in $\mathbb{Z}/3\mathbb{Z}$.

Theorem 1.2.2. *Let R be an integral domain, and let $p, q \neq 0$ be polynomials in $R[x]$. Then the following are true:*

- (1) $\deg pq = \deg p + \deg q$.
- (2) The units of $R[x]$ are precisely the units of R .
- (3) $R[x]$ is an integral domain.

Proof. Consider the leading terms $a_n x^n$ and $b_m x^m$ of p and q respectively. Then $a_n b_m x^{m+n}$ is the leading term of pq ; moreover we require $a_n b_m \neq 0$. Now, if $\deg pq < m + n$, then $ab = 0$, making a and b zero divisors of R ; impossible. Therefore $ab \neq 0$. It also follows that since no term of p is a zero divisor, then p cannot be a zero divisor of $R[x]$. Lastly, if $pq = 1$, then $\deg p + \deg q = 0$, so that pq is a constant polynomial. Noticing that constant polynomials are simply just elements of R , then p and q are units. ■

Theorem 1.2.3. *Let R be a ring. Let $R^{n \times n}$ be the set of all $n \times n$ matrices with entries in R and define the operations $+$ and \cdot by:*

$$(a_{ij}) + (b_{ij}) = (a_{ij} + b_{ij})$$

$$(a_{ij})(b_{ij}) = (c_{ij}), \text{ where } c_{ij} = \sum_{k=1}^n a_{ik} b_{kj}$$

Then $R^{n \times n}$ forms a ring under $+$ and \cdot .

Definition. For any ring R , we call the ring $R^{n \times n}$ the **matrix ring** of $n \times n$ matrices with entries in R .

Example 1.7. (1) Note that if R is a commutative ring, then for $n \geq 2$, $R^{n \times n}$ need not be commutative.

- (2) We call matrices of $R^{n \times n}$, for $n \in \mathbb{Z}^+$ **square matrices**. We call a matrix $(a_{ij}) \in R^{n \times n}$ **scalar** if $a_{ii} = 1$ for all $1 \leq i \leq n$ and $a_{ij} = 0$ whenever $i \neq j$.
- (3) If R has identity, then so does $R^{n \times n}$. We call the identity of $R^{n \times n}$ the **identity matrix** and denote it as the $n \times n$ scalar matrix I with 1 across the diagonal. We call the units of $R^{n \times n}$ **invertible** matrices, and denote the unit group of invertible matrices to be $GL(n, R)$ the general linear group of degree n over R .

- (4) Notice that $2\mathbb{Z}^{n \times n} \subseteq \mathbb{Z}^{n \times n} \subseteq \mathbb{Q}^{n \times n} \subseteq \mathbb{R}^{n \times n} \subseteq \mathbb{C}^{n \times n}$.
- (5) Let R be a ring, and $R^{n \times n}$ its matrix ring. Let $U^{n \times n} = \{(a_{ij}) : a_{pq} = 0 \text{ whenever } p > q\}$ the set of **upper triangular matrices**. Then $U^{n \times n} \subseteq R^{n \times n}$ is a subring.

Theorem 1.2.4. *Let R be a ring with identity, and let G be a finite group of order n . Let RG the set of all sums $a_1g_1 + \cdots + a_ng_n$, where $a_i \in R$ for all $1 \leq i \leq n$. Define the operations $+$ and \cdot by:*

$$(a_1g_1 + \cdots + a_ng_n) + (b_1g_1 + \cdots + b_ng_n) = (a_1 + b_1)g_1 + \cdots + (a_n + b_n)g_n$$

$$(a_1g_1 + \cdots + a_ng_n)(b_1g_1 + \cdots + b_ng_n) = c_1g_1 + \cdots + c_ng_n, \text{ where } c_k = \sum_{g_k = g_i g_j} a_i b_j$$

Then RG forms a ring with identity under $+$ and \cdot . Moreover, RG is commutative if, and only if G is abelian.

Definition. Let R be a ring with identity, and let G be a finite group of order n . We call the ring RG the **group ring** of G . We call the elements of RG **formal sums** of the elements of G .

Example 1.8. (1) Consider $D_8 = \langle r, t : r^4 = t^2 = 1, rt = tr^{-1} \rangle$ and \mathbb{Z} . Let $a, b \in \mathbb{Z}D_8$ where $a = r + r^2 - 2t$ and $b = -3r^2 + rt$. Then

$$a + b = r - 2r^2 + rt - t$$

$$ab = -5r^3 + r^3t + 7r^2t - 3$$

- (2) For any ring with identity R , and finite group G , $R \subseteq RG$, for take the elements of R to be the sums $a_1 + \cdots + a_n$. $G \subseteq RG$, for $g_i = 1g_i$; moreover, each g_i has an inverse in RG , so we call G the subgroup of units of RG .
- (3) Let G be a group with $\text{ord } G > 1$. Let $g \in G$ with $\text{ord } g = m$. Notice that the elements $(1 - g), (1 + g + \cdots + g^{m-1}) \in RG$ are nonzero, but that

$$(1 - g)(1 + g + \cdots + g^{m-1}) = 1 - g^m = 1 - 1 = 0$$

which makes $1 - g$ a zero divisor. In general, the ring RG will always have zero divisors.

- (4) Let G be a finite group. We call the rings $\mathbb{Z}G, \mathbb{Q}G, \mathbb{R}G$, and $\mathbb{C}G$ the **integral, rational, real, and complex** group rings of G , respectively. Notice that $\mathbb{Z}G \subseteq \mathbb{Q}G \subseteq \mathbb{R}G \subseteq \mathbb{C}G$. Moreover, if $H \leq G$ is a subgroup of G , then $RH \subseteq RG$ is a subring.

1.3 Ring Homomorphisms and Factor Rings.

Definition. Let R and S be rings. We call a map $\phi : R \rightarrow S$ a **ring homomorphism** if

- (1) ϕ is a group homomorphism with respect to addition.

- (2) $\phi(ab) = \phi(a)\phi(b)$ for any $a, b \in R$.

We denote the **kernel** of ϕ to be the kernel of ϕ as a group homomorphism. That is

$$\ker \phi = \{r \in R : \phi(r) = 0\}$$

Moreover, if ϕ is 1-1 and onto, we call ϕ an **isomorphism** and say that R and S are **isomorphic**, and write $R \simeq S$.

Example 1.9. (1) $\phi : \mathbb{Z} \rightarrow \mathbb{Z}/2\mathbb{Z}$ defined by $n \rightarrow 0$ if n is even and $n \rightarrow 1$ if n is odd is a ring homomorphism, with $\ker \phi = 2\mathbb{Z}$. Notice that $\phi(\mathbb{Z}) = \mathbb{Z}/2\mathbb{Z}$. ϕ is onto, but not 1-1.

- (2) Let $n \in \mathbb{Z}$ and consider the maps $\phi_n : \mathbb{Z} \rightarrow \mathbb{Z}$ by taking $x \rightarrow nx$. ϕ_n , in general is not a ring homomorphism, as $\phi(xy) = n(xy)$ but $\phi(x)\phi(y) = nxy = n^2(xy)$. ϕ_n , however is a group homomorphism for any n .

- (3) For any ring R , define the **valuation** map $\phi : R[x] \rightarrow R$ by taking $f(x) \rightarrow f(0)$; i.e. the polynomial f evaluated at 0. ϕ is a ring homomorphism. Moreover, notice that if $f(x) = a_0 + a_1x + \cdots + a_nx^n$, then $f(0) = a_0 \in R$. So that $\phi(R[x]) = R$. This makes ϕ onto. Now, take $\phi(f) = 0$. Then those are all polynomials with constant term $a_0 = 0$ (this does not make $\ker \phi = \langle e \rangle$). Again, ϕ is onto, but it is not 1-1.

Lemma 1.3.1. *Let R and S be rings, and $\phi : R \rightarrow S$ a ring homomorphism. Then*

- (1) $\phi(R)$ is a subring of S .

- (2) $\ker \phi$ is a subring of R .

Proof. Let $s_1, s_2 \in \phi(R)$. Then $s_1 = \phi(r_1)$ and $s_2 = \phi(r_2)$ for some $r_1, r_2 \in R$. Then $s_1s_2 = \phi(r_1)\phi(r_2) = \phi(r_1r_2) \in \phi(S)$. Additionally, $s^{-1} = \phi^{-1}(r) = \phi(r^{-1})$ for some $s \in S$, $r \in R$. This is sufficient to make S a subring of S .

By similar reasoning, if $r_1, r_2 \in \ker \phi$, then $\phi(r_1)\phi(r_2) = \phi(r_1r_2) = 0$ so that $r_1r_2 \in \ker \phi$, and $\phi(r^{-1}) = \phi^{-1}(r) = 0$ so $\phi^{-1} \in \ker \phi$. ■

Corollary. *For any $r \in R$ and $a \in \ker \phi$, then $ar \in \ker \phi$ and $ra \in \ker \phi$.*

Proof. We have $\phi(ar) = \phi(a)\phi(r) = \phi(a)0 = 0$ so $ar \in \ker \phi$. The same happens for ra . ■

Definition. Let R be a ring. We call a subring $I \subseteq R$ of R a **left ideal** in R if for any $r \in R$ and $a \in I$, we have $ar \in I$. Similarly, we call I a **right ideal** in R if $ra \in I$. We call I a **(two-sided) ideal** in R if it is both a left, and a right ideal and we say that the ideals I **absorb** r .

Lemma 1.3.2. *If R is a commutative ring, then every left ideal is a right ideal.*

Proof. Notice that $ar = ra$ for all $a, r \in R$. ■

Theorem 1.3.3. Let R be a ring, and I an ideal in R . Let R/I be the set of all $a + I$ with $a \in R$. Define operations $+$ and \cdot by

$$\begin{aligned}(a + I) + (b + I) &= (a + b) + I \\ (a + I)(b + I) &= ab + I\end{aligned}$$

Then R/I forms a ring under $+$ and \cdot .

Proof. Notice that $(a + I) + (b + I) = (a + b) + (I + I) = (a + b) + 2I = (a + b) + I$. Moreover, R/I inherits associativity in $+$ from addition in R . Now, take $0 + I = I$ as the additive identity and $-a + I$ as the inverse of $a + I$ for each I .

Now, notice, that $(a + I)(b + I) = ab + aI + bI + I^2 = ab + (I + I + I) = ab + I$ by distribution of multiplication over addition in R . Moreover, R/I also inherits associativity in \cdot of multiplication in R . Now, notice then that

$$(a + I)((b + I) + c + I) = (a + I)((b + c) + I) = a(b + c) + I = (ab + ac) + I = (ac + I) + (bc + I)$$

and

$$((a + I) + (b + I))(c + I) = ((a + b) + I)(c + I) = (a + b)c + I = (ac + bc) + I = (ac + I) + (bc + I)$$

Lastly, notice that $a + I$ is just the left coset of a by I in R as a group under addition. So that $+$ and \cdot are coset addition and multiplication, which are well defined. ■

Corollary. If R has identity 1, then R/I has identity $1 + I$. Moreover if R is commutative, then so is R/I .

Definition. Let R be a ring and I an ideal in R . We call the ring R/I under addition and multiplication of cosets the **factor ring** (or **quotient ring**) of R over I .

Example 1.10. (1) We call $(0) = \{0\}$ the **trivial ideal**, notice also that R is also an ideal.

(2) For any $n \in \mathbb{Z}$, notice that if $a \in \mathbb{Z}$ and $m \in n\mathbb{Z}$, then $m = nk$, for some $k \in \mathbb{Z}$ so that $am = n(ak) = ma \in n\mathbb{Z}$. So $n\mathbb{Z}$ is an ideal of \mathbb{Z} , with factor ring $\mathbb{Z}/n\mathbb{Z}$. So $\mathbb{Z}/n\mathbb{Z}$ is a factor ring on top of also being a factor group. We call the ring homomorphism $\phi : \mathbb{Z} \rightarrow \mathbb{Z}/n\mathbb{Z}$ by $a \rightarrow a \bmod n$ the **reduction homomorphism**.

(2) Let R a ring, and consider $R[x]$. Let I the set of all polynomials of degree greater than 2 together with 0. Then if $f \in I$, $\deg f > 2$ or $f = 0$. Then for any $g \in R[x]$, $\deg fg > 2$ or, $fg = 0$ and $\deg gf > 2$ or $gf = 0$. This makes I an ideal of $R[x]$. Moreover, $p, q \in I$ if and only if they have the same constant term. Notice then that $R[x]/I = \{a + bx : a, b \in R\}$.

Now, if R has no zero divisors, it is possible that $R[x]/I$ has zero divisors. Consider $\mathbb{Z}[x]/I$.

- (3) Let A a ring, and $X \neq \emptyset$. For the ring of functionss A^X , for a given $c \in X$, define the **valuation** map at c by $E_c : f(x) \rightarrow f(c)$. Notice that E_c is a ring homomorphism, so that $A^X / \ker E_c$ forms a factor ring. IN particular, if $A^X = A[x]$ the polynomial ring over A , and $c = 0$, then E_c is just the valuation map of polynomials.

Now, if $X = (0, 1]$, and $R = \mathbb{R}^{(0,1]}$, by the first isomorphism theorem, we have $\mathbb{R} \simeq \mathbb{R}^{(0,1]} / \ker E_c$, since $E_c(\mathbb{R}^{(0,1]}) = \mathbb{R}$.

- (4) Let $n \geq 2$ and consider $R^{n \times n}$. Let J an ideal of R . Then $J^{n \times n} = \{(a_{ij}) : a_{ij} \in J\}$ is an ideal of $R^{n \times n}$. Take the ring homomorphism

$$\begin{aligned} R^{n \times n} &\rightarrow (R/J)^{n \times n} \\ (a_{ij}) &\rightarrow (a_{ij} + J) \end{aligned}$$

Then $J^{n \times n}$ is the kernel of this homomorphism, so that

$$R^{n \times n} / J^{n \times n} \simeq (R/J)^{n \times n}$$

For example, with $n = 3$, we have

$$\mathbb{Z}^{3 \times 3} / 2\mathbb{Z}^{3 \times 3} \simeq (\mathbb{Z}/2\mathbb{Z})^{3 \times 3}$$

- (5) Let R a commutative ring with identity, and G a finite group of order n . Define the **augmentation** map to be the map

$$\begin{aligned} RG &\rightarrow R \\ \sum_{i=1}^n a_i g_i &\rightarrow \sum_{i=1}^n a_i \end{aligned}$$

We call the kernel of this map the **augmentation ideal** which is the set of all formal sums whose coefficients sum to 0. Another ideal of RG is the set $I = \{\sum a g_i : g_i \in G\}$ the set of all formal sums whose coefficients are all equal.

Theorem 1.3.4 (The First Isomorphism Theorem). *If $\phi : R \rightarrow S$ is a ring homomorphism from rings R into S , then $\ker \phi$ is an ideal of R and*

$$\begin{array}{ccc} & \phi(R) \simeq R / \ker \phi & \\ & \nearrow & \\ R & \xrightarrow{\quad \phi \quad} & S \\ \downarrow \pi & \searrow \bar{\phi} & \\ R / \ker \phi & & \end{array}$$

Proof. By the first isomorphism theorem for groups, ϕ is a group isomorphism. Now, let $K = \ker \phi$ and consider the map $\pi : R \rightarrow R/I$ by $a \xrightarrow{\pi} a+K$. Define the map $\bar{\phi} : R/K \rightarrow \phi(R)$ such that $\bar{\phi} \circ \pi = \phi$, then $\bar{\phi}$ defines the ring isomorphism. ■

Proof. The map $\pi : R \rightarrow R/I$ defined by $a \rightarrow a + I$, for any ideal I , is onto, with $\ker \pi = I$. ■

Theorem 1.3.5 (The Second Isomorphism Theorem). *Let $A \subseteq R$ a subring of R , and let B an ideal in R . Define $A + B = \{a + b : a \in A \text{ and } b \in B\}$. Then $A + B$ is a subring and $A \cap B$ is an ideal in A . Then*

$$A + B/B \simeq A/A \cap B$$

Theorem 1.3.6 (The Third Isomorphism Theorem). *Let I and J be ideals in a ring R , with $I \subseteq J$. Then J/I is an ideal of R/I and*


$$R/J = (R/I)/(J/I)$$

Theorem 1.3.7 (The Fourth Isomorphism Theorem). *Let I an ideal in a ring R , then the correspondence between A and A/I , for any subring $A \subseteq R$ is an inclusion preserving bijection between subrings of A containing I and R/I . Moreover, A is an ideal if, and only if A/I is an ideal.*

Example 1.11. We have $12\mathbb{Z}$ is an ideal of \mathbb{Z} , and that $\mathbb{Z}/12\mathbb{Z}$ has as ideals

$$\mathbb{Z}/12\mathbb{Z} \quad 2\mathbb{Z}/12\mathbb{Z} \quad 3\mathbb{Z}/12\mathbb{Z} \quad 4\mathbb{Z}/12\mathbb{Z} \quad 6\mathbb{Z}/12\mathbb{Z} \quad 12\mathbb{Z}/12\mathbb{Z}$$

Lemma 1.3.8. *Let R be a ring with ideals I and J . Then $I + J$, IJ and I^n , for any $n \geq 0$ are ideals of R and we have the lattice*



Example 1.12. (1) Consider the ideals $6\mathbb{Z}$ and $10\mathbb{Z}$ of \mathbb{Z} . Then $6\mathbb{Z} + 10\mathbb{Z}$ is the ideal consisting of all integers of the form $6x + 10y$. Now, for $x, y \in \mathbb{Z}$, since $(6, 10) = 2$,

we have that $6\mathbb{Z} + 10\mathbb{Z} \subseteq 2\mathbb{Z}$ since $6x + 10y = 2(3x + 5y)$. Now, we also have that $2 = 6 \cdot 2 + 10 \cdot -1$ so that $2 \in 6\mathbb{Z} + 10\mathbb{Z}$ which makes $2\mathbb{Z} \subseteq 6\mathbb{Z} + 10\mathbb{Z}$. Thus, we have $6\mathbb{Z} + 10\mathbb{Z} = 2\mathbb{Z}$. In general, we have that $m\mathbb{Z} + n\mathbb{Z} = d\mathbb{Z}$ where $d = (m, n)$ is the greatest common divisor of m and n . The ideal $6\mathbb{Z}10\mathbb{Z}$ gives all integers of the form $6x10y = 6 \cdot 10(xy) = 60(xy)$, so that $6\mathbb{Z}10\mathbb{Z} = 60\mathbb{Z}$.

- (2) Let $I \subseteq \mathbb{Z}[x]$ the ideal of polynomials with even constant term. Notice that $2, x = x + 0 \in I$ so that $4, x^2 \in I^2 = II$. So that $4 + x^2 \in I^2$ which is not in general divisible by elements in I .

1.4 Ideals.

Definition. Let R be a commutative ring with identity. We call the smallest ideal containing a nonempty subset A in R the **ideal generated** by A , and we write (A) . We call an ideal **principle** if it is generated by a single element of R , i.e. $I = (a)$ for some $a \in I$. We say that the ideal (A) is **finitely generated** if $|A|$ is finite, and if $A = \{a_1, \dots, a_n\}$, then we denote $(A) = (a_1, \dots, a_n)$.

Example 1.13. (1) In any commutative ring with identity, the trivial ideal and R are the ideals generated by 0 and 1, respectively, so we write them as (0) and $R = (1)$.

- (2) In \mathbb{Z} , we can write the ideals $n\mathbb{Z} = (n) = (-n)$. Notice that every ideal in \mathbb{Z} is a principle ideal. Moreover, for $m, n \in \mathbb{Z}$, $n|m$ if, and only if $n\mathbb{Z} \subseteq m\mathbb{Z}$. Notice that $m\mathbb{Z} + n\mathbb{Z} = d\mathbb{Z}$ is the ideal generated by m and n , where $d = (m, n)$ is the greatest common divisor of m and n . Indeed, by definition, $d|m, n$ so that $d\mathbb{Z} \subseteq m\mathbb{Z} + n\mathbb{Z}$, and if $c|m, n$, then $c|d$, making $m\mathbb{Z} + n\mathbb{Z} \subseteq d\mathbb{Z}$. Then $m\mathbb{Z} + n\mathbb{Z} = d\mathbb{Z}$ is the ideal generated by the greatest common divisor (m, n) and consists of all diophantine equations of the form

$$mx + ny = (m, n)$$

In general, we can define the **greatest common divisor** for integers n_1, n_2, \dots, n_m to be the smallest such integer d generating the ideal $n_1\mathbb{Z} + \dots + n_m\mathbb{Z} = d\mathbb{Z}$. We then write $d = (n_1, \dots, n_m)$.

- (3) Consider the ideal $(2, x)$ of $\mathbb{Z}[x]$. $(2, x)$ is not a principle ideal. We have that $(2, x) = \{2p_xq : p, q \in \mathbb{Z}[x]\}$, and that $(2, x) \neq \mathbb{Z}[x]$. Suppose that $(2, x) = (a)$ for some polynomial $a \in \mathbb{Z}[x]$, then $2 \in (a)$, so that $2 = p(x)a(x)$, of degree $\deg p + \deg a$. This makes p and a constant polynomials in $\mathbb{Z}[x]$. Now, since 2 is prime in \mathbb{Z} , then only values for p and q are $p = \pm 1$ and $a = \pm 2$. If $a(x) = \pm 1$, then every polynomial in $\mathbb{Z}[x]$ can be written as a polynomial in (a) , so that $(a) = \mathbb{Z}[x]$, impossible. If $a(x) = \pm 2$, then since $x \in (a)$, we get $x = 2q(x)$ where $q \in \mathbb{Z}[x]$. This cannot happen, so that $(a) \neq (2, x)$.
- (4) Consider $\mathbb{R}^{[0,1]}$ the ring of all functions $f : [0, 1] \rightarrow \mathbb{R}$. Let $M = \{f : f(\frac{1}{2}) = 0\}$. Then M is an ideal in $\mathbb{R}^{[0,1]}$, in fact, notice that it is the kernel of the valuation map at $\frac{1}{2}$.

Define $g : [0, 1] \rightarrow \mathbb{R}$ by:

$$g(x) = \begin{cases} 1, & \text{if } x \neq \frac{1}{2} \\ 0, & \text{if } x = \frac{1}{2} \end{cases}$$

then $f = fg$ by definition of both f and g . So that $M = (g)$ which makes M a principle ideal. M is not principle in general, consider $C^{[0,1]}$ the set of all realvalued continuous functions on $[0, 1]$.

- (5) Let G be a finite group and R a commutative ring with identity. The augmentation ideal in RG is generated by the set $\{g - 1 : g \in G\}$, and we write $(g_1 - 1, \dots, g_n - 1)$ where $\text{ord } G = n$. If G is cyclic, then the augmentation ideal is just $(g - 1)$, and is principle.

Lemma 1.4.1. *Let I an ideal in ring R with identity. Then*

- (1) $I = (1)$ if, and only if I contains a unit.
 (2) If R is commutative, then R is a field if, and only if its only ideals are (0) and (1) .

Proof. Recall that $R = (1)$. Now, if $I = (1)$, then $1 \in I$, and 1 is a unit. Conversely, suppose that $u \in I$ with u a unit. By definition, we have that $r = r \cdot 1 = r(uv) = r(vu) = (rv)u$, so that $1 \in I$. This makes $I = (1)$.

Now, if R is a field, then it is a commutative ring, moreover every $r \neq 0$ is a unit in R , which makes $r \in I$ for some ideal $I \neq (0)$. This makes every $I \neq (0)$ equal to (1) . Conversely, if (0) and (1) are the only ideals of the commutative ring R , then every $r \neq 0 \in (1)$, which makes them units. Hence all nonzero r is a unit in R . This makes R into a field. ■

Corollary. *If R is a field, then any nonzero ring homomorphism $\phi : R \rightarrow S$ is 1-1.*

Proof. If R is a field, then either $\ker \phi = (0)$ or $\ker \phi = (1)$. Now, since $\ker \phi \neq R$, we must have $\ker \phi = (0)$. ■

Definition. We call a ring D with identity a **division ring** if its only left and right ideals are (0) and (1) respectively.

Example 1.14. For any field F , the only two sided ideals of $F^{n \times n}$ are (0) and (1) , so that $F^{n \times n}$ is a division ring.

Definition. For any ideal M in a ring R , we call M **maximal** if $M \neq R$, and if N is an ideal with $M \subseteq N \subseteq R$, then either $M = N$ or $N = R$.

Lemma 1.4.2. *If R is a ring with identity, every proper ideal is contained in a maximal ideal.*

Proof. Let I a proper ideal of R . Let $\mathcal{S} = \{N : N \neq (1) \text{ is a proper ideal, and } I \subseteq N\}$. Then $\mathcal{S} \neq \emptyset$, as $I \in \mathcal{S}$, and the relation \subseteq partially orders \mathcal{S} . Let \mathcal{C} be a chain in and define

$$J = \bigcup_{A \in \mathcal{C}} A$$

We have that $J \neq \emptyset$ since $(0) \in J$. Now, let $a, b \in J$, then we have that either $(a) \subseteq (b)$ or $(b) \subseteq (a)$, but not both. In either case, we have $a - b \in J$ so that J is closed under additive inverse. Moreover, since $A \in \mathcal{C}$ is an ideal, by definition, J is closed with respect to absorption. This makes J an ideal.

Now, if $1 \in J$, then $J = (1)$ and J is not proper, and $A = (1)$ by definition of J . This is a contradiction as A must be proper. Therefore J must also be a proper ideal. Therefore, \mathcal{C} has an upperbound in \mathcal{S} , therefore, by Zorn's lemma, \mathcal{S} has a maximal element M , i.e. it has a maximal ideal M with $I \subseteq M$. ■

Lemma 1.4.3. *Let R be a commutative ring with identity. An ideal M is maximal if, and only if R/M is a field.*

Proof. If M is maximal, then there is no ideal $I \neq (1)$ for which $M \subseteq I \subseteq R$. By the fourth isomorphism theorem, the ideals of R containing M are in 1-1 correspondence with the those of R/M . Therefore M is maximal if, and only if the only ideals of R/M are (M) and $(1 + M)$. ■

Example 1.15. (1) Let $n \geq 0$ an integer. The ideal $n\mathbb{Z}$ is maximal in \mathbb{Z} if and only if $\mathbb{Z}/n\mathbb{Z}$ is a field. Therefore $n\mathbb{Z}$ is maximal if, and only if $n = p$ a prime in \mathbb{Z} . So the maximal ideals of \mathbb{Z} are those $p\mathbb{Z}$ where p is prime.

(2) $(2, x)$ is not principle in $\mathbb{Z}[x]$, but it is maximal in $\mathbb{Z}[x]$, as $\mathbb{Z}[x]/(2, x) \simeq \mathbb{Z}/2\mathbb{Z}$ which is a field.

(3) The ideal (x) is not maximal in $\mathbb{Z}/n\mathbb{Z}$, since $\mathbb{Z}/(x) \simeq \mathbb{Z}$, which is not a field. Moreover, $(x) \subseteq (2, x) \subseteq \mathbb{Z}[x]$. We construct this isomorphism by identifying $x = 0$, then all polynomials of $\mathbb{Z}[x]/(x)$ only have constant term in \mathbb{Z} .

(4) Let $a \in [0, 1]$, and $M_a = \{f : f(a) = 0\}$ the kernel of the valuation map at a . Then M is principle, moreover, we also have that since $f(a) \in \mathbb{R}$, then $\mathbb{R}^{[0,1]}/M_a \simeq \mathbb{R}$ which makes M_a maximal.

(5) If F is a field and G a finite group of order n , then the augmentation ideal $(g_1 - 1, \dots, g_n - 1)$ is maximal in FG . Let $\pi : \sum g_i a_i \rightarrow \sum a_i$, then $\ker \pi = (g_1 - 1, \dots, g_n - 1)$ and $\pi(FG) = F$. This makes $FG/(g_1 - 1, \dots, g_n - 1) \simeq F$.

Definition. We call an ideal P in a commutative ring R with identity **prime** if $P \neq (1)$ and if $ab \in P$ then either $a \in P$ or $b \in P$. Alternatively, if $(ab) \subseteq P$ then $(a) \subseteq P$ or $(b) \subseteq P$.

Example 1.16. The prime ideals of \mathbb{Z} are $p\mathbb{Z}$ with p prime together with (0) .

Lemma 1.4.4. *An ideal P in a commutative ring with identity, R , is prime if, and only if R/P is an integral domain.*

Proof. Suppose that P is prime, and let $(a + P)(b + P) = ab + P = P$. This gives us that $ab \in P$ and hence $a \in P$ or $b \in P$. Then either $a + P = P$ or $b + P = P$ in R/P . Conversely, if R/P is an integral domain, then for any $a + P, b + P$ $ab + P = P$ implies that either $a + P = P$ or $b + P = P$. Then $a \in P$ or $b \in P$, only when $ab \in P$. This makes P prime. ■

Corollary. *Every maximal ideal is a prime ideal.*

Example 1.17. (1) The prime ideals of \mathbb{Z} are $p\mathbb{Z}$, where p is prime, which are the maximal ideals of \mathbb{Z} .

(2) The ideal (x) in $\mathbb{Z}[x]$ is a prime ideal, but it is not maximal as $(x) \subseteq (2, x) \subseteq \mathbb{Z}[x]$.

1.5 Rings of Fractions.

Lemma 1.5.1. *Let R a commutative ring, and $D \subseteq R$ be nonempty with $0 \notin D$ such that D contains no zero divisors of R and that it is closed under multiplication. Define the relation \sim on $R \times D$ by*

$$(a, b) \sim (c, d) \text{ if, and only if } ad - bc = 0$$

Then \sim is an equivalence relation on $R \times D$.

Proof. We have $ab - ab = 0$ so that $(a, b) \sim (a, b)$. Moreover, if $ad - bc = 0$, then $bc - ad = 0$ so that $(a, b) \sim (c, d)$ implies $(c, d) \sim (a, b)$. Lastly, let $(a, b) \sim (c, d)$ and $(c, d) \sim (e, f)$. Then $ad - bc = 0$ and $cf - ed = 0$, so $af - eb = (ad - bc)f + d(cf - ed) = 0$ so that $(a, b) \sim (e, f)$. ■

Theorem 1.5.2. *Let R a commutative ring, and $D \subseteq R$ be nonempty with $0 \notin D$ such that D contains no zero divisors of R and that it is closed under multiplication. Then there exists a commutative ring Q with identity such that every element of D is a unit of Q .*

Proof. Define the equivalence relation \sim on $R \times D$ by

$$(a, b) \sim (c, d) \text{ if, and only if } ad - bc = 0$$

Label the equivalence classes of \sim over $R \times D$ as $\frac{a}{b} = \{(c, d) \in R \times D : ad - bc = 0\}$. Let

$$Q = R/\sim$$

The factor set of \sim over $R \times D$ and define binary operations $+$ and \cdot by

$$\begin{aligned} \frac{a}{b} + \frac{c}{d} &= \frac{ad + bc}{bd} \\ \frac{a}{b} \cdot \frac{c}{d} &= \frac{ac}{bd} \end{aligned}$$

Suppose that $\frac{a}{b} = \frac{a'}{b'}$ and $\frac{c}{d} = \frac{c'}{d'}$. Then $ab' - a'b = 0$ and $cd' - c'd = 0$. Then

$$\begin{aligned} (ad + bc)(b'd') &= adb'd' + bcb'd' \\ &= ab'dd' + cd'bb' \\ &= a'bdd' + c'dbb' \\ &= (a'd'c'd')bd \end{aligned}$$

So that $+$ is well define. By similar reasoning, \cdot is also well defined.

Now, let $\frac{a}{b}, \frac{c}{d} \in Q$. Then $\frac{a}{b} + \frac{c}{d} = \frac{ad+bc}{bd} \in Q$, as $ad+bc \in R$, and since $b, d \in D$, $bd \in D$. Moreover

$$\frac{a}{b} + \left(\frac{c}{d} + \frac{e}{f}\right) = \frac{a}{b} + \frac{cf+de}{df} = \frac{adf+bcf+bde}{bdf}$$

and

$$\left(\frac{a}{b} + \frac{c}{d}\right) + \frac{e}{f} = \frac{ad-bc}{bd} + \frac{e}{f} = \frac{adf+bcf+bde}{bdf}$$

so that $+$ is associative. Now, take $c = 0$ and $d \in D$, and we have

$$\frac{a}{b} + \frac{0}{d} = \frac{ad}{bd} = \frac{a}{b}$$

Since, $abd - abd = 0$ making $\frac{ad}{bd} = \frac{a}{b}$. Similarly, take $c = -a$ and $d \in D$ and we get

$$\frac{a}{b} + \frac{-a}{d} = \frac{0}{b}$$

So $\frac{0}{d}$ is the identity, and $\frac{-a}{d}$ is the inverse of $\frac{a}{b}$. Lastly, since R is commutative, this makes

$$\frac{a}{b} + \frac{c}{d} = \frac{c}{d} + \frac{a}{b}$$

Now, notice that since D is closed under multiplication, we have

$$\frac{a}{b} \frac{c}{d} = \frac{ac}{bd} \in Q$$

Moreover,

$$\frac{a}{b} \left(\frac{c}{d} \frac{e}{f}\right) = \frac{a}{b} \frac{ce}{df} = \frac{ace}{bdf} = \frac{ac}{bd} \frac{e}{f} = \left(\frac{a}{b} \frac{c}{d}\right) \frac{e}{f}$$

Additionally, since R is commutative, we get

$$\frac{a}{b} \frac{c}{d} = \frac{c}{d} \frac{a}{b}$$

Lastly, take $c = d$ and $d \in D$. Then

$$\frac{a}{b} \frac{d}{d} = \frac{ad}{bd} = \frac{a}{b}$$

This makes Q a commutative ring with identity. Moreover, every element of D is a unit in Q . Moreover $R \subseteq Q$ by taking $r \rightarrow \frac{r}{d}$ for some $d \in D$. ■

Corollary. *The ring Q satisfies the following*

- (1) *Q contains a copy of R as a subring, and every element of Q is of the form rd^{-1} . Moreover, if $D = R \setminus \{0\}$, then Q is a field.*
- (2) *Q is unique and the smallest ring containing a copy of R for which every element of D is a unit.*

Proof. Imbed R into Q first. Define the map $\iota : R \rightarrow Q$ by taking $r \rightarrow \frac{rd}{d}$ where $d \in D$. Notice that $\frac{rd}{d} = \frac{re}{e}$, so that ι is well defined. Now, since $\frac{d}{d}$ is the identity of Q , we have

$$\frac{rsd}{d} = \frac{arsd}{dd} = \frac{rd}{d} \frac{sd}{d}$$

So that ι is a ring homomorphism. Now, since no $d \in D$ is a zero divisor, we have that $\ker \iota = (0)$. This makes ι 1-1. Therefore, by the first isomorphism theorem, we get

$$\iota(R) \simeq R$$

Since $\iota(R) \subseteq Q$, ι is the required imbedding.

Now, if $D = R \setminus \{0\}$, this makes every $r \in R$, nonzero into a unit of Q . Then Q has no zero divisors making it an integral domain, thus Q is a field.

Lastly, let $\phi : R \rightarrow S$ be a 1-1 ring homomorphism such that every $\phi(d)$ is a unit in S , where S is a commutative ring with unit and $d \in D$. Define the map $\Phi : Q \rightarrow S$ by taking $rd^{-1} \rightarrow \phi(rd^{-1})$. Then Φ is a 1-1 ring homomorphism, so that by the first isomorphism theorem,

$$\Phi(Q) \simeq S$$

This makes Q unique. ■

Definition. Let R a commutative ring, and $D \subseteq R$ be nonempty with $0 \notin D$ such that D contains no zero divisors of R and that it is closed under multiplication. Define the equivalence relation \sim on $R \times D$ by

$$(a, b) \sim (c, d) \text{ if, and only if } ad - bc = 0$$

and let

$$Q = R \times D / \sim$$

Then we call the commutative ring Q , with identity $1 = \frac{d}{d}$, the **ring of fractions** of R . If $D = R \setminus \{0\}$ and R is an integral domain, we call Q the **field of fractions**.

Lemma 1.5.3. *If R is an integral domain, and Q its field of fractions, and F is a field containing $R' \simeq R$, then the subfield of F generated by R is isomorphic to Q .*

Proof. Let $\phi : R \rightarrow R'$ the ring isomorphism between R and R' . Then the $\phi : R \rightarrow F$ is 1-1. Define then the map $\Phi : Q \rightarrow F$ by $rd^{-1} \rightarrow \phi(rd^{-1})$. Then Φ is 1-1 and by the first isomorphism theorem, $\Phi(Q) \simeq Q$. Moreover, $\Phi(R) = \phi(R) = R' \subseteq \Phi(Q)$. Now, for all $r, s \in R$, we have $\phi(rs^{-1}) \in \Phi(Q)$ and since every element of Q is of the form rs^{-1} , any subfield containing R' contains $\Phi(Q)$. ■

Example 1.18. (1) The field of fractions of \mathbb{Z} is \mathbb{Q} . Indeed, the construction of the ring of fractions of a commutative ring with identity is inspired by constructing \mathbb{Q} from \mathbb{Z} .

(2) The field of fractions of \mathbb{Q} is \mathbb{Q} itself. In general if F is a field, it is its own field of fractions.

(3) The field of fractions of $\mathbb{Z}[\sqrt{D}]$ is $\mathbb{Q}[\sqrt{D}]$.

(4) $2\mathbb{Z}$ as a subring has no zero divisors, so the field of fractions of $2\mathbb{Z}$ is also \mathbb{Q} .

1.6 Sun Tzu's Theorem.

Definition. Let $\{R_\alpha\}$ a collection of commutative rings with identity. We define the **direct product** of $\{R_\alpha\}$ to be the direct product of $\{R_\alpha\}$ as a group, made into a ring by the operation $(r_\alpha), (s_\alpha) \rightarrow (r_\alpha s_\alpha)$. We write $R = R_1 \times R_2 \times \dots$ when $\{R_\alpha\}$ is a countable collection.

Definition. We call the ideals $A, B \subseteq R$ of a ring R **comaximal** if $A + B = R$.

Example 1.19. If $(m, n) = 1$, then the ideals $n\mathbb{Z}, m\mathbb{Z}$ with $m\mathbb{Z} + n\mathbb{Z} = \mathbb{Z}$ are comaximal. $m\mathbb{Z} + n\mathbb{Z} = \mathbb{Z}$ is the set of all diophantine equations of the form

$$mx + ny = 1$$

Theorem 1.6.1 (Sun-Tzu's Theorem). *Let A_1, \dots, A_k be ideals in a commutative ring R with identity. Then the map*

$$\begin{aligned} R &\rightarrow R/A_1 \times \dots \times R/A_k \\ r &\rightarrow (r + A_1, \dots, r + A_k) \end{aligned}$$

is a ring homomorphism with kernel

$$K = \bigcap_{i=1}^k A_i$$

Moreover if for all $1 \leq i, j \leq k$ with $i \neq j$, A_i and A_j are comaximal, then this map is onto with $\bigcap A_i = \prod A_i$ so that

$$R/\prod_{i=1}^k A_i \simeq R/A_1 \times \dots \times R/A_k$$

Proof. Let $k = 2$, and $A_1 = A$ and $A_2 = B$. Consider the map

$$\begin{aligned} \phi : R &\rightarrow R \\ r &\rightarrow (r + A, r + B) \end{aligned}$$

Then $rs \rightarrow (rs + A, rs + B) = (r + A, r + B)(s + A, s + B)$ so that ϕ is a ring homomorphism. Now, let $r \in \ker \phi$, then $(r + A, r + B) = (A, B)$ so that $r \in A \cap B$, conversely if $r \in A \cap B$ then $r + A = A$ and $r + B = B$ so that $r \in \ker \phi$. Therefore

$$\ker \phi = A \cap B$$

Now, suppose that A and B are comaximal, that is, $A + B = (1)$. Then there is an $x \in A$, and a $y \in B$ such that $x + y = 1$. Then $\phi(x) = (0, 1)$ and $\phi(y) = (1, 0)$ and $x = 1 - y \in 1 + B$. Now, take $r + A, s + B$, then

$$\phi(rx + sy) = \phi(r)\phi(x) + \phi(s)\phi(y) = (r + A, r + B)(0, 1) + (s + A, s + B)(1, 0) = (r + A, s + B)$$

this makes ϕ onto, moreover notice that $AB \subseteq A \cap B$, and if $A + B = (1) = R$, then for every $x \in AB$, $c = c \cdot 1 = cx + cy \in AB$ so that $A \cap B = AB$.

Now, by induction on $k \geq 2$, take $A = A_1$ and $B = A_2 \dots A_k$ by repeating the above argument, we get the result. ■

Corollary. Let $n = p_1^{a_1} \dots p_k^{a_k} \in \mathbb{Z}^+$ be the prime factorization of n , where $p_1 \neq \dots \neq p_k$. Then

$$\mathbb{Z}/n\mathbb{Z} \simeq U(\mathbb{Z}/p_1^{a_1}\mathbb{Z}) \times \dots \times U(\mathbb{Z}/p_k^{a_k}\mathbb{Z})$$

Remark. Sun-Tzu's theorem is most commonly known as the Chinese Remainder theorem, however it is the belief of the author that the name of the theorem should credit the author whenever possible. Also note that the Sun-Tzu of this theorem *is not* the same Sun-Tzu who penned *The Art of War*.

Chapter 2

Domains.

2.1 Eculidian Domains.

Bibliography

- [1] D. Dummit, *Abstract algebra*. Hoboken, NJ: John Wiley & Sons, Inc, 2004.
- [2] I. N. Herstein, *Topics in algebra*. New York: Wiley, 1975.