

Algebraic Geometry.

Alec Zabel-Mena

September 27, 2023

Contents

1	Affine Algebraic Sets	5
1.1	Affine n -Space and Algebraic Sets	5
1.2	Ideals of Algebraic Sets	9
1.3	Hilbert's Basis Theorem	11
1.4	Irreducible Components	12

Chapter 1

Affine Algebraic Sets

1.1 Affine n -Space and Algebraic Sets

Definition. Let k be a field. We define **affine n -space** over k to be the cartesian product $\mathbb{A}^n(k) = \underbrace{k \times \cdots \times k}_{n\text{-times}}$. If the field k is understood, we write \mathbb{A}^n . We call the elements of $\mathbb{A}^n(k)$ **affine points**. We call $\mathbb{A}^1(k)$ and $\mathbb{A}^2(k)$ the **affine line** and **affine plane** over k , respectively.

Definition. Let k be a field, and let $f \in k[x_1, \dots, x_n]$. We call an affine point $P \in \mathbb{A}^n(k)$ a **zero**, or **root** of f if $f(P) = 0$, where $f(P)$ is understood to be $f(a_1, \dots, a_n)$, where $P = (a_1, \dots, a_n)$. We call the set of zeros of f , $V(f)$ the **hypersurface** defined by f . We call hypersurfaces in $\mathbb{A}^2(k)$ **affine plane curves**. If $\deg f = 1$, we call $V(f)$ a **hyperplane**. We call hypersurfaces in $\mathbb{A}^1(k)$ **lines**.

Example 1.1. The following curves in figure 1.1 define algebraic sets.

Definition. Let k be a field, and S any set of polynomials in $k[x_1, \dots, x_n]$. We define the **set of zeros** of S to be the set $V(S) = \{P \in \mathbb{A}^n(k) : f(P) = 0 \text{ for all } f \in S\}$. We call a subset X of $\mathbb{A}^n(k)$ an **affine algebraic set** if $X = V(S)$ for some set S of polynomials.

Lemma 1.1.1. *The following are true for any field k .*

(1) *If \mathfrak{a} is an ideal in $k[x_1, \dots, x_n]$ generated by a set $S \subseteq k[x_1, \dots, x_n]$, then $V(\mathfrak{a}) = V(S)$.*


(2) *If $\{\mathfrak{a}_\alpha\}$ is a collection of ideals of $k[x_1, \dots, x_n]$, then*

$$V\left(\bigcup \mathfrak{a}_\alpha\right) = \bigcap V(\mathfrak{a}_\alpha)$$

(3) *If $\mathfrak{a} \subseteq \mathfrak{b}$ are ideals, then $V(\mathfrak{b}) \subseteq V(\mathfrak{a})$.*

(4) *If $f, g \in k[x_1, \dots, x_n]$, then $V(fg) = V(f) \cup V(g)$.*

(5) *$V(0) = \mathbb{A}^n(k)$ and $V(1) = \emptyset$.*

Figure 1.1: Affine Algebraic Sets in $\mathbb{A}^2(\mathbb{R})$ and $\mathbb{A}^3(\mathbb{R})$.

Proof. First, let S be a set of polynomials in $k[x_1, \dots, x_n]$. Let $\mathfrak{a} = (S)$ the ideal generated by S . Then if $f \in S$ is a polynomial, $f \in I$. Then if $P \in \mathbb{A}^n$ is a zero of f in S , it is a zero of f in \mathfrak{a} , hence $V(S) \subseteq V(\mathfrak{a})$. Conversely, we have that if $f \in \mathfrak{a}$, then by supposition, $f(x_1, \dots, x_n) = f_1(x_1, \dots, x_n) + \dots + f_n(x_1, \dots, x_n) + \dots$. Now, if $f(P) = 0$ in I , then we have $f_i(P) = 0$ for every i . This makes $f(P) = 0$ in S , so that $V(\mathfrak{a}) \subseteq V(S)$.

Now, consider the collection $\{\mathfrak{a}_\alpha\}$ of ideals in $k[x_1, \dots, x_n]$. Let $P \in V(\bigcup \mathfrak{a}_\alpha)$. Then for every $f \in \bigcup \mathfrak{a}_\alpha$, $f(P) = 0$ for each α . So that $P \in \bigcap V(\mathfrak{a}_\alpha)$. Again, on the otherhand, if $P \in \bigcap V(\mathfrak{a}_\alpha)$, $P \in V(\mathfrak{a}_\alpha)$ for all α so that $P \in V(\bigcup \mathfrak{a}_\alpha)$.

Let \mathfrak{a} and \mathfrak{b} ideals in $k[x_1, \dots, x_n]$, where $\mathfrak{a} \subseteq \mathfrak{b}$. Let $P \in V(\mathfrak{b})$. Then for every polynomial $f \in \mathfrak{b}$, $f(P) = 0$, so that $f(P) = 0$ when $f \in \mathfrak{a}$, hence $P \in V(\mathfrak{a})$. This makes $V(\mathfrak{b}) \subseteq V(\mathfrak{a})$.

Consider now the polynomials $f, g \in k[x_1, \dots, x_n]$. Certainly if $P \in V(fg)$ it is a root of fg ; i.e. $fg(P) = 0$. This makes $f(P) = 0$ or $g(P) = 0$ so that $V(fg) \subseteq V(f) \cup V(g)$. On the otherhand if P is a root of f , or a root of g , it is a root of fg making $V(f) \cup V(g) \subseteq V(fg)$, and equality is established.

Finally, observe that the zero polynomial $0(x_1, \dots, x_n)$ has all its coefficients 0, so that any point $P \in \mathbb{A}^n$ is a zero. This makes $V(0) = \mathbb{A}^n$. Likewise, the constant polynomial $1(x_1, \dots, x_n)$ has its 0-th coefficient 1 so that it has not points $P \in \mathbb{A}^n$ as roots. That is $V(1) = \emptyset$. ■

Corollary. *Finite unions of algebraic sets are algebraic.*

Example 1.2. (1) Let k be a field, and consider $\mathbb{A}^1(k)$. Let $f \in k[x]$ be a polynomial of degree n . Then f has at most n roots in k . Now, if \mathfrak{a} is an ideal in k , since k is a PID, we also get $\mathfrak{a} = (f)$ for some $f \in k[x]$. That is $|V(\mathfrak{a})| \leq n$, and so any algebraic set in $\mathbb{A}^1(k)$ is necessarily finite, except, possibly $\mathbb{A}^1(k)$.

(2) Let k be a finite field with p^m elements, where $p, m \in \mathbb{Z}^+$ and p is prime. Then k is the splitting field of the polynomial $f(x_n) = x_n^{p^m} - x_n$ over the finite field \mathbb{F}_p . Suppose then that there is no set S of polynomials in $k[x_1, \dots, x_n]$ for which $X = V(S)$, for some $X \in \mathbb{A}^n(k)$. Choose then a point $P \in X$ and a polynomial $g \in S$. Then we have $g(x_1, \dots, x_n) = g_1(\tilde{X})x_n + \dots + g_n(\tilde{X})x_n$. Notice that if P is a root of f ; i.e. $P \in V(f)$; i.e. $P^{p^m} - P = 0$, then since $P^{p^m} - P$ is a generator for k as a multiplicative group, it generates S . That is, S must contain the point P as a root for g , notice $P^{p^m} = P$ so that $g(P) = g_1(P)P + \dots + g_n(P)P = 0$ in k . This contradicts that $X \neq V(S)$. This makes every set of $\mathbb{A}^n(k)$ algebraic for any finite field.

(3) By the corollary to lemma 1.1.1, we have that finite unions of algebraic sets are algebraic. Now, consider the field \mathbb{Q} , and let $f_q(x) = x + \frac{q}{2}$ in $\mathbb{Q}[x]$. We have that there are $X \subseteq \mathbb{A}^1(\mathbb{Q})$ algebraic, in which $X = V(f_q)$. Notice however, that the polynomial

$$f(x) = \prod_{q \in \mathbb{Q}} f_q(x)$$

has no roots in \mathbb{Q} , as that would imply that for some $n \in \mathbb{Z}^+$, $\sqrt[n]{2} \in \mathbb{Q}$. That is, there is no $X \subseteq \mathbb{A}^1(\mathbb{Q})$ for which $X = V(\prod f_q) = \bigcup V(f_q)$. In general, the countable union of algebraic sets need not be algebraic.

Example 1.3. (1) Let k be a field, and $X = \{(t, t^2, t^3) \in \mathbb{A}^3(k) : t \in k\}$. If k is finite, this is algebraic. Suppose that k is infinite, and consider the polynomial $f(x_1, x_2, x_3) = x_1 + x_2^2 + x_3^3$. Notice that the point $0 \in X$ is a root of f , and that if P is a root of f , then $P \in X$. That is, $X = V(f)$ making X algebraic.

(2) Let $X = \{(\cos t, \sin t) \in \mathbb{A}^2(\mathbb{R}) : t \in \mathbb{R}\}$. Consider the polynomial $f(x, y) = x^2 + y^2 - 1$. Since we have that $\cos^2 t + \sin^2 t = 1$, $X = V(f)$ and X is algebraic.

(3) Let $X = \{(r, \sin t) \in \mathbb{A}^2(\mathbb{R}) : r = \sin t, t \in \mathbb{R}\}$. Consider the polynomial $f(x, y) = x - y$. Then $X = V(f)$.

Lemma 1.1.2. Let k be a field and $C \subseteq \mathbb{A}^2(k)$ an affine plane curve. Let $L \subseteq \mathbb{A}^2(k)$ a line not contained in C . Then C and L intersect at no more than n points; that is, $C \cap L$ is finite with at most n points.

Proof. Let $C = V(f)$ where $f \in k[x, y]$ is a polynomial of degree n , and let $L = V(l)$ where $l(x, y) = y - ax + b$, for some $a, b \in k$. We have that $f(x, y) = f_1(x)y + f_2(x)y^2$. Now, notice that if X, Y is a root of l , then $l(X, Y) = Y - aX + b = 0$, so that $Y = aX + b$. Now, consider a point $P = (X, Y) \in C \cap L = V(f) \cap V(l)$. Then $f(X, Y) = f(X, aX + b) = f_1(X)(aX + b) + f_2(X)(aX + b)^2$. Since f has finitely many roots, there are finitely many $P = (X, Y)$ satisfying $f(X, Y) = 0$. Moreover, f has at most n roots. We finally observe that $C \cap L = V(f(x, ax + b))$. Which shows that $C \cap L$ is finite, and has at most n points. ■

Example 1.4. The following sets are not algebraic.

- (1) $X = \{(x, y) \in \mathbb{A}^2(\mathbb{R}) : y = \sin x\}$. Let L be a line in $\mathbb{A}^2(\mathbb{R})$. Notice then that L intersects X at infinitely many points, so that X cannot be algebraic.
- (2) $X = \{(z, w) \in \mathbb{A}^2(\mathbb{C}) : |z|^2 + |w|^2 = 1\}$, where $|x + iy|^2 = x^2 + y^2$ for all $x, y \in \mathbb{R}$. Let $f(z, w) = |z|^2 + |w|^2 - 1$, and suppose that $X = V(f)$. Let L be a line in $\mathbb{A}^2(\mathbb{C})$. Then $|L \cap X| = 4$; however $\deg f = 2$, so that X cannot be algebraic.
- (3) $X = \{(\cos t, \sin t, t) \in \mathbb{A}^3(\mathbb{R}) : t \in \mathbb{R}\}$. As in (1), there is a line L intersecting X at infinitely many points.

Theorem 1.1.3. *Let k be an algebraically closed field, Then for $n \geq 1$, the complement of an algebraic set is infinite.*

Proof. Observe that since k is algebraically closed, k is infinite, so that $\mathbb{A}^n(k)$ is infinite. Now, suppose $n = 1$, and let $f \in k[x]$ a nonconstant polynomial, and let $X = V(f)$ an algebraic set. Since f has at most finitely many roots, we get $|X|$ is finite, so that $\mathbb{A}^1(k) \setminus X$ is infinite. Moreover since $k[x]$ is a PID, every algebraic set is of the form $X = V(f)$.

Now, suppose that $n > 1$, Let $S \subseteq k[x_1, \dots, x_n]$. Let X be an algebraic set with $X = V(S)$. Then $S = (f_1, \dots, f_m, \dots)$. Now, if $P \in \mathbb{A}^{n-1}(k)$, then each $f_i(P, x_n) \in k[x_n]$ has finitely many roots. So that the polynomial $f_1(P, x_n) + \dots + f_m(P, x_n) + \dots$ has finitely many roots. This makes X finite, and hence $\mathbb{A}^n(k) \setminus X$ is infinite. ■

Corollary. *If $f \in k[x_1, \dots, x_n]$ is nonconstant, then $V(f)$ is infinite.*

Proof. consider $f \in k[x_1, \dots, x_n]$ nonconstant. Observe that

$$f(x_1, \dots, x_n) = \sum f_i(x_1, \dots, x_{n-1})x_n^i$$

Where $f_i \in k[x_1, \dots, x_{n-1}]$. Now, suppose that $P = (a_1, \dots, a_{n-1})$, then

$$f(P, x_n) = \sum f_i(a_1, \dots, a_{n-1})x_n^i$$

has at most n roots in $k[x_n]$. However, notice that since $\mathbb{A}^n(k)$ is infinite, there are infinitely many choices for P , so that if $Q = (P, a_n)$ is a root of f , then f has infinitely many roots. That is, $V(f)$ is infinite. ■

Lemma 1.1.4. *Let k be a field, and let $X \subseteq \mathbb{A}^n(k)$ and $Y \subseteq \mathbb{A}^m(k)$ algebraic sets. Then $X \times Y$ is an algebraic set in $\mathbb{A}^{n+m}(k)$.*

Proof. Since $\mathbb{A}^m(k)$ and $\mathbb{A}^n(k)$ are cartesian products, we have that $\mathbb{A}^m(k) \times \mathbb{A}^n(k) = \mathbb{A}^{m+n}(k)$. Then $X \times Y = (X, Y)$. Now, let $S \subseteq k[x_1, \dots, x_m]$ and $T \subseteq k[x_1, \dots, x_n]$ such that $X = V(S)$ and $Y = V(T)$. Let $P \in X \times Y$, then $P = (A, B)$ where $A = (a_1, \dots, a_m)$ and $B = (b_1, \dots, b_n)$. Let $f = f_1 + \dots + f_d + \dots \in S$ and $g = g_1 + \dots + g_l \in T$. Consider then $f \times g((x_1, \dots, x_m), (y_1, \dots, y_n)) = f(x_1, \dots, x_m)g(y_1, \dots, y_n)$. Since $f(A) = 0$ and $g(B) = 0$, then $f \times g(P) = f(A)g(B) = 0$ so that $P \in V(f) \times V(g)$. Conversely, let $P \in V(f) \times V(g)$. Then $P = (A, B)$ where $A \in \mathbb{A}^m(k)$ and $B \in \mathbb{A}^n(k)$, and $f \times g(P) = f(A)g(B) = 0$. Since $A \in V(f)$ and $B \in V(g)$, we get $f(A) = 0$ and $g(B) = 0$, so that $P \in X \times Y$. This makes $X \times Y = V(f) \times V(g)$. ■

1.2 Ideals of Algebraic Sets

Lemma 1.2.1. *Let k be a field, and $X \subseteq \mathbb{A}^n(k)$. Consider the set $I(X) = \{f \in k[x_1, \dots, x_n] : f(P) = 0 \text{ for all } P \in X\}$. Then $I(X)$ forms an ideal of $k[x_1, \dots, x_n]$.*

Proof. Let $f, g \in I(X)$. Then for all $P \in X$, $f(P) = 0$, and $g(P) = 0$, so that $f + g(P) = f(P) + g(P) = 0$. Moreover, $-f(P) = 0$ as well. So I is a subgroup of $k[x_1, \dots, x_n]$ under addition. Now, take $f \in I(X)$ and $g \in k[x_1, \dots, x_n]$. Then $fg(P) = 0$ for all $P \in X$ which makes $I(X)$ into an ideal. ■

Definition. Let k be a field and $X \subseteq \mathbb{A}^n(k)$. We define the **ideal** of X to be the ideal $I(X) = \{f \in k[x_1, \dots, x_n] : f(P) = 0 \text{ for all } P \in X\}$ of $k[x_1, \dots, x_n]$.

Lemma 1.2.2. *Let k be a field. The following are true for all $X, Y \subseteq \mathbb{A}^n(k)$ and for all $S \subseteq k[x_1, \dots, x_n]$.*

- (1) *If $X \subseteq Y$, then $I(Y) \subseteq I(X)$.*
- (2) *$I(\emptyset) = k[x_1, \dots, x_n]$ and $I(\mathbb{A}^n(k)) = (0)$.*
- (3) *$S \subseteq I(V(S))$ and $X \subseteq V(I(X))$.*
- (4) *$V(I(V(S))) = V(S)$ and $V(I(V(I(X)))) = I(X)$.*

Proof. Let $X, Y \subseteq \mathbb{A}^n(k)$, with $X \subseteq Y$. Let $f \in I(Y)$, then for all $P \in Y$, $f(P) = 0$. Now, since $P \in X$, we get for all $P \in X$ $f(P) = 0$ so that $f \in I(X)$.

Observe now that the polynomial $1(x_1, \dots, x_n) = 1$ has no points in $\mathbb{A}^n(k)$ as roots, so that $I(\emptyset) = k[x_1, \dots, x_n]$. Likewise, for the polynomial $0(x_1, \dots, x_n) = 0$, every point in $\mathbb{A}^n(k)$ is a root, so that $I(\mathbb{A}^n(k)) = (0)$.

For the third assertion, let $S \subseteq k[x_1, \dots, x_n]$. If $f \in V(S)$, then for every $P \in V(S)$, $f(P) = 0$, by definition. This makes $S \subseteq I(V(S))$. Likewise, if $X \subseteq \mathbb{A}^n(k)$ and $P \in X$, then for all $f \in I(X)$, $f(P) = 0$, so that $P \in V(I(X))$.

Lastly, let $P \in V(S)$, and $f \in I(V(S))$. By definition, $f(P) = 0$ so that $V(S) \subseteq V(I(V(S)))$. Conversely, let $P \in V(I(V(S)))$ then for every $f \in I(V(S))$, $f(P) = 0$, which puts $P \in V(S)$ so that $V(I(V(S))) \subseteq V(S)$. Likewise, by similar reasoning we conclude that $V(I(V(I(X)))) = I(X)$. ■

Corollary. *If k is an infinite field, then for any $a_1, \dots, a_n \in k$, $I(a_1, \dots, a_n) = (x_1 - a_1, \dots, x_n - a_n)$.*

Proof. Let $f \in I(a_1, \dots, a_n)$. Since k is infinite, and $f(a_1, \dots, a_n) = 0$,

$$f(x_1, \dots, x_n) = \sum g_i(x_1, \dots, x_n)(x_i - a_i)$$

so $f \in (x_1 - a_1, \dots, x_n - a_n)$. Conversely, if $f \in (x_1 - a_1, \dots, x_n - a_n)$, we observe that $f \in I(a_1, \dots, a_n)$. ■

Definition. Let \mathfrak{a} be an ideal of a ring R . We define the **radical** of \mathfrak{a} to be the set

$$\text{Rad } \mathfrak{a} = \{a \in R : a^n \in \mathfrak{a}, \text{ for some } n \in \mathbb{Z}^+\}$$

We call I a **radical ideal** if $I = \text{Rad } I$.

Lemma 1.2.3. *Let R be a ring, and \mathfrak{a} an ideal of R . Then $\text{Rad } \mathfrak{a}$ is also an ideal of R .*

Proof. Let $a, b \in \text{Rad } \mathfrak{a}$, then $a^m \in \mathfrak{a}$ and $b^n \in \mathfrak{a}$ for some $m, n \in \mathbb{Z}^+$. Now, observe that

$$(a + b)^{m+n} = a^{m+n} + \sum_{i=1}^{m+n-2} \binom{m+n}{i} a^i b^{m+n-i} + b^{m+n}$$

Now, $a^{m+n} = a^m a^n \in \mathfrak{a}$ and $b^{m+n} = b^n b^m \in \mathfrak{a}$ by the ideal properties of \mathfrak{a} . Moreover, notice if $i \geq n$, then $a^i b^{m+n-i} \in \mathfrak{a}$; on the otherhand, if $m \leq m+n-i$, then $a^i b^{m+n-i} \in \mathfrak{a}$. This makes each $a^i b^{m+n-i} \in \mathfrak{a}$, and that $(a+b)^{m+n} \in \mathfrak{a}$. Also observe that if $a^n \in \mathfrak{a}$, then $(-a)^n = -(a^n) \in \mathfrak{a}$. So that $\text{Rad } \mathfrak{a}$ is an additive subgroup of R .

Lastly, suppose that if $a \in \text{Rad } R$, and $r \in R$, then we have $(ra)^n = r^n a^n \in \mathfrak{a}$ for some $n \in \mathbb{Z}^+$. Thus $ra \in \text{Rad } \mathfrak{a}$. This makes $\text{Rad } \mathfrak{a}$ an ideal of R . ■

Corollary. *$\text{Rad } \mathfrak{a}$ is a radical ideal of R .*

Proof. Observe that $\text{Rad } \mathfrak{a} \subseteq \text{Rad}(\text{Rad } \mathfrak{a})$. Now, let $a \in \text{Rad}(\text{Rad } \mathfrak{a})$, then $a^n \in \text{Rad } \mathfrak{a}$ for some $n \in \mathbb{Z}^+$, so that $(a^n)^m = a^{mn} \in \mathfrak{a}$ for some $m \in \mathbb{Z}^+$. This makes $a \in \text{Rad } \mathfrak{a}$. So $\text{Rad}(\text{Rad } \mathfrak{a}) \subseteq \text{Rad } \mathfrak{a}$. This makes $\text{Rad } \mathfrak{a}$ radical. ■

Lemma 1.2.4. *Any prime ideal in a ring R is radical.*

Proof. Let \mathfrak{p} be a prime ideal. We have that $\mathfrak{p} \subseteq \text{Rad } \mathfrak{p}$. Now, let $a \in \text{Rad } \mathfrak{p}$. Then for some $n \in \mathbb{Z}^+$, we have that $a^n \in \mathfrak{p}$. Since \mathfrak{p} is prime, either $a \in \mathfrak{p}$ or $a^{n-1} \in \mathfrak{p}$. If $a \in \mathfrak{p}$, we are done; otherwise we have $a^{n-1} = aa^{n-2} \in \mathfrak{p}$. Repeating this process recursively, we obtain that $a \in \mathfrak{p}$, so that $\mathfrak{p} = \text{Rad } \mathfrak{p}$. ■

Lemma 1.2.5. *Let k be a field, then for any $X \subseteq \mathbb{A}^n(k)$, $I(X)$ is a radical ideal.*

Proof. For any $f \in I(X)$, notice that $f^n(P) = f(f^{n-1}(P)) = \cdots = \underbrace{f(f(\cdots(P)))}_{n \text{ times}}$ ■

Example 1.5. Observe that $\mathbb{R}[x]/(x^2 + 1) \simeq \mathbb{C}$ is a field, so that $(x^2 + 1)$ is a maximal ideal, hence a prime ideal, and hence, a radical ideal. Observe also that $V(x^2 + 1) = \emptyset$, so that $I(V(x^2 + 1)) = \mathbb{R}[x]$. Therefore, $(x^2 + 1)$ is not the ideal of any nonempty set of $\mathbb{A}^1(\mathbb{R})$.

Lemma 1.2.6. *If X and Y are algebraic sets in $\mathbb{A}^n(k)$, then $I(X) = I(Y)$ if, and only if $X = Y$.*

Proof. If $X = Y$, then we can observe that $I(X) = I(Y)$. Conversely, suppose that $I(X) = I(Y)$, and let $f \in I(X)$. Then for all $P \in X$, we have $f(P) = 0$. Since $I(X) = I(Y)$, we must have that $P \in Y$ so that $X \subseteq Y$. In similar fashion, we get that $Y \subseteq X$. ■

Theorem 1.2.7. *Let k be a field. The ideal $(x_1 - a_1, \dots, x_n - a_n)$ of $k[x_1, \dots, x_n]$ is a maximal ideal of $k[x_1, \dots, x_n]$ and the natural map*

$$k \rightarrow k[x_1, \dots, x_n]/(x_1 - a_1, \dots, x_n - a_n)$$

defines an isomorphism.

Proof. Define the map $\phi : k[x_1, \dots, x_n] \rightarrow k$ defined by the rule $f(x_1, \dots, x_n) \rightarrow f(a_1, \dots, a_n)$ where $a_1, \dots, a_n \in k$. Then notice that $\ker \phi = (x_1 - a_1, \dots, x_n - a_n)$. Now, consider $f(x_1, \dots, x_n) = 1 + 0x_1 + \dots + 0x_n \in k[x_1, \dots, x_n]$. Then $f(a_1, \dots, a_n) = 1 + 0a_1 + \dots + 0a_n = 1 \in \phi(k[x_1, \dots, x_n])$. So that ϕ is onto. By the first isomorphism theorem for ring homomorphisms, we get

$$k[x_1, \dots, x_n] / (x_1 - a_1, \dots, x_n - a_n) \simeq k$$

So that $(x_1 - a_1, \dots, x_n - a_n)$ is a maximal ideal. Notice also that $\Phi = \pi \circ \phi$ where $\pi : k \rightarrow k[x_1, \dots, x_n] / (x_1 - a_1, \dots, x_n - a_n)$ is the natural map. So π defines the isomorphism. ■

1.3 Hilbert's Basis Theorem

Definition. We call a ring in which every ideal is finitely generated a **Noetherian ring**.

Theorem 1.3.1 (Hilbert's Basis Theorem). *If R is a Noetherian ring, then so is the polynomial ring $R[x]$.*

Proof. Let \mathfrak{a} be an ideal of $R[x]$, and let L be the set of all leading coefficients of polynomials in \mathfrak{a} . Notice that since $0 \in \mathfrak{a}$, then $0 \in L$, so that L is nonempty. Moreover, let $f(x) = ax^d + \dots$ and $g(x) = bx^e + \dots$ polynomials in \mathfrak{a} of degree $\deg f = d$ and $\deg g = e$, with leading coefficients $a, b \in R$. Then for any $r \in R$, we have the coefficient $ra - b = 0$, or $ra - b$ is the leading coefficient of the polynomial $rx^e f - x^d g \in \mathfrak{a}$. In either case, we get $ra - b \in L$. This makes L an ideal of R . Now, since R is Noetherian L is finitely generated; let $L = (a_1, \dots, a_n)$. Then for every $1 \leq i \leq n$, let $f_i \in \mathfrak{a}$ the polynomial of degree $\deg f_i = e_i$ whose leading coefficient is a_i . Take, then $N = \max \{e_1, \dots, e_n\}$. Then for any $d \in \mathbb{Z}/N\mathbb{Z}$, let L_d be the set of all leading coefficients of polynomials in \mathfrak{a} , of degree d , together with 0. Let $f_{di} \in \mathfrak{a}$ a polynomial of degree $\deg f_{di} = d$ with leading coefficient b_{di} . We wish to show that

$$\mathfrak{a} = (f_1, \dots, f_n) \cup (f_{d1}, \dots, f_{dn})$$

Let $\mathfrak{a}' = (f_1, \dots, f_n) \cup (f_{d1}, \dots, f_{dn})$. By construction, since the generators were chosen from \mathfrak{a} , $\mathfrak{a}' \subseteq \mathfrak{a}$. Now, if $\mathfrak{a} \neq \mathfrak{a}'$. Then there is a nonzero polynomial $f \in \mathfrak{a}$ of minimum degree not contained in \mathfrak{a}' (i.e. $f \notin \mathfrak{a}'$). Let $\deg f = d$, and let a be the leading coefficient of f . Suppose that $d \geq N$. Since $a \in L$, a is an R -linear combination of the generators of L ; i.e.

$$a = r_1 a_1 + \dots + r_n a_n$$

where $r_1, \dots, r_n \in R$. Let

$$g = r_1 x^{d-e_1} f_1 + \dots + r_n x^{d-e_n} f_n$$

then $g \in \mathfrak{a}'$ and has degree $\deg g = d$ and leading coefficient a . Hence $f - g \in \mathfrak{a}'$ is of smaller degree, and by the minimality of f , $f - g = 0$, which makes $f = g \in \mathfrak{a}'$; a contradiction. Therefore $\mathfrak{a} = \mathfrak{a}'$

Now, if $d < N$, then $a \in L_d$, and so is an R -linear combination of generators of L_d ; that is

$$a = r_1 b_{d1} + \dots + r_n b_{dn}$$

where $r_1, \dots, r_n \in R$. Then let

$$g = r_1 f_{d1} + \dots + r_n f_{dn}$$

then $g \in \mathfrak{a}'$ is a polynomial of degree $\deg g = d$ and leading coefficient a ; which gives us the above contradiction.

Therefore, $\mathfrak{a} = \mathfrak{a}'$, and since \mathfrak{a}' is finitely generated, $R[x]$ is Noetherian. \blacksquare

Corollary. *Let k be a field. Then the polynomial ring in n variables $k[x_1, \dots, x_n]$ is Noetherian.*

Theorem 1.3.2. *Every algebraic set is the intersection of a finite number of hypersurfaces.*

Proof. Let \mathfrak{a} be an ideal in the ring $k[x_1, \dots, x_n]$ for some field k , and consider the set $V(\mathfrak{a})$. Since $k[x_1, \dots, x_n]$ is Noetherian, then $\mathfrak{a} = (f_1, \dots, f_n)$, so that

$$V(\mathfrak{a}) = V(f_1) \cap \dots \cap V(f_n)$$

\blacksquare

Theorem 1.3.3. *Let \mathfrak{a} be an ideal in a ring R , and consider the natural map $\pi : R \rightarrow R/\mathfrak{a}$. The following are true.*

- (1) *For every ideal \mathfrak{b}' of R/\mathfrak{a} , $\pi^{-1}(\mathfrak{b}') = \mathfrak{b}$ is an ideal of R containing \mathfrak{a} . Moreover, for any ideal \mathfrak{b} of R containing \mathfrak{a} , then $\pi(\mathfrak{b}) = \mathfrak{b}'$.*
- (2) *The ideal \mathfrak{b}' of R/\mathfrak{a} is a radical ideal if, and only if \mathfrak{b} is a radical ideal in R .*
- (3) *If \mathfrak{b} is finitely generated in R , then \mathfrak{b}' is finitely generated in R/\mathfrak{a} . Moreover, R/\mathfrak{a} is Noetherian if R is Noetherian.*

Proof. Let \mathfrak{b}' be an ideal of R/\mathfrak{a} . Since the natural map π is onto, there is an ideal $\mathfrak{b} \in R$ for which $\mathfrak{b} = \pi^{-1}(\mathfrak{b}')$. Now, let $a, b \in \mathfrak{b}$, then $\pi(a), \pi(b) \in \mathfrak{b}'$, so that $\pi(a + b) \in \mathfrak{b}'$ and $-\pi(a) \in \mathfrak{b}'$. Moreover, if $a \in \mathfrak{b}$, and $r \in R$, then $r\pi(a) = \pi(ra) \in \mathfrak{b}'$, since \mathfrak{b}' is an ideal. Now, since $\ker \pi = \mathfrak{a}$, we have that $\mathfrak{a} \subseteq \mathfrak{b}$. So that \mathfrak{b} is an ideal containing \mathfrak{a} . By similar reasoning, if \mathfrak{b} is an ideal containing \mathfrak{a} , then $\mathfrak{b}' = \pi(\mathfrak{b})$ is also an ideal.

Now, suppose that \mathfrak{b} is a radical ideal. That is, $\mathfrak{b} = \text{Rad } \mathfrak{b}$. Since $\mathfrak{b} = \pi^{-1}(\mathfrak{b}')$, we have $\pi^{-1}(\mathfrak{b}') = \text{Rad } \pi^{-1}(\mathfrak{b}')$. Now, suppose that \mathfrak{b} is a prime ideal, then if $ab \in \mathfrak{b}$, either $a \in \mathfrak{b}$ or $b \in \mathfrak{b}$. This implies whenever $\pi(ab) \in \mathfrak{b}'$, either $\pi(a) \in \mathfrak{b}'$ or $\pi(b) \in \mathfrak{b}'$. This makes \mathfrak{b}' prime. Similarly, if \mathfrak{b}' is prime so is \mathfrak{b} . Finally, by definition of a maximal ideal, \mathfrak{b} is maximal if, and only if \mathfrak{b}' is maximal.

Finally, suppose that \mathfrak{b} is finitely generated, then $\mathfrak{b} = (a_1, \dots, a_n) = \pi^{-1}(\mathfrak{b}')$ for $a_1, \dots, a_n \in R$. Then every element of \mathfrak{b} is the sum of a_1, \dots, a_n . That is, $b = r_1 a_1 + \dots + r_n a_n$ for every $b \in \mathfrak{b}$, and $r_1, \dots, r_n \in R$. Now, since $b \in \mathfrak{b} = \pi^{-1}(\mathfrak{b}')$, then $\pi(b) = r_1 \pi(a_1) + \dots + r_n \pi(a_n) \in \mathfrak{b}'$, so that $\mathfrak{b}' = (\pi(a_1), \dots, \pi(a_n))$. \blacksquare

Corollary. *Let k be a field and \mathfrak{a} an ideal of $k[x_1, \dots, x_n]$. Any ring of the form $k[x_1, \dots, x_n]/\mathfrak{a}$ is a Noetherian ring.*

1.4 Irreducible Components

Bibliography

- [1] D. Dummit, *Abstract algebra*. Hoboken, NJ: John Wiley & Sons, Inc, 2004.
- [2] I. N. Herstein, *Topics in algebra*. New York: Wiley, 1975.
- [3] M. Atiyah and I. MacDonald, *Introduction to Commutative Algebra*. Addison-Wesley Series in Mathematics, CRC Press.
- [4] D. Eisenbud, *Commutative Algebra: Wit a View Toward Algebraic Geometry*. Graduate Texts in Mathematics, Springer Verlag.
- [5] R. Hartshorne, *Algebraic Geometry*. Graduate Texts in Mathematics, Springer Verlag.
- [6] W. Fulton, *Algebraic Curves: An Introduction to ALgebraic Geometry*. Advanced Book Classics, Addison-Wesley.