

Topology

Alec Zabel-Mena

June 28, 2022

Chapter 1

Topological Spaces and Continuous Functions.

1.1 Topological Spaces.

Definition. A **topology** on a set X is a collection \mathcal{T} of subsets of X such that:

- (1) $\emptyset \in \mathcal{T}$ and $X \in \mathcal{T}$
- (2) For any collection $\{U_\alpha\}$ of subsets of X , $\bigcup_\alpha U_\alpha \in \mathcal{T}$.
- (3) For any finite collection $\{U_i\}_{i=1}^n$ of subsets of X , $\bigcap_{i=1}^n U_i \in \mathcal{T}$.

We call the pair (X, \mathcal{T}) a **topological space**, and we call the elements of \mathcal{T} **open sets**.

Example 1.1. (1) Let X be any set, the collection of all subsets of X , 2^X is a topology on X , which we call the **discrete topology**. We call the topology $\mathcal{T} = \{\emptyset, X\}$ the **indiscrete topology**.

- (2) The set of three points $\{a, b, c\}$ has the 9 following topologies in figure 1.1.




Figure 1.1: The Topologies on $\{a, b, c\}$.

- (3) Let X be any set, and let $\mathcal{T}_f = \{U \subseteq X : X \setminus U \text{ is finite, or } X \setminus U = X\}$. Then \mathcal{T}_f is a topology and called the **finite complement topology**.
- (4) Let X be any set, and let $\mathcal{T}_c = \{U \subseteq X : X \setminus U \text{ is countable, or } X \setminus U = X\}$. Then \mathcal{T}_c is a topology on X called the **countable complement topology**.

Definition. Let X be a set, and let \mathcal{T} and \mathcal{T}' be topologies on X . We say that \mathcal{T} is **coarser** than \mathcal{T}' , and \mathcal{T}' **finer** than \mathcal{T} if $\mathcal{T} \subseteq \mathcal{T}'$. If two topologies are either coarser, or finer than each other, we call them **comparable**.

Example 1.2. The topologies \mathcal{T}_f and \mathcal{T}_c are comparable, and we see that $\mathcal{T}_f \subseteq \mathcal{T}_c$, so \mathcal{T}_f is coarser than \mathcal{T}_c , and \mathcal{T}_c is finer than \mathcal{T}_f .

1.2 The Basis and Subbasis for a Topology.

Definition. If X is a set, the **basis** for a topology on X is a collection \mathcal{B} of subsets of X , called **basis elements**, such that:

- (1) For every $x \in X$, there is a $B \in \mathcal{B}$ such that $x \in B$.
- (2) For $B_1, B_2 \in \mathcal{B}$, if $x \in B_1 \cap B_2$, then there is a $B_3 \in \mathcal{B}$ such that $x \in B_3 \subseteq B_1 \cap B_2$.

We define the topology \mathcal{T} **generated** by \mathcal{B} to be collection of open sets: $\mathcal{T} = \{U \subseteq X : \text{for all } x \in U, \text{ there exists a } B \in \mathcal{B} \text{ such that } x \in B\}$.

Theorem 1.2.1. *Let X be a set, and \mathcal{B} a basis of X , then the collection of subsets of X , $\mathcal{T} = \{U \subseteq X : \text{for all } x \in U, \text{ there exists a } B \in \mathcal{B} \text{ such that } x \in B\}$ is a topology on X .*


Proof. Let \mathcal{B} be a basis for a topology in X , and consider \mathcal{T} as defined above. Clearly, $\emptyset \in \mathcal{T}$ and so is X .

Now let $\{U_\alpha\}$ be a collection of subsets of X , and let $U = \bigcup U_\alpha$. Then if $x \in U$ for some α , there is a B_α such that $x \in B_\alpha \subseteq U_\alpha$, thus $x \in B_\alpha \subseteq U$.

Now let $x \in U_1 \cap U_2$, and choose $B_1, B_2 \in \mathcal{B}$ such that $x \in B_1 \subseteq U_1$ and $x \in B_2 \subseteq U_2$. Then by definition, there is a B_3 for which $x \in B_3 \subseteq B_1 \cap B_2$. Now suppose for arbitrary n , that $U = \bigcap_{i=1}^n U_i \in \mathcal{T}$, for some finite collection $\{U_i\}$ of subsets of X . Then by let $B_n, B_{n+1} \in \mathcal{B}$ such that $x \in B_n \subseteq U$ and $x \in B_{n+1} \subseteq U_{n+1}$. Then by our hypothesis, there is a B for which $x \in B \subseteq B_n \cap B_{n+1}$, thus $U \cap U_{n+1} = \bigcap_{i=1}^{n+1} U_i \in \mathcal{T}$. This make \mathcal{T} a topology on X . ■

Example 1.3. (1) Let \mathcal{B} be the set of all circular regions in the plane $\mathbb{R} \times \mathbb{R}$, then \mathcal{B} satisfies the conditions needed for a basis.

- (2) The collection \mathcal{B}' in $\mathbb{R} \times \mathbb{R}$ of all rectangular region also forms a basis for a topology on $\mathbb{R} \times \mathbb{R}$.
- (3) For any set X , the set of all 1-point subsets of X forms a basis for a topology on X .

Figure 1.2: The basis for \mathcal{B} and \mathcal{B}' in $\mathbb{R} \times \mathbb{R}$ (see example (2)).

Lemma 1.2.2. *Let X be a set, and \mathcal{B} be a basis for a topology \mathcal{T} on X . Then $\mathcal{T} = \{\bigcup B : B \in \mathcal{B}\}$.*

Proof. Given a collection $\{B_i\}_{i=1}^{\infty}$ of basis elements in \mathcal{B} , since they are all in \mathcal{T} , their unions are also in \mathcal{T} . Conversely, given $U \in \mathcal{T}$, then for every point $x \in U$, choose a $B_x \in \mathcal{B}_x$ such that $x \in B_x \subseteq U$, then $U = \bigcup_{x \in U} B_x$. ■

Lemma 1.2.3. *Let (X, \mathcal{T}) be a topological space, and let $\mathcal{C} \subseteq \mathcal{T}$ be a collection of open sets of X such that for every $x \in U$, there is a $C \in \mathcal{C}$ such that $x \in C \subseteq U$. Then \mathcal{C} is the basis for a \mathcal{T} on X .*

Proof. Take any $x \in X$, then there is a $C \in \mathcal{C}$ such that $x \in C \subseteq U$, thus the first condition for a basis is satisfied. Now let $x \in C_1 \cap C_2$ for $C_1, C_2 \in \mathcal{C}$, since $C_1 \cap C_2$ is open in X , there is a $C_3 \in \mathcal{C}$ such that $x \in C_3 \subseteq C_1 \cap C_2$. Therefore \mathcal{C} is a basis for a topology on X .

Now let $\mathcal{T}_{\mathcal{C}}$ be the topology generated by \mathcal{C} , now for $U \in \mathcal{T}$, we have by the hypothesis, that $U \in \mathcal{T}_{\mathcal{C}}$; and by lemma 1.2.2, $W \in \mathcal{T}_{\mathcal{C}}$ is the union of elements of \mathcal{C} , which is a subcollection of \mathcal{T} , thus $W \in \mathcal{T}$. Therefore $\mathcal{T}_{\mathcal{C}} = \mathcal{T}$. ■

Lemma 1.2.4. *Let \mathcal{B} and \mathcal{B}' be bases for topologies \mathcal{T} and \mathcal{T}' on X . Then the $\mathcal{T} \subseteq \mathcal{T}'$ if and only if for all $x \in X$, and all $B \in \mathcal{B}$, there is a $B' \in \mathcal{B}'$ such that $x \in B' \subseteq B$.*

Proof. Suppose first that $\mathcal{T} \subseteq \mathcal{T}'$, and let $x \in X$, and choose $B \in \mathcal{B}$ such that $x \in B$, then B is open in \mathcal{T} , thus it is open in \mathcal{T}' , thus there is a $B' \in \mathcal{B}'$ such that $x \in B' \subseteq B$. Conversely, suppose there is a $B' \in \mathcal{B}'$ for which $x \in B' \subseteq B$ for all $x \in X$, $B \in \mathcal{B}$. Take $x \in U \in \mathcal{T}$, since \mathcal{B} generates \mathcal{T} , $x \in B \subseteq U$, since $B' \subseteq B$, this implies that $U \in \mathcal{T}'$ and $\mathcal{T} \subseteq \mathcal{T}'$. ■

Definition. If \mathcal{B} is the collection of open intervals (a, b) in \mathbb{R} , we call the topology generated by \mathcal{B} the **standard topology** on \mathbb{R} , and we denote it simply by \mathbb{R} .

Definition. If \mathcal{B} is the collection of half open intervals $[a, b)$ in \mathbb{R} , we call the topology generated by \mathcal{B} the **lower limit topology** on \mathbb{R} , and we denote it simply by \mathbb{R}_l . If \mathcal{B}' is the collection of all half open intervals $(a, b]$ in \mathbb{R} , then we call the topology generated by \mathcal{B}' the **upper limit topology** on \mathbb{R} , and denote it \mathbb{R}_L .

Definition. If \mathcal{B} is the collection of all open intervals of the form $(a, b) \setminus \frac{1}{\mathbb{Z}^+}$, where $\frac{1}{\mathbb{Z}^+} = \{\frac{1}{n} : n \in \mathbb{Z}^+\}$, we call the topology generated by \mathcal{B} the $\frac{1}{\mathbb{Z}^+}$ -**topology** on \mathbb{R} , and we denote it $\mathbb{R}_{\frac{1}{\mathbb{Z}^+}}$.

Lemma 1.2.5. *The topologies \mathbb{R}_l , \mathbb{R}_L , and $\mathbb{R}_{\frac{1}{\mathbb{Z}^+}}$ are all strictly finer than \mathbb{R} , but are not comparable with each other.*

Proof. Let (a, b) be a basis element for \mathbb{R} , and let $x \in (a, b)$, the basis element $[x, b) \in \mathbb{R}_l$ lies in (a, b) and contains x , however, there can be no interval (a, b) in $[x, b)$ as $x \leq a$, thus $\mathbb{R} \subset \mathbb{R}_l$; a similar argument holds for \mathbb{R}_L .

Similarly, for $(a, b) \in \mathbb{R}$, the basis element $(a, b) \setminus \frac{1}{\mathbb{Z}^+}$ of $\mathbb{R}_{\frac{1}{\mathbb{Z}^+}}$ lies in (a, b) , however, choose the basis $B = (-1, 1) \setminus \frac{1}{\mathbb{Z}^+}$, and choose $0 \in B$, since \mathbb{Z}^+ is dense in \mathbb{R} , there is no interval (a, b) containing 0 and lying in B , thus $\mathbb{R} \subseteq \mathbb{R}_{\frac{1}{\mathbb{Z}^+}}$.

Now choose $[0, 1)$ in \mathbb{R}_l , and choose $\frac{1}{k} \in [0, 1)$ such that $k \in \mathbb{Z}^+$. Now $(0, 1) \subseteq [0, 1)$, so we cannot say that $[0, 1)$ is a basis for \mathbb{R} , and moreover, $[0, 1) \setminus \frac{1}{\mathbb{Z}^+}$ cannot be said to be a basis in $\mathbb{R}_{\frac{1}{\mathbb{Z}^+}}$, thus \mathbb{R}_l and $\mathbb{R}_{\frac{1}{\mathbb{Z}^+}}$ are incomparable, a similar argument holds for \mathbb{R}_L .

Lastly, let (a, b) be in \mathbb{R} and choose $x \in (a, b)$. Then $(a, x]$ and $[x, b)$ are both in (a, b) , however it is clear that $(a, x]$ and $[x, b)$ cannot be contained in each other, thus \mathbb{R}_l and \mathbb{R}_L are incomparable. ■

Definition. A **subbasis**, \mathcal{S} , for a topology on X is a collection of subsets of X whose union equals X . We call the **topology generated by \mathcal{S}** to be the collection of all unions of finite intersections of elements of \mathcal{S} , that is:

$$\mathcal{T} = \left\{ \bigcup_{i=1}^n S_i : S_i \in \mathcal{S} \text{ for } 1 \leq i \leq n \right\}$$

Theorem 1.2.6. Let \mathcal{S} be a subbasis for a topology on X . Then the collection $\mathcal{T} = \left\{ \bigcup_{i=1}^n S_i : S_i \in \mathcal{S} \text{ for } 1 \leq i \leq n \right\}$ is a topology on X .

Proof. It is sufficient to show that the collection \mathcal{B} of all finite intersections of elements of \mathcal{S} is a basis for a topology on X . By lemma 1.2.1, for $x \in X$, it belongs to an element S of \mathcal{S} , and therefore, to an element of \mathcal{B} . Now let $B_1 = \bigcap_{i=1}^m S_i$ and $B_2 = \bigcap_{j=1}^n S'_j$ be basis elements of \mathcal{B} . The intersection $B_1 \cap B_2$ is a finite intersection of elements of \mathcal{S} , and hence also belongs in \mathcal{B} , and hence we can take another basis element B_3 such that $x \in B_3 \subseteq B_1 \cap B_2$. ■

1.3 The Order Topology.

Definition. Let X be a set with a simple order relation, and suppose that $|X| > 1$. Let \mathcal{B} be the collection of sets of the following forms:

- (1) All open intervals $(a, b) \in X$.
- (2) All half open intervals $[a_0, b)$ where a_0 is the least element (if any) of X .
- (3) All half open intervals of the form $(a, b_0]$ where b_0 is the greatest element (if any) of X .

Then \mathcal{B} forms the basis for a topology on X called the **order topology**

Theorem 1.3.1. The collection \mathcal{B} forms a basis.

Proof. Consider $x \in X$, if x is the least element of X , then it lies in all intervals of type (2), if it is the largest, then it lies in all intervals of type (3). If x is neither the least nor largest element, then $x \in (a_0, b_0)$ with a_0 and b_0 the least and largest elements (if any) of X . If no such elements exist, then $x \in (a, b)$, for some lowerbound a and upperbound b . Thus, in all three cases, there is a basis element containing x .

Now suppose $B_1, B_2 \in \mathcal{B}$ such that $x \in B_1 \cap B_2$. If B_1 and B_2 are both of type (1), then let $B_1 = (a, b)$, $B_2 = (c, d)$, then $B_1 \cap B_2$ is an open interval of type (1), now fix B_1 to be of type one. If B_2 is of type (2), then letting $B_2 = [a_0, c)$, then $x \in [a_0, d)$ for some $d \in X$. Likewise, if $B_2 = (c, b_0]$, is of type (3), we get a similar result. Moreover, the results are analogous if we fix B_2 and let B_1 range between intervals of the three types. Thus in all cases, there is a $B_3 \in \mathcal{B}$ such that $x \in B_3 \subseteq B_1 \cap B_2$. ■

Example 1.4. (1) The standard topology on \mathbb{R} is the order topology on \mathbb{R} induced by the usual order relation. We have that \mathbb{R} under this topology has no intervals of type (2), nor (3), so all bases elements in the standard topology are open intervals in \mathbb{R} .

- (2) Consider the dictionary order on $\mathbb{R} \times \mathbb{R}$. Since $\mathbb{R} \times \mathbb{R}$ has no intervals of type (2), nor (3), the bases of $\mathbb{R} \times \mathbb{R}$ under the dictionary order are the open intervals of the form $(a \times b, c \times d)$ Where $a \leq c$, and $b < d$.




Figure 1.3: The order topology on $\mathbb{R} \times \mathbb{R}$.

- (3) The positive integers \mathbb{Z}^+ with the least element 1 form an ordered set under the usual order. Taking $n > 1$, we see the bases of \mathbb{Z}^+ under the order topology are of the form $(n-1, n+1) = \{n\}$ and $[1, n) = \{1, \dots, n-1\}$. Thus the order topology on \mathbb{Z}^+ is the discrete topology.
- (4) The set $X = \{1, 2\} \times \mathbb{Z}^+$ over the dictionary order is also an ordered set, with the least element 1×1 . Denote $1 \times n$ as a_n and $2 \times n$ as b_n . Then X consist of the elements $a_1, a_2, \dots, b_1, b_2, \dots$.

Now take $\{b_1\}$, then any open set containing b_1 must have a basis about b_1 , and also contains points a_i with $i \in \mathbb{Z}^+$; thus the order topology on X is not the discrete topology.

Definition. Let X be an ordered set, and let $a \in X$. There are two subsets in X , $(a, \infty) = \{x \in X : x > a\}$ and $(-\infty, a) = \{x \in X : x < a\}$ called **open rays** of X . There are also two sets $[a, \infty) = \{x \in X : x \geq a\}$ and $(-\infty, a] = \{x \in X : x \leq a\}$ called **closed rays** of X .

Theorem 1.3.2. *Let X be an ordered set. Then the collection of all open rays in X form a subbasis for the order topology on X .*

Proof. Let \mathcal{S} be the collection of all open rays of X , let $a < b$ and $(a, \infty), (-\infty, b) \in \mathcal{S}$, then $(a, b) = (a, \infty) \cap (-\infty, b)$. Now take:

$$S = \bigcup_{a, b \in X} (a, b)$$

then $S \subseteq X$, likewise, since S runs through all intersections of open rays of X , it contains all open intervals in X , hence $X \subseteq S$, and so $X = S$ as required. ■

1.4 The Product Topology.

Definition. Let X and Y be topological spaces. We define the **product topology** on $X \times Y$ to be the topology having as basis the collection

$$\mathcal{B} = \{U \times V \subseteq X \times Y : U \text{ is open in } X \text{ and } V \text{ is open in } Y\}$$

Theorem 1.4.1. *The collection $\mathcal{B} = \{U \times V \subseteq X \times Y : U \text{ is open in } X \text{ and } V \text{ is open in } Y\}$ forms a basis for the product topology on $X \times Y$.*

Proof. Clearly, we have that $X \times Y$ is a basis element of \mathcal{B} . Now take $U_1 \times V_1$ and $U_2 \times V_2$ in \mathcal{B} . Since $U_1 \times V_1 \cap U_2 \times V_2 = U_1 \cap U_2 \times V_1 \cap V_2$, since $U_1 \cap U_2$ and $V_1 \cap V_2$ are open in X and Y respectively, then we have that $U_1 \times V_1 \cap U_2 \times V_2$ is a basis element as well. ■


Theorem 1.4.2. *If \mathcal{B} is the basis for a topology on X , and \mathcal{C} is the basis for a topology on Y , then the collection:*

$$\mathcal{D} = \{B \times C : B \in \mathcal{B} \text{ and } C \in \mathcal{C}\}$$

Is a basis for the topology on $X \times Y$.

Proof. By lemma 1.2.3, let W be an open set of $X \times Y$, and let $x \times y \in W$. Then there is a basis $U \times V$ such that $x \times y \in U \times V \subseteq W$. Since \mathcal{B} and \mathcal{C} are bases of X and Y respectively, choosing $B \in \mathcal{B}$ and $C \in \mathcal{C}$, we have that $x \in B \subseteq U$, and $y \in C \subseteq V$, thus $x \times y \in B \times C \subseteq U \times V \subseteq W$. Therefore, \mathcal{D} is the basis for a topology on $X \times Y$. ■

Example 1.5. The product of the standard topology on \mathbb{R} with itself is called the **standard topology on $\mathbb{R} \times \mathbb{R}$** , and has as basis the collection of all products of open sets in \mathbb{R} . By theorem 1.4.2, if we take the collection of all open intervals $(a, b) \times (c, d)$ in $\mathbb{R} \times \mathbb{R}$, we form a basis. Constructing this basis geometrically gives the interior of a rectangle, whose boundaries are the intervals (a, b) and (c, d) .


Figure 1.4: A basis element for $\mathbb{R} \times \mathbb{R}$

Definition. Let $\pi_1 : X \times Y \rightarrow X$ be defined such that $\pi_1(x, y) = x$, and define $\pi_2 : X \times Y \rightarrow Y$ such that $\pi_2(x, y) = y$. We call π_1 and π_2 **projections** of $X \times Y$ onto its first and second **factors**; that is onto X and Y , respectively.

Clearly, π_1 and π_2 are both onto. Now let U be open in X , then $\pi_1^{-1}(U) = U \times Y$ is open in $X \times Y$; similarly, $\pi_2^{-1}(V) = X \times V$ is also open in $X \times Y$, for V open in Y .

Theorem 1.4.3. *The collection $\mathcal{S} = \{\pi_1^{-1}(U) : U \text{ is open in } X\} \cup \{\pi_2^{-1}(V) : V \text{ is open in } Y\}$ is a subbasis for the product topology on X .*

Proof. Let \mathcal{T} be the product topology on $X \times Y$, and let \mathcal{T}' be the topology generated by \mathcal{S} . Since every element of \mathcal{S} is open in \mathcal{T} , $\mathcal{T} \subseteq \mathcal{T}'$. Conversely, consider the basis element $U \times V$ of \mathcal{T} , then $\pi_1^{-1}(U) \cap \pi_2^{-1}(V) = U \times Y \cap X \times V = U \times V$, thus $\mathcal{T} \subseteq \mathcal{T}'$. Therefore, \mathcal{S} is a subbasis for the product topology. ■

Figure 1.5: The inverse images, $\pi_1^{-1}(U)$ and $\pi_2^{-1}(V)$, of the projections π_1 and π_2 onto the $X \times Y$ plane.

1.5 The Subspace Topology.

Theorem 1.5.1. *Let X be a topological space with topology \mathcal{T} , and let $Y \subseteq X$. Then the collection:*

$$\mathcal{T}_Y = \{Y \cap U : U \in \mathcal{T}\}$$

forms a topology on Y .

Proof. Clearly, $Y \cap \emptyset = \emptyset \in \mathcal{T}_Y$ and $Y \cap X = Y \in \mathcal{T}_Y$. Now consider the collection $\{U_\alpha\}$. Then $\bigcup (Y \cap U_\alpha) = Y \cap \bigcup U_\alpha$, similarly, for $\{U_i\}_{i=1}^n$, $\bigcap (Y \cap U_i) = Y \cap \bigcap U_i$, hence \mathcal{T} is a topology on Y . ■

Definition. Let X be a topological space, and let $Y \subseteq X$. We call the \mathcal{T} defined in theorem 1.5.1 the **subspace topology** on Y . We say that $U \subseteq Y$ is **open in Y** if $U \in \mathcal{T}_Y$.

Lemma 1.5.2. *Let \mathcal{B} be the basis for a topology on X . Then the collection $\mathcal{B}_Y = \{B \cap Y : B \in \mathcal{B}\}$, where $Y \subseteq X$, is a basis for the subspace topology on Y .*

Proof. Let U be open in X , and let $y \in Y \cap U$, and choose $B \in \mathcal{B}$ such that $y \in B \subseteq U$, then $y \in B \cap Y \subseteq U \cap Y$, then by lemma 1.2.2, \mathcal{B}_Y is the basis for the subspace topology on Y . ■

Lemma 1.5.3. *Let Y be a subspace of X , If $U \subseteq Y$ is open in Y and Y is open in X , then U is open in X .*

Proof. Let $U \in \mathcal{T}_Y$, then for some $V \subseteq X$, $U = Y \cap V$. Now since Y is open in X , and so is V , then it follows that U is also open in X . ■

Remark. What this lemma says is that given a topological space X , and a subspace Y of X , then the subspace topology of Y is coarser than the topology on X , i.e. $\mathcal{T}_Y \subseteq \mathcal{T}$.

Theorem 1.5.4. *If A is a subspace of X , and B is a subspace of Y , then the product topology on $A \times B$ is the topology that $A \times B$ inherits as a subspace of $X \times Y$.*

Proof. We have that $U \times V$ is the basis element for $X \times Y$, with U open in X , and V open in Y . Thus $(U \times V) \cap (A \times B) = (U \cap A) \times (V \cap B)$ is a basis element for the subspace topology on $X \times Y$. Since $U \cap A$ and $V \cap B$ are open in the subspace topologies of A and B respectively, then $(U \cap A) \times (V \cap B)$ is a basis for the product topology on $A \times B$. ■

Example 1.6. (1) Consider $[0, 1] \subseteq \mathbb{R}$. In the subspace topology of $[0, 1]$, we have as basis elements of the form $(a, b) \cap [0, 1]$, with $(a, b) \subseteq \mathbb{R}$. If we have that $(a, b) \subseteq [0, 1]$, then $(a, b) \cap [0, 1] = (a, b)$. On the other hand, if $a \in [0, 1]$ or $b \in [0, 1]$, then we get $(a, b) \cap [0, 1] = (a, 1]$ or $(a, b) \cap [0, 1] = [0, b]$, lastly if neither a nor b are in $[0, 1]$, then we have $(a, b) \cap [0, 1] = [0, 1]$ only if $[0, 1] \subseteq (a, b)$, and $(a, b) \cap [0, 1] = \emptyset$ otherwise.

The only one of these sets open in \mathbb{R} under the standard topology is $(0, 1)$.

(2) For $[0, 1] \cup \{2\} \subseteq \mathbb{R}$, the singleton $\{2\}$ is open in the subspace topology on $[0, 1] \cup \{2\}$; for observe, that $(\frac{3}{5}, \frac{5}{2}) \cap ([0, 1] \cup \{2\}) = \{2\}$, however, in the order topology, on that same set, $\{2\}$ is not open. Any basis element on $[0, 1] \cup \{2\}$ containing 2 is of the form $(a, 2]$, where $a \in [0, 1] \cup \{2\}$.

- (3) The dictionary order on $[0, 1] \times [0, 1]$ is a restriction of the dictionary order on $\mathbb{R} \times \mathbb{R}$. Now the set $\{\frac{1}{2}\} \times (\frac{1}{2}, 1]$ is open in the subspace topology on $[0, 1] \times [0, 1]$, but it is not open in the dictionary order on the same set.

Definition. We call the set $[0, 1] \times [0, 1]$ on the dictionary order the **ordered square**, and we denote it by I_0^2 .

Definition. Let X be an ordered set. We say that a nonempty subset $Y \subset X$ is **convex** in X if for each pair of points $a, b \in Y$, with $a < b$, then the open interval $(a, b) \subseteq X$ is also contained in Y .




Figure 1.6: A convex set, and a nonconvex set.

Example 1.7. Let X be any ordered set. Then by definition, all open intervals and rays in X are convex in X .

Theorem 1.5.5. Let X be an ordered set on the order topology, and let $Y \subseteq X$ be convex in X . Then the order topology on Y is the same as the subspace topology on Y .

Proof. Consider $(a, \infty) \subseteq X$. If $a \in Y$, then $(a, \infty) \cap Y = \{x \in Y : x > a\}$, which is by definition an open ray on Y . Now if $a \notin Y$, then a is either a lowerbound, or an upperbound. Then $(a, \infty) \cap Y = \emptyset$ and $(-\infty, a) \cap Y = Y$ if a is an upperbound, similarly, if a is a lowerbound we get $(a, \infty) \cap Y = Y$ and $(-\infty, a) \cap Y = \emptyset$.

Since $(a, \infty) \cap Y$ and $(-\infty, a) \cap Y$ form a subbasis on the subspace topology on Y , and since they are also open in the order topology, then the order topology contains the subspace topology.

Now if (a, ∞) is an open ray in Y , then $(a, \infty) = (b, \infty) \cap Y$, with (b, ∞) some open ray in X , hence (a, ∞) is open in the subspace topology of Y , and since it also forms the subbasis for the order topology, we have that the order topology is contained within the subspace topology. Thus both topologies are equal. ■




Figure 1.7: An illustration of theorem 1.5.5.

1.6 Closed Sets and Limit Points.

Definition. A subset A of a topological space X is said to be **closed** if $X \setminus A$ is open.

Example 1.8. (1) Consider $[a, b] \subseteq \mathbb{R}$, we have that $\mathbb{R} \setminus [a, b] = (-\infty, a) \cup (b, \infty)$ which is open in \mathbb{R} . So $[a, b]$ is closed.

- (2) In $\mathbb{R} \times \mathbb{R}$, the set $A = \{x \times y : x, y \geq 0\}$ (i.e the first quadrant of the plane) is closed, for $\mathbb{R} \times \mathbb{R} \setminus A = ((-\infty, 0) \times \mathbb{R}) \cup (\mathbb{R} \times (-\infty, 0))$, which is open in $\mathbb{R} \times \mathbb{R}$.
- (3) Consider the finite complement topology \mathcal{T}_f on a set X . We have that $X \setminus X = \emptyset \in \mathcal{T}_f$, so X is closed, similarly, \emptyset is also closed. Likewise, if $A \subseteq X$ is a finite set, then $X \setminus A$ is also finite, and hence A is also closed. Thus, we have that all the closed sets of \mathcal{T}_f are those finite subsets of X . As a consequence, this example also illustrates that sets can be both closed and open.
- (4) In the discrete topology 2^X , every open set is closed. This is another example where open sets are also closed sets.
- (4) Consider $[0, 1] \cup (2, 3)$ in the subspace topology on \mathbb{R} . We have that $[0, 1]$ is open in the subspace topology on \mathbb{R} ; $[0, 1] = [0, 1] \cup (2, 3) \cap (-\frac{2}{3}, \frac{3}{2})$, similarly, $(2, 3)$ is also open. Now taking $[0, 1] \cup (2, 3) \setminus (2, 3) = [0, 1]$, which is open, so $[0, 1]$ is closed in the subspace topology on \mathbb{R} , by the same reasoning, so is $(2, 3)$.

Theorem 1.6.1. Let X be a topological space. Then:

- (1) \emptyset and X are closed.
- (2) Arbitrary intersections of closed sets are closed.
- (3) Finite unions of closed sets are closed.

Proof. We have that $X \setminus \emptyset = X$ and $X \setminus X = \emptyset$, both of which are open in X , so they are also closed in X . Now let $\{U_\alpha\}$ be a collection of closed sets of X . We have that:

$$X \setminus \bigcap_{\alpha} U_{\alpha} = \bigcup_{\alpha} X \setminus U_{\alpha}.$$

Similarly, for $\{U_i\}_{i=1}^n$, we have

$$X \setminus \bigcup_{i=1}^n U_i = \bigcap_{i=1}^n X \setminus U_i.$$

Both of which are open in X . This completes the proof. ■

Definition. If Y is a subspace of X , we say that A is **closed in Y** if $A \subseteq Y$ and A is closed in the subspace topology on Y .

Theorem 1.6.2. *Let Y be a subspace of X . Then A is closed in Y if and only if A equals the intersection of a closed set of X with Y .*

Proof. Suppose that A is closed in Y , then $Y \setminus A$ is open in Y , hence we have that $Y \setminus A = U \cap Y$ for some open set U of X . Now $X \setminus U$ is closed in X , and with $A \subseteq Y$, we have that $A = Y \cap (X \setminus U)$.

Conversely, suppose that $A = C \cap Y$, with C closed in X . Then $X \setminus C$ is open in X , hence $(X \setminus C) \cap Y$ is open in Y , now since $(X \setminus C) \cap Y = Y \setminus A$, which is open, we have that A is closed in Y . ■

Theorem 1.6.3. *Let Y be a subspace of X . If A is closed in Y , and Y is closed in X , then A is closed in X ; that is, closure is transitive.*

Proof. By theorem 1.6.2, if A is closed in Y , then $A = C \cap Y$ with C closed in X , now since Y is closed in X , then $Y = D \cap X$ with D closed in X . Thus $A = (C \cap D) \cap X$, therefore, A is closed in X . ■

We now go over the concepts of the closure, and the interior of a set.

Definition. Let $A \subseteq X$, with X a topological space. The **interior** of A is defined to be the union of all open sets in A . The **closure** of A is defined to be the intersection of all closed sets containing A . We denote the interior and the closure of A as $\text{Int } A$ and $\text{cl } A$ respectively

We have by the very definitions that $\text{Int } A \subseteq A \subseteq \text{cl } A$

Lemma 1.6.4. *$\text{Int } A = A$ only when A is open, and $\text{cl } A = A$ only when A is closed.*

Proof. Now, if A is open, then it is in the union of all open sets of A , hence $A \subseteq \text{Int } A$, likewise, if A is closed, then since $\text{cl } A$ is the intersection of all closed sets containing A , we get $\text{cl } A \subseteq A$. ■

Corollary. *A is closed and open if and only if $\text{Int } A = \text{cl } A$.*

Theorem 1.6.5. *Let Y be a subspace of X , and let $A \subseteq Y$. Then $\text{cl } A \cap Y$ is the closure of A in Y .*

Proof. Let $\text{cl } A$ be the closure of A in X . Since $\text{cl } A$ is closed in X , by theorem 1.6.2, $\text{cl } A \cap Y$ is closed in Y , now we have that $A \subseteq \text{cl } A \cap Y$, and since $\text{cl } A = \bigcap U$, then $\text{cl } A \subseteq \text{cl } A \cap Y$.

Conversely, suppose that $\text{cl } A$ is closed in Y , again by theorem 1.6.2, we have that $\text{cl } A = C \cap Y$, where C is closed in X , since $A \subseteq \text{cl } A$, then $A \subseteq C$, and since C is closed, then $\text{cl } A \subseteq C$, thus $\text{cl } A \cap Y \subseteq \text{cl } A$. ■

Definition. Let X be a topological space, and let $x \in X$. We call an open set U of X a **neighborhood** of x if $x \in U$.

Theorem 1.6.6. *If $A \subseteq X$, with X a topological space, then $\text{cl } A$ is a neighborhood of $x \in X$ if and only if for every neighborhood U of x , $A \cap U \neq \emptyset$.*

Proof. We prove the contrapositive. If $x \notin \text{cl } A$, then $U = X \setminus \text{cl } A$ is an open set containing x , disjoint from A . Conversely, suppose there is a neighborhood U of x , with U disjoint from A , then $X \setminus U$ is closed, and therefore contains the closure of A , thus $x \notin \text{cl } A$. ■

Corollary. *$\text{cl } A$ is a neighborhood of x if and only if for every basis element B of X , containing x , intersects A .*

Proof. This is a direct application of theorem 1.6.6, since basis elements are open sets. ■

Example 1.9. (1) We have the closure of $(0, 1]$ in \mathbb{R} is the closed interval $[0, 1]$, since every neighborhood of 0 intersects $(0, 1]$. Now every point outside of $[0, 1]$ has a neighborhood disjoint from $[0, 1]$ (take the neighborhood $(2, 3)$ of 2).

$$(2) \text{cl } \frac{1}{\mathbb{Z}^+} = \{0\} \cup \frac{1}{\mathbb{Z}^+} \text{ and } \text{cl } \{0\} \cup (1, 2) = \{0\} \cup [1, 2].$$

$$(3) \text{cl } \mathbb{Q} = \mathbb{R}, \text{cl } \mathbb{Z}^+ = \mathbb{Z}^+, \text{cl } \mathbb{R}^+ = \mathbb{R}^+ \cup \{0\}. \text{ This first follows from the density of } \mathbb{Q} \text{ in } \mathbb{R}. \text{ Every neighborhood } n \in \mathbb{Z}^+ \text{ intersects } \mathbb{Z}^+, \text{ so } \text{cl } \mathbb{Z}^+ \subseteq \mathbb{Z}^+, \text{ and we have that the neighborhood } (0, 1) \text{ of } 0 \text{ intersects } \mathbb{R}^+, \text{ so } \text{cl } \mathbb{R}^+ \subseteq \mathbb{R}^+ \cup \{0\}.$$

Definition. If $A \subseteq X$, with X a topological space, and if $x \in X$, we say that x is a **limit point** of A if every neighborhood of x intersects A at some distinct point. That is: $x \in \text{cl } X \setminus \{x\}$.

Example 1.10. (1) Consider $(0, 1]$, we have that $0 \in [0, 1] = \text{cl } (0, 1] = \text{cl } [0, 1] \setminus \{0\}$, so 0 is a limit point of $(0, 1]$, the same can be said for any $x \in (0, 1]$.

- (2) For $\frac{1}{\mathbb{Z}^+}$, 0 is once again a limit point. Let $x \in \mathbb{R}$ be nonzero, and let $[x, b)$ be the neighborhood of x in the lower limit topology. Then $[x, b) \cap \frac{1}{\mathbb{Z}^+} = \emptyset$ or $\{x\}$, hence, 0 is the only limit point of $\frac{1}{\mathbb{Z}^+}$.
- (3) $\text{cl } \{0\} \cup (1, 2) = \{0\} \cup [1, 2]$ has all of its limit points in $[1, 2]$. Likewise, every point in \mathbb{R} is a limit point of \mathbb{Q} . \mathbb{Z}^+ has no limit points in \mathbb{R} , and the limit points of \mathbb{R}^+ are all the points of $\text{cl } \mathbb{R}^+$.

Theorem 1.6.7. *Let $A \subseteq X$, X a topological space, and let A' be the set of all limit points in A . Then $\text{cl } A = A \cup A'$.*

Proof. Let $x \in A'$, then every neighborhood of x intersects A at some distinct point x' , by definition, so by theorem 1.6.6, $x \in \text{cl } A$, hence $A' \subseteq \text{cl } A$, so $A \cup A' \subseteq \text{cl } A$. Now, let $x \in \text{cl } A$. If $x \in A$, we are done. Otherwise, since every neighborhood of x intersects A , we have that they intersect at distinct points, thus $x \in A'$, therefore $\text{cl } A \subseteq A \cup A'$. ■

Corollary. $A \subseteq X$ is closed if and only if $A' \subseteq A$.

Proof. If A is closed, then $\text{cl } A = A = A \cup A'$, thus $A' \subseteq A$. The converse is obvious. ■

Definition. Let X be a topological space. A sequence of points of X $\{x_n\}$ is said to **converge** to a point $x \in X$ if for every neighborhood U of x , there is an $N \in \mathbb{Z}^+$ such that $x_n \in U$ for all $n \geq N$.

Example 1.11. Consider the following topological space on $\{a, b, c\}$ in figure ??, and define




Figure 1.8: A topology on $\{a, b, c\}$, is not a Hausdorff space.

the sequence $\{x_n\}$ by $x_n = b$ for all $n \in \mathbb{Z}^+$. The neighborhoods of a , b , and c are $U_a = \{a, b\}$, $U_b = \{b\}$, and $U_c = \{b, c\}$. Now let $N > 0$, then we see that for all $n \geq N$, that $b \in U_a$, $b \in U_b$, and $b \in U_c$ thus b converges to a and to c , and itself,

Definition. A topological space X is called a **Hausdorff space** if for each pair of distinct points x_1 , and x_2 , there are neighborhoods U_1 and U_2 of x_1 and x_2 respectively such that U_1 and U_2 are disjoint.

Example 1.12. The topology of the previous example in figure 1.8 is not a Hausdorff space.

Theorem 1.6.8. Every finite point set in a Hausdorff space is closed.

Proof. Let X be a Hausdorff space, and let $x_0 \in X$. We have that $\text{cl } \{x_0\} = \bigcap_{\{x_0\} \in U} U$. Now let $x \neq x_0 \in X$. Since $x \notin \{x_0\}$, and X is Hausdorff, the intersections of the neighborhoods of x and x_0 is empty, thus $x \notin \text{cl } \{x_0\}$, therefore $\text{cl } \{x_0\} = \{x_0\}$.

We can then extend this proof to finite point sets of size n by induction. ■

Now the condition that finite point sets be closed need not depend on whether or not X is a Hausdorff space. In fact, we can assume the following for some topological spaces.

Axiom 1.6.1 (The T_1 Axiom). In any topological space, every finite point set of X is closed.

Theorem 1.6.9. Let X be a topological space satisfying the T_1 axiom, and let $A \subseteq X$. Then a point x is a limit point of A if and only if every neighborhood of x contains infinitely many points of A .

Proof. Let U_x be a neighborhood of x . If U_x intersects A at infinitely many points of A , then it intersects A at a point distinct from x , thus x is a limit point of A .

Conversely suppose that x is a limit point of A , and let $U_x \cap A$ be finite, then $U_x \cap (A \setminus \{x\})$ is also finite. Now let $U_x \cap (A \setminus \{x\}) = \{x_1, \dots, x_m\}$. By the T_1 axiom, $\{x_1, \dots, x_m\}$ is closed, so $X \setminus \{x_1, \dots, x_m\}$ is open, thus $U_x \cap (X \setminus \{x_1, \dots, x_m\})$ is a neighborhood of x that does not intersect $A \setminus \{x\}$, which contradicts that x is a limit point. ■

Theorem 1.6.10. *If X is a Hausdorff space, then a sequence of points of X converges to at most one point in X .*

Proof. Let $\{x_n\}$ be a sequence of points converging to x , and let $y \neq x$ and let U_x and U_y be neighborhoods of x and y respectively. Then $U_x \cap U_y = \emptyset$. Now since $\{x_n\}$ converges to x , we have that for $N > 0$, $x_n \in U_x$ whenever $n \geq N$. Then $x_n \notin U_y$, and so $\{x_n\}$ cannot converge to y . ■

Definition. Let $\{x_n\}$ be a sequence in a Hausdorff space X . If $\{x_n\}$ converges to a point $x \in X$, we call x the **limit** of $\{x_n\}$ as n **approaches** ∞ and we write $\lim_{n \rightarrow \infty} x_n = x$ or $\{x_n\} \rightarrow x$ as $n \rightarrow \infty$.

Theorem 1.6.11. *The following are true:*

- (1) *Every simply ordered set under the order topology is Hausdorff.*
- (2) *The product of two Hausdorff spaces is Hausdorff.*
- (3) *The subspace of a Hausdorff space is Hausdorff.*

Proof. (1) Let X be an ordered set under the order topology. Take $x, y \in X$ distinct, and suppose without loss of generality that $x < y$. Then consider the neighborhoods $(-\infty, x]$ and $[y, \infty)$ of x and y respectively. Then $(-\infty, x] \cap [y, \infty) = \emptyset$.

- (2) Let X and Y be Hausdorff, and consider $X \times Y$ in the product topology. Let $x_1 \times y_1$ and $x_2 \times y_2$ be distinct points, and let $U_{x_1}, U_{x_2}, V_{y_1}$ and V_{y_2} be basis elements of x_1, x_2, y_1 , and y_2 respectively. Then they are neighborhoods of those elements respectively.

Now we have that $U_{x_1} \times V_{y_1}$ and $U_{x_2} \times V_{y_2}$ are basis elements of $x_1 \times y_1$ and $x_2 \times y_2$, respectively, and hence neighborhoods of those elements respectively. Then we have $(U_{x_1} \times V_{y_1}) \cap (U_{x_2} \times V_{y_2}) = (U_{x_1} \cap U_{x_2}) \times (V_{y_1} \cap V_{y_2}) = \emptyset \times \emptyset = \emptyset$.

- (3) Let X be Hausdorff, and let Y be a subspace of X . Let x_1 and x_2 be distinct points, and let U_{x_1} and U_{x_2} be their neighborhoods. Since Y is open in X , then so are $Y \cap U_{x_1}$ and $Y \cap U_{x_2}$, so they are also neighborhoods of x_1 and x_2 respectively. Then $(Y \cap U_{x_1}) \cap (Y \cap U_{x_2}) = Y \cap (U_{x_1} \cap U_{x_2}) = \emptyset$. ■

1.7 Continuous Functions.

Definition. Let X and Y be topological spaces. We say that a mapping $f : X \rightarrow Y$ is **continuous** if for each open set V in Y , $f^{-1}(V)$ is open in X .

Now if $f : X \rightarrow Y$ is continuous, then for every open set V of Y , $f^{-1}(V)$ is open in X . Now suppose that \mathcal{B} is a basis of Y , then $V = B_\alpha$, hence $f^{-1}(B_\alpha)$ is open in X .

Similarly, if \mathcal{S} is a subbasis of Y , then for any basis element B of Y , $B = \bigcap_{i=1}^n S_i$, which then implies that $f^{-1}(B) = \bigcap_{i=1}^n f^{-1}(S_i)$, thus $f^{-1}(S_i)$ is also open in X for $1 \leq i \leq n$.

Example 1.13. (1) Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a continuous realvalued function. Then for each open interval $I \subseteq \mathbb{R}$, $f^{-1}(I)$ is an open interval in \mathbb{R} , so take $x_0 \in \mathbb{R}$ and $\varepsilon > 0$, and let $I = (f(x_0) - \varepsilon, f(x_0) + \varepsilon)$, then since $x_0 \in f^{-1}(I)$, there is a basis $(a, b) \subseteq f^{-1}(I)$ about x_0 . Then take $\delta = \min\{x_0 - a, x_0 - b\}$, then $x \in (a, b)$ whenever $0 < |x - x_0| < \delta$, and we get that $f(x) \in I$, that is, $|f(x) - f(x_0)| < \varepsilon$. This is the definition of continuity defined in the real analysis. We can prove that the converse holds also.

If $f : \mathbb{R} \rightarrow \mathbb{R}$ is continuous at a point x_0 , then for every $\varepsilon > 0$, there is a $\delta > 0$ such that $|f(x) - f(x_0)| < \varepsilon$ whenever $0 < |x - x_0| < \delta$. Then we notice that x and x_0 are distinct, furthermore, $x_0 - \delta < x < x_0 + \delta$, hence $x \in (x_0 - \delta, x_0 + \delta)$ implies that $f(x) \in (f(x_0) - \varepsilon, f(x_0) + \varepsilon)$. Letting $V_\delta(x_0) = (x_0 - \delta, x_0 + \delta)$ and $V_\varepsilon(f(x_0)) = (f(x_0) - \varepsilon, f(x_0) + \varepsilon)$, we have that whenever $x \in V_\delta(x_0)$, then $f(x) \in V_\varepsilon(f(x_0)) \subseteq f^{-1}(V_\delta(x_0))$. And so the topological definition of continuity is equivalent to the real analytic definition of continuity.

(2) Let $f : \mathbb{R} \rightarrow \mathbb{R}_l$ be defined such that $f(x) = x$ for all $x \in \mathbb{R}$. Take $[a, b) \subseteq \mathbb{R}_l$, we have that $f^{-1}([a, b)) = [a, b)$, which is not open in \mathbb{R} (under the standard topology), hence f is not continuous. However, the map $g : \mathbb{R}_l \rightarrow \mathbb{R}$ defined the same way is continuous since $g^{-1}([a, b))$ is open in \mathbb{R}_l .

Theorem 1.7.1. Let X and Y be topological spaces, and let $f : X \rightarrow Y$ be a mapping of X into Y . Then the following are equivalent:

- (1) f is continuous.
- (2) For every $A \subseteq X$, $f(\text{cl } A) \subseteq \text{cl } f(A)$.
- (3) For every closed set $B \subseteq Y$, $f^{-1}(B)$ is closed in X .
- (4) For each $x \in X$ and each neighborhood V of $f(x)$, there is a neighborhood U of x such that $f(U) \subseteq V$.

Proof. Let f be continuous and let $A \subseteq X$. Consider the neighborhood V of $f(x)$, then $f^{-1}(V)$ is open in X , and intersects A at a point y . Then $V \cap f(A) = f(y)$, thus $f(x) \in \text{cl } f(A)$.

Now let B be closed in Y , and let $A = f^{-1}(B)$. Then we have that $f(A) = f(f^{-1}(B)) \subseteq B$, thus $x \in \text{cl } A$.

Now let V be open in Y , so that $B = Y \setminus V$ is closed in Y , and $f^{-1}(B) = f^{-1}(Y) \setminus f^{-1}(V) = X \setminus f^{-1}(V)$ which is closed in X , hence $f^{-1}(V)$ is open in X .

Now let $x \in X$, and let V be a neighborhood of $f(x)$. Then $U = f^{-1}(V)$ is a neighborhood of x for which $f(U) \subseteq V$. Finally let V be open in Y , and let $x \in f^{-1}(V)$, then $f(x) \in V$, so there is a neighborhood U_x of x for which $f(U_x) \subseteq V$, then $U_x \subseteq f^{-1}(V)$, then $f^{-1}(V)$ is a union of open sets, and hence open in X . ■

Definition. Let X and Y be topological spaces, and $f : X \rightarrow Y$ be a 1 – 1 mapping of X onto Y . We call f a **homeomorphism** if both f and f^{-1} are continuous.

Lemma 1.7.2. *Let X and Y be topological spaces and let $f : X \rightarrow Y$ be a homeomorphism. Then $f(U)$ is open if and only if U is open.*

Proof. We have that both $f : X \rightarrow Y$ and $f^{-1} : Y \rightarrow X$ are continuous 1 – 1 of X and Y onto each other (respectively). Now let U be open in X , then $U = f^{-1}(V)$, for some set V open in Y . Notice then, that $f(U) = f(f^{-1}(V)) = V$, thus $f(U)$ is open in Y . Conversely, let $V = f(U)$ be open in Y for some open set U in X , then $U = f^{-1}(V)$, so by definition of continuity, U is open in X . ■

Definition. Let X and Y be topological spaces and let $f : X \rightarrow Y$ be a continuous 1 – 1 mapping of X into Y , and consider $f(X)$ as a subspace of Y . We call $f : X \rightarrow f(X)$ a **topological imbedding** if f is a homeomorphism of X onto $f(X)$.

Example 1.14. (1) The map $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = 3x + 1$ is a homeomorphism whose inverse is $f^{-1}(y) = \frac{1}{3}(y - 1)$, both f and f^{-1} are continuous.

(2) The map $f : (-1, 1) \rightarrow \mathbb{R}$ defined by $f(x) = \frac{x^2}{1-x^2}$ has as its inverse the map $f : \mathbb{R} \rightarrow (-1, 1)$ defined by $f^{-1}(y) = \frac{2y}{1+\sqrt{1+4y^2}}$. Both f and f^{-1} are continuous, so f is a homeomorphism.

(3) The map $g : \mathbb{R}_+ \rightarrow \mathbb{R}$ defined by $g(x) = x$ is not a homeomorphism, despite being continuous, as $g^{-1}(1)$ is undefined.

(4) Let S^1 be the unit circle in \mathbb{R}^2 , which is a subspace of \mathbb{R}^2 , and define $f : [0, 1) \rightarrow S^1$ by $f(t) = (\cos(2t\pi), \sin(2t\pi))$. Clearly f is 1 – 1 onto S^1 , and continuous, however f^{-1} is not continuous as $f([0, \frac{1}{4}))$ is not open in S^1 as $f(0)$ is in no open set of \mathbb{R}^2 such that $U \cap S^1 = f([0, 1))$.

(5) Consider the mappings $g : [0, 1) \rightarrow \mathbb{R}^2$ by $g(t) = (\cos(2t\pi), \sin(2t\pi))$. Now g is 1 – 1 and continuous, and we have that $g([0, 1)) \subseteq S^1$, however since g is not a homeomorphism, g fails to be a topological embedding.

Theorem 1.7.3 (Constructions for continuous functions.). *Let X and Y be topological spaces, then:*

- (1) (Constant construction) *If $f : X \rightarrow Y$ maps $x \rightarrow y_0$ for all $x \in X$, then f is continuous.*
- (2) (Inclusion) *If $A \subseteq X$ is a subspace, then the inclusion mapping $\iota : A \rightarrow X$ is continuous.*

- (3) (Construction by composition) If $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ are continuous, then $f \circ g : X \rightarrow Z$ is also continuous.
- (4) (Domain restriction) If $f : X \rightarrow Y$ is continuous and $A \subseteq X$, then $f : A \rightarrow Y$ is continuous.
- (5) (Range restriction) if $f : X \rightarrow Y$, and $Z \subseteq Y$ such that $f(X) \subseteq Z$, then $f : X \rightarrow Z$ is continuous.
- (6) (Range expansion) If $f : X \rightarrow Y$ is continuous, and $Y \subseteq Z$ is a subspace of Z , then $f : X \rightarrow Z$ is continuous.
- (7) (Local Formulation) The map $f : X \rightarrow Y$ is continuous if X can be written as the union of open sets U_α such that $f : U_\alpha \rightarrow Y$ is continuous for all α .

Proof. (1) Let $f(x) = y_0$ for all $x \in X$, and let V be open in Y , then $f^{-1}(V) = X$ or \emptyset depending on if $y_0 \in V$ or not. In either case, $f^{-1}(V)$ is open.

(2) If U is open in X , then $f^{-1}(U) = U \cap A$ which is open in the subspace topology of X .

(3) If U is open in Z , $g^{-1}(U)$ is open in Y , hence $f^{-1}g^{-1}(U)$ is open in X .

(4) Notice that $f_A = \iota \circ f = f : A \rightarrow Y$ which is continuous by (2) and (3).

(5) Let $f : A \rightarrow Y$ be continuous and let $f(X) \subseteq Z \subseteq Y$. Let B be open in Z , so $B = Z \cap U$ for some U open in Y . Now by hypothesis, we have that $f^{-1}(U) \subseteq f^{-1}(B)$, hence $f^{-1}(B)$ is open in X , thus $f : X \rightarrow Z$ is continuous.

(6) Let f be as in (5), and let $Y \subseteq Z$ be a subspace of Z . Then the mapping $h : X \rightarrow Z$ defined by $h = \iota \circ f$ is continuous.

(7) Let $X = \bigcup U_\alpha$ where U_α is open in X , and $f : U_\alpha \rightarrow Y$ is continuous for all α . Let V be open in Y , then $f^{-1}(V) \cap U_\alpha = f_{U_\alpha}^{-1}(V)$, and since f is continuous on U_α , then $f^{-1}(V) = \bigcup f_{U_\alpha}^{-1}(V)$ is open in X . ■

Theorem 1.7.4 (The pasting lemma). Let $X = A \cup B$ with A and B closed in X , and let $f : A \rightarrow Y$ and $g : B \rightarrow Y$ be continuous. If $f(x) = g(x)$ for all $x \in A \cap B$, then we can construct a mapping $h : X \rightarrow Y$ defined by $h(x) = \begin{cases} f(x), & x \in A \\ g(x), & x \in B \end{cases}$. Then h is continuous.

Proof. Let C be closed in Y , then $h^{-1}(C) = f^{-1}(C) \cup g^{-1}(C)$. Since f and g are continuous, then $f^{-1}(C)$ and $g^{-1}(C)$ are closed in A and B , respectively. Thus $h^{-1}(C)$ is closed in X . ■

Example 1.15. Define $h : \mathbb{R} \rightarrow \mathbb{R}$ by $h(x) = \begin{cases} x, & x \leq 0 \\ \frac{x}{2}, & x \geq 0 \end{cases}$. We have that x and $\frac{x}{2}$ are continuous on their respective domains, intersecting at 0, i.e. $x : (-\infty, 0] \rightarrow \mathbb{R}$, $\frac{x}{2} : [0, \infty) \rightarrow \mathbb{R}$, and $\{0\} = (-\infty, 0] \cap [0, \infty)$. Thus h is continuous on \mathbb{R} .

However, $k, l : \mathbb{R} \rightarrow \mathbb{R}$ defined by $k(x) = \begin{cases} x - 2, & x \leq 0 \\ x + 2, & x \geq 0 \end{cases}$ and $l(x) = \begin{cases} x - 2, & x < 0 \\ x + 2, & x \geq 0 \end{cases}$ are not continuous. We have that their domains intersect at 0, but that $k(0) = \pm 2$, (so k isn't even a function). Likewise, $(-\infty, 0) \cap [0, \infty) = \emptyset$, which is open in \mathbb{R} see 1.9.




Figure 1.9: The mappings h , k , and l .

Theorem 1.7.5. Let $f : A \rightarrow X \times Y$ be defined by $f(a) = (f_1(a), f_2(a))$, where $f_1 : A \rightarrow X$ and $f_2 : A \rightarrow Y$. Then f is continuous if and only if f_1 and f_2 are continuous.

Proof. Let $\pi_1 : X \times Y \rightarrow X$ and $\pi_2 : X \times Y \rightarrow Y$ be projections onto X and Y respectively. Since $\pi_1^{-1}(U) = U \times Y$ and $\pi_2^{-1}(V) = X \times V$ are both open in $X \times Y$, π_1 and π_2 are continuous. Then notice that $f_1(a) = \pi_1 \circ f(a)$ and $f_2(a) = \pi_2 \circ f(a)$, both of which are continuous.

Now suppose that f_1 and f_2 are continuous. We have that $a \in f^{-1}(U \times V)$ if and only if $f_1(a) \in U$ and $f_2(a) \in V$, then $f_1^{-1}(U)$ and $f_2^{-1}(V)$ are open in A , hence so is $f^{-1}(U \times V) = f_1^{-1}(U) \cap f_2^{-1}(V)$. ■

Definition. We define the **parametrized curve** of the plane \mathbb{R}^2 to be the continuous function $f : [a, b] \rightarrow \mathbb{R}^2$ defined by $f(t) = (x(t), y(t))$. If f is in a vector field, then we define $f(t) = x(t)i + y(t)j$ where $i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $j = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$.

Example 1.16. The function $f(t) = ((\cos(t)), \sin(t))$ is a parametrization of the curve $x^2 + y^2 = 1$, i.e. the unit circle S^1 .

Bibliography

- [1] J. Munkres, *Topology*. New York, NY: Pearson, 2018.