

Commutative Algebra

Alec Zabel-Mena

June 21, 2023

Contents

| | | |
|----------|---|----------|
| 1 | Rings and Ideals | 5 |
| 1.1 | Definitions and Examples | 5 |
| 1.2 | Polynomial Rings | 7 |
| 1.3 | Ring Homomorphisms and Factor Rings | 8 |
| 1.4 | Properties of Ideals | 10 |
| 1.5 | Euclidean Domains | 13 |
| 1.6 | Principal Ideal Domains. | 16 |
| 1.7 | Unique Factorization Domains. | 18 |
| 1.8 | The Nilradical and Jacobson Radical | 20 |
| 1.9 | Operations on Ideals | 21 |
| 1.10 | Extensions and Contractions of Ideals | 25 |
| 1.11 | Multivariate Polynomial Rings | 27 |
| 1.12 | Noetherian Rings | 29 |

Chapter 1

Rings and Ideals

1.1 Definitions and Examples

Definition. A **commutative ring** A is a set together with two binary operations $+$: $(a, b) \rightarrow a + b$ and \cdot : $(a, b) \rightarrow ab$ called **addition** and **multiplication** such that:

- (1) A is an Abelian group over $+$, where we denote the identity element as 0 and the inverse of each $a \in A$ as $-a$.
- (2) For any $a, b \in A$, $ab \in A$ and $a(bc) = (ab)c$. That is, A is closed under multiplication, and multiplication is associative.
- (3) $a(b + c) = ab + ac$ and $(a + b)c = ac + bc$.
- (4) $ab = ba$ for all $a, b \in A$.

If there exists an element $1 \in A$ such that $a1 = 1a = a$, then we call A a ring with **identity**. If $1 = 0$, we call A the **zero ring** and write $A = 0$.

Definition. A commutative ring k with identity $1 \neq 0$ is called a **field** if for all $a \in k$, where $a \neq 0$, there exists a $b \in A$ such that $ab = 1$.

Lemma 1.1.1. *Let A be a commutative ring with identity. Then the following are true for all $a, b \in A$.*

- (1) $0a = a0 = 0$.
- (2) $(-a)b = a(-b) = -(ab)$.
- (3) $(-a)(-b) = ab$
- (4) $1 \neq 0$, then 1 is unique and $-a = (-1)a$.

Proof. (1) Notice $0a = (0 + 0)a = 0a + 0a$, so that $0a = 0$. Likewise, $a0 = 0$ by the same reasoning.

- (2) Notice that $b - b = 0$, so $a(b - b) = ab + a(-b) = 0$, so that $a(-b) = -(ab)$. The same argument with $(a - a)b$ gives $(-a)b = -(ab)$.

- (3) By the inverse laws of addition in A , we have $-(a(-b)) = -(-(ab))$, so that $(-a)(-b) = ab$.
- (4) Suppose A has identity $1 \neq 0$, and suppose there is an element $2 \in A$ for which $2a = a2 = a$ for all $a \in A$. Then we have that $1 \cdot 2 = 1$ and $1 \cdot 2 = 2$, making $1 = 2$; so 1 is unique. Now, we have that $a + (-a) = 0$, so that $1(a + (-a)) = 1a + 1(-a) = 1a + (-a) = 0$. So $(-a) = -(1a) = (-1)a$ by (2). ■

Definition. Let A be a ring. We call an element $a \in A$ a **zero divisor** if $a \neq 0$ and there exists an element $b \neq 0$ such that $ab = 0$. Similarly, we call $a \in A$ a **unit** if there is a $b \in A$ for which $ab = ba = 1$. We call an element a **nilpotent** if there exists some $n \in \mathbb{Z}^+$ for which $x^n = 0$.

Definition. Let A be a ring. We call the set of all units in A the **group of units** and denote it $\mathcal{U}(A)$, or A^* .

Lemma 1.1.2. *Let A be a commutative ring with identity $1 \neq 0$. Then the group of units $\mathcal{U}(A)$ forms an Abelian group under multiplication.*

Proof. Let $a, b \in A$ be units in A . Then there are $c, d \in A$ for which $ac = ca = 1$ and $bd = db = 1$. Consider then ab . Then $ab(dc) = a(bd)c = ac = 1$ and $(dc)ab = d(ca)b = db = 1$ so that ab is also a unit in A . Moreover $\mathcal{U}(A)$ inherits the associativity of \cdot and 1 serves as the identity element of A^* . Lastly, if $a \in A^*$ is a unit there is a $b \in A$ for which $ab = ba = 1$. This also makes b a unit in A , and the inverse of a . Now, since A is a commutative ring, the multiplication in $\mathcal{U}(A)$ is commutative, making $\mathcal{U}(A)$ Abelian. ■

Corollary. *a is a zero divisor if, and only if it is not a unit.*

Proof. Suppose that $a \neq 0$ is a zero divisor. Then there is a $b \in A$ such that $b \neq 0$ and $ab = 0$. Then for any $v \in A$, $v(ab) = (va)b = 0$ so that a cannot be a unit. On the other hand let a be a unit, and $ab = 0$ for some $b \neq 0$. Then there is a $v \in A$ for which $v(ab) = (va)b = 1b = b = 0$. Then $b = 0$ which is a contradiction. ■

Corollary. *If k is a field, then it has no zero divisors.*

Proof. Notice by definition of a field, every element is a unit, except for 0. ■

Definition. A commutative ring with identity $1 \neq 0$ is called an **integral domain** if it has no zero divisors.

Lemma 1.1.3. *Any finite integral domain is a field.*

Proof. Let A be a finite integral domain and consider the map on A , by $x \rightarrow ax$. By above, this map is 1-1, moreover since A is finite, it is also onto. So there is a $b \in A$ for which $ab = 1$, making a a unit. Since a is arbitrarily chosen, this makes A a field. ■

Corollary. *If k is a field it is a (not necessarily finite) integral domain.*

Definition. A **subring** of a ring A is a subgroup of A closed under multiplication.

1.2 Polynomail Rings

Theorem 1.2.1. *Let A be a commutative ring with identity, and define $A[x] = \{f(x) = a_0 + a_1x + \cdots + a_nx^n : a_0, \dots, a_n \in A\}$. Define the operations $+$ and \cdot on $A[x]$ for $f(x) = a_0 + a_1x + \cdots + a_nx^n$ and $g(x) = b_0 + b_1x + \cdots + b_nx^n$ by:*

$$f + g = (a_0 + b_0) + (a_1 + b_1)x + \cdots + (a_n + b_n)x^n$$

$$fg = c_0 + c_1x + \cdots + c_kx^k \text{ where } c_j = \sum_{i=0}^j a_ib_{j-i} \text{ and } k = n + m$$

Then $A[x]$ is a commutative ring with identity.

Definition. Let A be a commutative ring with identity. We call the ring $A[x]$ the **ring of polynomials** in x with **coefficients** in A whose elements of the form

$$f(x) = a_0 + a_1x + \cdots + a_nx^n$$

where $n \geq 0$ are called **polynomails**. If $a_n \neq 0$, then the **degree** of f is denoted $\deg f = n$, and f is called **monic** if $a_n = 1$. We call $+$ and \cdot the **addition** and **multiplication** of polynomials.

Example 1.1. (1) Take A any commutative ring with identity and form $A[x]$. One can verify that the polynomial $0(x) = 0 + 0x + \cdots + 0x^n + \cdots = 0$, in this case we call 0 the **zero polynomail**. Similarly, the additive inverse of $f(x) = a_0 + a_1x + \cdots + a_nx^n$ is the polynomial $-f(x) = -a_0 - a_1x - \cdots - a_nx^n$. Now, since $A[x]$ has identity, the **identity** polynomial is $1(x) = 1 + 0x + \cdots = 1$, that is, it is the identity in A . Lastly, we call a polynomial f with $\deg f = 0$ a **constant polynomail**. Notice that 0 and 1 are constant polynomials.

(2) $\mathbb{Z}[x]$, $\mathbb{Q}[x]$, $\mathbb{A}[x]$ and $\mathbb{C}[x]$ are the polynomial rings in x with coefficients in \mathbb{Z} , \mathbb{Q} , \mathbb{A} , and \mathbb{C} respectively.

(3) Notice that the rings $\mathbb{Z}[\omega]$ and $\mathbb{Z}[i]$ are polynomial rings in ω and i , respectively, with coefficients in \mathbb{Z} , and where $\omega = \sqrt{D}$ if $D \not\equiv 1 \pmod{4}$ or $\omega = \frac{1+\sqrt{D}}{2}$ otherwise, and $i^2 = -1$. Notice that the highest degree a polynomial in $\mathbb{Z}[i]$ can achieve is $\deg = 1$; however, one may be able to form polynomial rings in other variables with coefficients in $\mathbb{Z}[i]$, i.e. take $Z[x]$, where $Z = \mathbb{Z}[i]$.

(4) $\mathbb{Z}/_3\mathbb{Z}[x]$ is the polynomial ring with coefficients in $\mathbb{Z}/_3\mathbb{Z}$.

Theorem 1.2.2. *Let A be an integral domain, and let $p, q \neq 0$ be polynomials in $A[x]$. Then the following are true:*

(1) $\deg pq = \deg p + \deg q$.

(2) *The units of $A[x]$ are precisely the units of A*

(3) $A[x]$ is an integral domain.

Proof. Consider the leading terms $a_n x^n$ and $b_m x^m$ of p and q respectively. Then $a_n b_m x^{m+n}$ is the leading term of pq ; moreover we require $a_n b_m \neq 0$. Now, if $\deg pq < m + n$, then $ab = 0$, making a and b zero divisors of A ; impossible. Therefore $ab \neq 0$. It also follows that since no term of p is a zero divisor, then p cannot be a zero divisor of $A[x]$. Lastly, if $pq = 1$, then $\deg p + \deg q = 0$, so that pq is a constant polynomial. Noticing that constant polynomials are simply just elements of A , then p and q are units. ■

1.3 Ring Homomorphisms and Factor Rings

Definition. Let A and B be commutative rings with identity. We call a map $\phi : A \rightarrow B$ a **ring homomorphism** if

- (1) ϕ is a group homomorphism with respect to addition.
- (2) $\phi(ab) = \phi(a)\phi(b)$ for any $a, b \in A$.
- (3) $\phi(1_A) = 1_B$.

We denote the **kernel** of ϕ to be the kernel of ϕ as a group homomorphism. That is

$$\ker \phi = \{r \in A : \phi(r) = 0\}$$

Moreover, if ϕ is 1-1 and onto, we call ϕ an **isomorphism** and say that A and B are **isomorphic**, and write $A \simeq B$.

Lemma 1.3.1. *Let A and B be commutative rings with identity, and $\phi : A \rightarrow B$ a ring homomorphism. Then*

- (1) $\phi(A)$ is a subring of B .
- (2) $\ker \phi$ is a subring of A .

Proof. Let $s_1, s_2 \in \phi(A)$. Then $s_1 = \phi(r_1)$ and $s_2 = \phi(r_2)$ for some $r_1, r_2 \in A$. Then $s_1 s_2 = \phi(r_1)\phi(r_2) = \phi(r_1 r_2) \in \phi(B)$. Additionally, $s^{-1} = \phi^{-1}(r) = \phi(r^{-1})$ for some $s \in B$, $r \in A$. This is sufficient to make B a subring of B .

By similar reasoning, if $r_1, r_2 \in \ker \phi$, then $\phi(r_1)\phi(r_2) = \phi(r_1 r_2) = 0$ so that $r_1 r_2 \in \ker \phi$, and $\phi(r^{-1}) = \phi^{-1}(r) = 0$ so $\phi^{-1} \in \ker \phi$. ■

Corollary. *For any $r \in A$ and $a \in \ker \phi$, then $ar \in \ker \phi$ and $ra \in \ker \phi$.*

Proof. We have $\phi(ar) = \phi(a)\phi(r) = \phi(a)0 = 0$ so $ar \in \ker \phi$. The same happens for ra . ■

Definition. Let A be a commutative ring with identity. We call a subset \mathfrak{a} of A an **ideal** of A if it is a subgroup under $+$, and for any $r \in A$, and $a \in \mathfrak{a}$, $ra \in \mathfrak{a}$.

Theorem 1.3.2. Let A be a commutative ring with identity, and $I\mathfrak{a}$ an ideal in A . Let A/\mathfrak{a} be the set of all $a + \mathfrak{a}$ with $a \in A$. Define operations $+$ and \cdot by

$$\begin{aligned}(a + \mathfrak{a}) + (b + \mathfrak{a}) &= (a + b) + \mathfrak{a} \\ (a + \mathfrak{a})(b + \mathfrak{a}) &= ab + \mathfrak{a}\end{aligned}$$

Then A/\mathfrak{a} forms a commutative ring with identity under $+$ and \cdot .

Proof. Notice that $(a + \mathfrak{a}) + (b + \mathfrak{a}) = (a + b) + (\mathfrak{a} + \mathfrak{a}) = (a + b) + 2\mathfrak{a} = (a + b) + \mathfrak{a}$. Moreover, A/\mathfrak{a} inherits associativity in $+$ from addition in A . Now, take $0 + \mathfrak{a} = \mathfrak{a}$ as the additive identity and $-a + I$ as the inverse of $a + \mathfrak{a}$ for each a .

Now, notice, that $(a + \mathfrak{a})(b + \mathfrak{a}) = ab + a\mathfrak{a} + b\mathfrak{a} + \mathfrak{a}^2 = ab + (\mathfrak{a} + \mathfrak{a} + \mathfrak{a}) = ab + \mathfrak{a}$ by distribution of multiplication over addition in A . Moreover, A/\mathfrak{a} also inherits associativity and commutativity in \cdot from multiplication in A . Now, notice then

$$(a + \mathfrak{a})((b + \mathfrak{a}) + c + \mathfrak{a}) = (a + \mathfrak{a})((b + c) + \mathfrak{a}) = a(b + c) + \mathfrak{a} = (ab + ac) + \mathfrak{a} = (ac + \mathfrak{a}) + (bc + \mathfrak{a})$$

Observe also that if 1 is the identity of A , then $1 + \mathfrak{a}$ is the identity of A/\mathfrak{a} as $a + \mathfrak{a}$. Since $(a + \mathfrak{a})(1 + \mathfrak{a}) = a + \mathfrak{a}$.

Lastly, notice that $a + \mathfrak{a}$ is just the left coset of a by \mathfrak{a} in A as a group under addition. So that $+$ and \cdot are coset addition and multiplication, which are well defined. ■

Definition. Let A be a commutative ring with identity and \mathfrak{a} an ideal in A . We call the ring A/\mathfrak{a} under addition and multiplication of cosets the **factor ring** (or **quotient ring**) of A over \mathfrak{a} .

Theorem 1.3.3 (The First Isomorphism Theorem). If $\phi : A \rightarrow B$ is a ring homomorphism from rings A into B , then $\ker \phi$ is an ideal of A and

$$\begin{array}{ccc} & \phi(A) \simeq A/\ker \phi & \\ & \nearrow \bar{\phi} & \\ A & \xrightarrow{\phi} & B \\ \downarrow \pi & & \uparrow \\ A/\ker \phi & & \end{array}$$

Proof. By the first isomorphism theorem for groups, ϕ is a group isomorphism. Now, let $K = \ker \phi$ and consider the map $\pi : A \rightarrow A/\mathfrak{a}$ by $a \mapsto a + K$. Define the map $\bar{\phi} : A/K \rightarrow \phi(A)$ such that $\bar{\phi} \circ \pi = \phi$, then $\bar{\phi}$ defines the ring isomorphism. ■

Proof. The map $\pi : A \rightarrow A/\mathfrak{a}$ defined by $a \mapsto a + \mathfrak{a}$, for any ideal \mathfrak{a} , is onto, with $\ker \pi = \mathfrak{a}$. ■

Theorem 1.3.4 (The Second Isomorphism Theorem). *Let $\mathfrak{a} \subseteq A$ a subring of A , and let \mathfrak{b} an ideal in A . Define $\mathfrak{a} + \mathfrak{b} = \{a + b : a \in \mathfrak{a} \text{ and } b \in \mathfrak{b}\}$. Then $\mathfrak{a} + \mathfrak{b}A$ is a subring and $\mathfrak{a} \cap \mathfrak{b}$ is an ideal in A . Then*


$$\mathfrak{a}\mathfrak{b}/\mathfrak{b} \simeq \mathfrak{a}/\mathfrak{a} \cap \mathfrak{b}$$

Theorem 1.3.5 (The Third Isomorphism Theorem). *Let \mathfrak{a} and \mathfrak{b} be ideals in a ring A , with $\mathfrak{a} \subseteq \mathfrak{b}$. Then $\mathfrak{b}/\mathfrak{a}$ is an ideal of A/\mathfrak{a} and*

$$A/\mathfrak{b} = (A/\mathfrak{a})/(\mathfrak{b}/\mathfrak{a})$$

Theorem 1.3.6 (The Fourth Isomorphism Theorem). *Let \mathfrak{a} an ideal in a ring A , then the correspondence between A and A/\mathfrak{a} , for any subring of A is an inclusion preserving bijection between subrings of A containing \mathfrak{a} and A/\mathfrak{a} . Moreover, A is an ideal if, and only if A/\mathfrak{a} is an ideal.*

Lemma 1.3.7. *Let A be a ring with ideals \mathfrak{a} and \mathfrak{b} . Then $\mathfrak{a} + \mathfrak{b}$, $\mathfrak{a}\mathfrak{b}$ and \mathfrak{a}^n , for any $n \geq 0$ are ideals of A and we have the lattice*



1.4 Properties of Ideals

Definition. Let A be a commutative ring with identity. We call the smallest ideal containing a nonempty subset S in A the **ideal generated** by S , and we write (S) . We call an ideal **principle** if it is generated by a single element of A , i.e. $\mathfrak{a} = (a)$ for some $a \in \mathfrak{a}$. We say that the ideal (S) is **finitely generated** if $|S|$ is finite, and if $S = \{a_1, \dots, a_n\}$, then we denote $(S) = (a_1, \dots, a_n)$.

Example 1.2. (1) In any commutative ring with identity, the trivial ideal and A are the ideals generated by 0 and 1, respectively, so we write them as (0) and $A = (1)$.

(2) In \mathbb{Z} , we can write the ideals $n\mathbb{Z} = (n) = (-n)$. Notice that every ideal in \mathbb{Z} is a principle ideal. Moreover, for $m, n \in \mathbb{Z}$, $n|m$ if, and only if $n\mathbb{Z} \subseteq m\mathbb{Z}$. Notice that $m\mathbb{Z} + n\mathbb{Z} = d\mathbb{Z}$ is the ideal generated by m and n , where $d = (m, n)$ is the greatest

common divisor of m and n . Indeed, by definition, $d|m, n$ so that $d\mathbb{Z} \subseteq m\mathbb{Z} + n\mathbb{Z}$, and if $c|m, n$, then $c|d$, making $m\mathbb{Z} + n\mathbb{Z} \subseteq d\mathbb{Z}$. Then $m\mathbb{Z} + n\mathbb{Z} = d\mathbb{Z}$ is the ideal generated by the greatest common divisor (m, n) and consists of all diophantine equations of the form

$$mx + ny = (m, n)$$

In general, we can define the **greatest common divisor** for integers n_1, n_2, \dots, n_m to be the smallest such integer d generating the ideal $n_1\mathbb{Z} + \dots + n_m\mathbb{Z} = d\mathbb{Z}$. We then write $d = (n_1, \dots, n_m)$.

- (3) Let $m, n \in \mathbb{Z}$. Then the least common multiple of m, n , $[m, n]$ is $[m, n]\mathbb{Z} = m\mathbb{Z} \cap n\mathbb{Z}$. Indeed, if $c = [m, n]$ is the least common multiple of m, n , then we have that $m|c$ and $n|c$, making $c \in m\mathbb{Z} \cap n\mathbb{Z}$; similarly, for any $c' \in m\mathbb{Z} \cap n\mathbb{Z}$, $c|c'$, by definition which puts $c' \in c\mathbb{Z}$. In general, for $n_1, \dots, n_m \in \mathbb{Z}$, we define the **least common multiple** of n_1, \dots, n_m to be the largest such integer c generating the ideal $c\mathbb{Z} = n_1\mathbb{Z} \cap \dots \cap n_m\mathbb{Z}$. And we write $c = [n_1, \dots, n_m]$.
- (4) Let $m, n \in \mathbb{Z}^+$ be coprime, i.e. $(m, n) = 1$. Then we can obtain $mn = [m, n]$ by observing the ideals generated by mn , (m, n) , and $[m, n]$.
- (5) Consider the ideal $(2, x)$ of $\mathbb{Z}[x]$. $(2, x)$ is not a principle ideal. We have that $(2, x) = \{2p_xq : p, q \in \mathbb{Z}[x]\}$, and that $(2, x) \neq \mathbb{Z}[x]$. Suppose that $(2, x) = (a)$ for some polynomial $a \in \mathbb{Z}[x]$, then $2 \in (a)$, so that $2 = p(x)a(x)$, of degree $\deg p + \deg a$. This makes p and a constant polynomials in $\mathbb{Z}[x]$. Now, since 2 is prime in \mathbb{Z} , then only values for p and q are $p = \pm 1$ and $a = \pm 2$. If $a(x) = \pm 1$, then every polynomial in $\mathbb{Z}[x]$ can be written as a polynomial in (a) , so that $(a) = \mathbb{Z}[x]$, impossible. If $a(x) = \pm 2$, then since $x \in (a)$, we get $x = 2q(x)$ where $q \in \mathbb{Z}[x]$. This cannot happen, so that $(a) \neq (2, x)$.

Lemma 1.4.1. *Let \mathfrak{a} an ideal in ring A with identity. Then*

- (1) $\mathfrak{a} = (1)$ if, and only if \mathfrak{a} contains a unit.
- (2) If A is commutative, then A is a field if, and only if its only ideals are (0) and (1) .

Proof. Recall that $A = (1)$. Now, if $\mathfrak{a} = (1)$, then $1 \in \mathfrak{a}$, and 1 is a unit. Conversely, suppose that $u \in \mathfrak{a}$ with u a unit. By definition, we have that $r = r \cdot 1 = r(uv) = r(vu) = (rv)u$, so that $1 \in \mathfrak{a}$. This makes $\mathfrak{a} = (1)$.

Now, if A is a field, then it is a commutative ring, moreover every $r \neq 0$ is a unit in A , which makes $r \in \mathfrak{a}$ for some ideal $\mathfrak{a} \neq (0)$. This makes every $\mathfrak{a} \neq (0)$ equal to (1) . Conversely, if (0) and (1) are the only ideals of the commutative ring A , then every $r \neq 0 \in (1)$, which makes them units. Hence all nonzero r is a unit in A . This makes A into a field. ■

Corollary. *If k is a field, then any nonzero ring homomorphism ϕ defined on k is 1-1.*

Proof. If k is a field, then either $\ker \phi = (0)$ or $\ker \phi = (1)$. Now, since $\ker \phi \neq A$, we must have $\ker \phi = (0)$. ■

Definition. For any ideal \mathfrak{m} in a ring A , we call \mathfrak{m} **maximal** if $\mathfrak{m} \neq A$, and if \mathfrak{a} is an ideal with $\mathfrak{m} \subseteq \mathfrak{a} \subseteq A$, then either $\mathfrak{m} = \mathfrak{a}$ or $\mathfrak{a} = A$.

Lemma 1.4.2. *If A is a commutative ring with identity, every proper ideal is contained in a maximal ideal.*

Proof. Let \mathfrak{a} a proper ideal of A . Let $\mathcal{S} = \{N : N \neq (1) \text{ is a proper ideal, and } \mathfrak{a} \subseteq N\}$. Then $\mathcal{S} \neq \emptyset$, as $\mathfrak{a} \in \mathcal{S}$, and the relation \subseteq partially orders \mathcal{S} . Let \mathcal{C} be a chain in \mathcal{S} and define

$$J = \bigcup_{\mathfrak{a} \in \mathcal{C}} \mathfrak{a}$$

We have that $J \neq \emptyset$ since $(0) \in J$. Now, let $a, b \in J$, then we have that either $(a) \subseteq (b)$ or $(b) \subseteq (a)$, but not both. In either case, we have $a - b \in J$ so that J is closed under additive inverse. Moreover, since $\mathfrak{a} \in \mathcal{C}$ is an ideal, by definition, J is closed with respect to absorption. This makes J an ideal.

Now, if $1 \in J$, then $J = (1)$ and J is not proper, and $\mathfrak{a} = (1)$ by definition of J . This is a contradiction as \mathfrak{a} must be proper. Therefore J must also be a proper ideal. Therefore, \mathcal{C} has an upperbound in \mathcal{S} , therefore, by Zorn's lemma, \mathcal{S} has a maximal element \mathfrak{m} , i.e. it has a maximal ideal \mathfrak{m} with $\mathfrak{a} \subseteq \mathfrak{m}$. ■

Lemma 1.4.3. *Let A be a commutative ring with identity. An ideal \mathfrak{m} is maximal if, and only if A/\mathfrak{m} is a field.*

Proof. If \mathfrak{m} is maximal, then there is no ideal $I \neq (1)$ for which $\mathfrak{m} \subseteq \mathfrak{a} \subseteq A$. By the fourth isomorphism theorem, the ideals of A containing \mathfrak{a} are in 1-1 correspondence with the those of A/\mathfrak{m} . Therefore \mathfrak{m} is maximal if, and only if the only ideals of A/\mathfrak{m} are (\mathfrak{m}) and $(1+\mathfrak{m})$. ■

Example 1.3. (1) Let $n \geq 0$ an integer. The ideal $n\mathbb{Z}$ is maximal in \mathbb{Z} if and only if $\mathbb{Z}/n\mathbb{Z}$ is a field. Therefore $n\mathbb{Z}$ is maximal if, and only if $n = p$ a prime in \mathbb{Z} . So the maximal ideals of \mathbb{Z} are those $p\mathbb{Z}$ where p is prime.

(2) $(2, x)$ is not principal in $\mathbb{Z}[x]$, but it is maximal in $\mathbb{Z}[x]$, as $\mathbb{Z}[x]/(2, x) \simeq \mathbb{Z}/2\mathbb{Z}$ which is a field.

(3) The ideal (x) is not maximal in $\mathbb{Z}/n\mathbb{Z}$, since $\mathbb{Z}/(x) \simeq \mathbb{Z}$, which is not a field. Moreover, $(x) \subseteq (2, x) \subseteq \mathbb{Z}[x]$. We construct this isomorphism by identifying $x = 0$, then all polynomials of $\mathbb{Z}[x]/(x)$ only have constant term in \mathbb{Z} .

Definition. We call an ideal \mathfrak{p} in a commutative ring A with identity **prime** if $\mathfrak{p} \neq (1)$ and if $ab \in \mathfrak{p}$ then either $a \in \mathfrak{p}$ or $b \in \mathfrak{p}$. Alternatively, if $(ab) \subseteq \mathfrak{p}$ then $(a) \subseteq \mathfrak{p}$ or $(b) \subseteq \mathfrak{p}$.

Example 1.4. The prime ideals of \mathbb{Z} are $p\mathbb{Z}$ with p prime together with (0) .

Lemma 1.4.4. *An ideal \mathfrak{p} in a commutative ring with identity, A , is prime if, and only if A/\mathfrak{p} is an integral domain.*

Proof. Suppose that \mathfrak{p} is prime, and let $(a + \mathfrak{p})(b + \mathfrak{p}) = ab + \mathfrak{p} = \mathfrak{p}$. This gives us that $ab \in \mathfrak{p}$ and hence $a \in \mathfrak{p}$ or $b \in \mathfrak{p}$. Then either $a + \mathfrak{p} = \mathfrak{p}$ or $b + \mathfrak{p} = \mathfrak{p}$ in A/\mathfrak{p} . Conversely, if A/\mathfrak{p} is an integral domain, then for any $a + \mathfrak{p}, b + \mathfrak{p}$ $ab + \mathfrak{p} = \mathfrak{p}$ implies that either $a + \mathfrak{p} = \mathfrak{p}$ or $b + \mathfrak{p} = \mathfrak{p}$. Then $a \in \mathfrak{p}$ or $b \in \mathfrak{p}$, only when $ab \in \mathfrak{p}$. This makes \mathfrak{p} prime. ■

Corollary. *Every maximal ideal is a prime ideal.*

Example 1.5. (1) The prime ideals of \mathbb{Z} are $p\mathbb{Z}$, where p is prime, which are the maximal ideals of \mathbb{Z} .

(2) The ideal (x) in $\mathbb{Z}[x]$ is a prime ideal, but it is not maximal as $(x) \subseteq (2, x) \subseteq \mathbb{Z}[x]$.

Definition. Let A be a commutative ring with identity. We call A a **local ring** if it has one, and only one maximal ideal. We define the **residue field** of A to be the field $k = A/\mathfrak{m}$. We call a commutative ring with identity a **semi-local ring** if it has only finitely many maximal ideals.

Example 1.6. The ring \mathbb{Z} is not a local ring, it is not even semi-local, since every prime ideal (p) of \mathbb{Z} , where $p \in \mathbb{Z}^+$ is prime, is also maximal.

Lemma 1.4.5. *Let A be a commutative ring with identity. Then the following are true.*

- (1) *If $\mathfrak{m} \neq (1)$ is an ideal of A such that every element of $A \setminus \mathfrak{m}$ is a unit, then A is a local ring having \mathfrak{m} as its maximal ideal.*
- (2) *If \mathfrak{m} is a maximal ideal of A such that every element of $1 + \mathfrak{m}$ is a unit, then A is a local ring.*

Proof. Suppose that $\mathfrak{m} \neq (1)$. We have by lemma 1.4.2 that \mathfrak{m} is contained in a maximal ideal. Moreover, \mathfrak{m} contains no units by lemma 1.4.1. Since $x \in A \setminus \mathfrak{m}$ is a unit, we get $(x) = (1)$, which makes \mathfrak{m} the only maximal ideal of A and A is a local ring.

Now, suppose that \mathfrak{m} is maximal, and take $x \in A \setminus \mathfrak{m}$. Then the ideal $(x, \mathfrak{m}) = (1)$, so that there exists a $y \in A$, and $t \in \mathfrak{m}$ for which $xy - t = 1$; i.e. $xy = 1 - t$, which makes x a unit. By above, this makes A a local ring. ■

1.5 Euclidian Domains

Definition. Let A be a commutative ring with identity. We call a map $N : A \rightarrow \mathbb{N}$, with $N(0) = 0$ a **norm**, or, **degree**. If $N(a) \geq 0$, for all $a \in A$, then we call N **nonnegative**. If $N(a) > 0$ for all $a \in A$ then we call N **positive**.

Definition. Let A be a commutative ring with identity, and $N : A \rightarrow \mathbb{N}$ a norm. We say that A is a **Euclidean domain** if for all $a, b \in A$, with $b \neq 0$, there exist elements $q, r \in A$ such that

$$a = qb + r \text{ where } r = 0 \text{ or } N(r) < N(b)$$

We call q the **quotient** and r the **remainder** of a when **divided** by b .

Example 1.7. (1) Let k be any field, and $N : k \rightarrow \mathbb{N}$ defined by $N(a) = 0$ for all $a \in k$. Then N makes k into a Euclidean domain. Take $a, b \in k$, with $b \neq 0$, and $q = ab^{-1}$. Then $a = qb + r$ where $r = 0$.

(2) The integers \mathbb{Z} is a Euclidean domain with norm $N(a) = |a|$, the absolute value of a . In fact, the motivation for Euclidean rings comes from the division theorem, or Euclid's theorem for integers.

(3) Let k be a field, and consider $k[x]$. Let $N : k[x] \rightarrow \mathbb{N}$ be defined by $N : f \rightarrow \deg f$. Then $k[x]$ is a Euclidean domain. If k is not a field, then it is not necessarily true that $k[x]$ is a Euclidean domain.

(4) Let $D \in \mathbb{Z}^+$ be squarefree, and consider $\mathbb{Z}[\sqrt{D}]$. Define $N : \mathbb{Z}[\sqrt{D}] \rightarrow \mathbb{N}$ to be the absolute value of the field norm, that is $N(a + b\sqrt{D}) = \|a + b\sqrt{D}\|^2 = a^2 + Db^2$. We notice that $\mathbb{Z}[\sqrt{D}]$ is an integral domain, but it is not a Euclidean domain. This depends on our choice of D . Let $D = -1$ so that $\sqrt{D} = i$, and $i^2 = -1$. Then the Gaussian integers, $\mathbb{Z}[i]$, is a Euclidean domain. Let $x = a + ib$, $y = c + id$ with $y \neq 0$. In $\mathbb{Q}[i]$, the field of fractions, we have that $\frac{x}{y} = r + is$, where

$$r = \frac{ac + bd}{\|y\|^2} \text{ and } s = \frac{bc - ad}{\|y\|^2}$$

Now, let p and q be the integers closest to r and s , respectively so that

$$|r - p| \leq \frac{1}{2} \text{ and } |s - q| \leq \frac{1}{2}$$

Let $w = (r - p) + i(s - q)$, and take $z = wy$. Then we have $z = x - (p + iq)y$, so that $x = (p + iq)y + z$, moreover, we have $N(w) = (r - p)^2 + (s - q)^2 \leq \frac{1}{4} + \frac{1}{4} = \frac{1}{2}$. Since $\|\cdot\|$ is multiplicative, we have

$$N(w)N(y) \leq \frac{1}{2}N(y)$$

which makes $\mathbb{Z}[i]$ into a Euclidean domain.

Lemma 1.5.1. *Every ideal in a Euclidean domain A , is a principle ideal.*

Proof. If $I = (0)$, we are done. Now, let $N : A \rightarrow \mathbb{N}$ be the norm of A , and consider the image $N(I) = \{N(a) : a \in I\}$. By the well ordering principle, $N(I)$ has a minimum element $N(d)$ for some $d \neq 0$ in I . Notice that $(d) \subseteq I$. Now, let $a \in I$. Since A is a Euclidean domain, there exist $q, r \in A$ for which

$$a = qd + r \text{ where } r = 0 \text{ or } N(r) < N(d)$$

Then notice that

$$r = a - qd$$

putting $r \in I$ and $N(r) \in N(I)$. Since $N(d)$ is the minimum element, we must have $r = 0$ so that $a = qd$, which puts $a \in (d)$. Therefore $I = (d)$, making I principle. ■

Example 1.8. (1) The polynomial ring $\mathbb{Z}[x]$ is not a Euclidean domain. The ideal $(2, x)$ is not principle.

- (2) Consider $\mathbb{Z}[\sqrt{-5}]$, i.e. $D = -5$. Suppose the ideal $(3, 2 + \sqrt{-5})$ is a principle ideal, that is $(3, 2 + \sqrt{-5}) = (a + b\sqrt{-5})$ for some $a, b \in \mathbb{Z}$. Then we get that $3 = x(a + b\sqrt{-5})$ and $2 + \sqrt{-5} = y(a + b\sqrt{-5})$. Then notice that $N(x) = a^2 + 5b^2 = 9$, and since $a^2 + 5b^2 \in \mathbb{Z}^+$, we must have that $a^2 + 5b^2 = 1, 3, 9$.
- (i) If $a^2 + 5b^2 = 9$, then $N(x) = 1$ making $x = \pm 1$ and $a + b\sqrt{-5} = \pm 3$, which cannot happen since $2 + \sqrt{-5}$ is not divisible by 3.
 - (ii) the equation $a^2 + 5b^2 = 3$ cannot happen since it has no integer solutions. This makes
 - (iii) $a^2 + b^2 = 1$, which makes $(a + b\sqrt{-5}) = \mathbb{Z}[\sqrt{-5}]$, moreover, we get the equation $3x + y(2 + \sqrt{-5}) = 1$ for any $x, y \in \mathbb{Z}[\sqrt{-5}]$. Multiplying both sides by $2 - \sqrt{-5}$, we get that $3|(2 - \sqrt{-5})$ which is impossible.

In all three cases, we were led to an impossibility, hence $\mathbb{Z}[\sqrt{-5}]$ cannot be a Euclidean domain.

Definition. Let A be a commutative ring with identity, and $a, b \in A$ with $b \neq 0$. We say that b **divides** a if there is an $x \in A$ for which $a = bx$. We write $b|a$. We also say that a is a **multiple** of b .

Definition. Let A be a commutative ring with identity. We call a nonzero element $d \in A$ a **greatest common divisor** of elements $a, b \in A$ if

- (1) $d|a$ and $d|b$.
- (2) If $c \in A$ is nonzero such that $c|a$ and $c|b$, then $c|d$.

We write $d = (a, b)$.

Lemma 1.5.2. *Let A be a commutative ring with identity. For any $a, b \in A$ a nonzero element $d \in A$ is the greatest common divisor if*

- (1) $(a, b) \subseteq (d)$.
- (2) If $c \in A$ is nonzero with $(a, b) \subseteq (c)$, then $(d) \subseteq (c)$.

In particular, $d = (a, b)$.

Proof. The first two statements follow from definition, and the last follows lemma 1.5.1. ■

Lemma 1.5.3. *If A is a commutative ring with identity, and $a, b \in A^*$, such that $(a, b) = (d)$ for some $d \in A^*$, then d is the greatest common divisor of a and b .*

Lemma 1.5.4. *Let A be an inetegral domain. If $c, d \in A$ generate the same principle ideal, i.e. $(d) = (c)$, then $d = uc$ for some unit $u \in A$.*

Proof. If $c = 0$ or $d = 0$, we are done. Suppose then that $c, d \neq 0$. Since $(d) = (c)$, we have that $d = xc$ and $c = yd$ for some $x, y \in A$. Then $d = (xy)d$, which makes $d(1 - xy) = 0$. Since $d \neq 0$, we get $xy = 1$, making x and y units of A . ■

Definition. We call an integral domain in which every principle ideal is generated by two elements a **Bezout domain**.

Lemma 1.5.5. *Every Euclidean domain is a Bezout domain.*

Theorem 1.5.6 (The Extended Euclidean Algorithm). *Let A be a Euclidean and $a, b \neq 0$ elements of A . Let $d = r_n$ be the least nonzero remainder obtained by dividing a from b recursively $n + 1$ times. Then*

- (1) $d = (a, b)$ is the greatest common divisor of a and b .
- (3) There exist $x, y \in A$ for which $ax + by = d$.

Proof. By lemma 1.5.1, we get that the ideal (a, b) is principle, so there exists a greatest common divisor of a and b . Now, let $d = r_n$ be obtained by dividing a and b recursively $(n + 1)$ times. Then the $(n + 1)^{st}$ equation gives $r_{n-1} = q_{n+1}r_n$, so that $r_n | r_{n-1}$. Now, by induction on n if $r_n | r_{k+1}$ and $r_n | r_k$ then the $(k + 1)^{st}$ equation gives $r_{k-1} = q_{k+1}r_k + r_{k+1}$, which implies that $r_n | r_{k-1}$. Therefore we get in the 1^{st} equation that $r_n | b$, and in the 0^{th} that $r_n | a$. That is, $d | a$ and $d | b$.

Now, notice that $r_0 \in (a, b)$ and that $r_1 = b - qr_0 \in (b, r_0) \subseteq (a, b)$. By induction on r_n , if $r_{k-1}, r_n \in (a, b)$ then

$$r_{k+1} = r_{k-1} - q_{k+1}r_k \in (r_{k-1}, r_n) \subseteq (a, b)$$

which puts $r_n \in (a, b)$ making $d = (a, b)$ the greatest common divisor. ■

1.6 Principle Ideal Domains.

Definition. An integral domain A is called a **principle ideal domain (PID)** if every ideal in A is principle.

Example 1.9. (1) Every Euclidean domain is a PID, as dictated by lemma ???. Hence the rings \mathbb{Z} and $\mathbb{Z}[i]$ are PIDs, however, the polynomial ring $\mathbb{Z}[x]$ is not principle, recall the ideal $(2, x)$.

- (2) The ring $\mathbb{Z}[\sqrt{-5}]$ is not a PID, consider the ideal $(3, 2 + \sqrt{-5})$. However, notice that $(3, 1 + \sqrt{-5})(3, 1 - \sqrt{-5}) = (3)$ is principle, despite $(3, 1 + \sqrt{-5})$ and $(3, 1 - \sqrt{-5})$ are not principle.

- (3) The ring $\mathbb{Z}[1 + \frac{\sqrt{-19}}{2}]$ is a PID, but not a Euclidean domain.

Lemma 1.6.1. *Let A be a principle ideal domain and let d be a generator for the ideal (a, b) , for $a, b \in A$. Then the following are true.*

- (1) $d = (a, b)$; i.e. d is the greatest common divisor of a and b .
- (2) There exist $x, y \in A$ for which $ax + by = d$.
- (3) d is unique up to unit.

Lemma 1.6.2. *Every nonzero prime ideal in a principle ideal domain A is maximal.*

Proof. Let $(p) \neq (0)$ be a prime ideal of A . Let (m) be an ideal of A containing (p) . Then $p \in (m)$ so that $p = rm$ for some $r \in A$. Now, since p is prime, and $rm \in (p)$, then either $r \in (p)$ or $m \in (p)$. If $m \in (p)$, then $(p) = (m)$. Otherwise, if $r \in (p)$, then $r = ps$ for some $s \in A$. Then $p = rm = pms = p(ms)$ which makes $ms = 1$, hence m is a unit, which makes $(m) = (0)$. ■

Corollary. *If A is any commutative ring, such that the polynomial ring $A[x]$ is a principle ideal domain, then A is necessarily a field.*

Proof. If $A[x]$ is a PID, then $A \subseteq A[x]$, as a subring, must be an integral domain. Consider now, the ideal (x) , then $A[x]_{(x)} \simeq A$ which makes (x) prime by lemma 1.4.4. Therefore (x) is maximal, which then makes A a field by lemma 1.4.3. ■

Definition. Let A be a commutative ring, and $N : A \rightarrow \mathbb{N}$ a norm. We call N a **Dedekin-Hasse norm** if N is a positive norm such that for all $a, b \in N$, either $a \in (b)$, or there exists an element $c \in (a, b)$ such that $N(c) < N(b)$.

Lemma 1.6.3 (The Dedekin-Hasse Criterion). *An integral domain A is a PID if, and only if it has a Dedekin-Hasse norm.*

Proof. Let $\mathfrak{b} \neq (0)$ an ideal of A . Let $a \in \mathfrak{b}$ a nonzero element, so that $(a, b) \subseteq \mathfrak{b}$. Since N is Dedekin-Hasse, and by minimality of b , we get that $a \in (b)$ so that $\mathfrak{b} = (b)$ is principle. ■

Example 1.10. Consider the ring $\mathbb{Z}[1 + \frac{\sqrt{-19}}{2}]$. With norm $N = \|\cdot\|^2$ the field norm. Let $x, y \in \mathbb{Z}[1 + \frac{\sqrt{-19}}{2}]$ be nonzero elements and that $\frac{x}{y} \notin \mathbb{Z}[1 + \frac{\sqrt{-19}}{2}]$. Write

$$\frac{x}{y} = \frac{a + b\sqrt{-19}}{c} \in \mathbb{Q}[1 + \frac{\sqrt{-19}}{2}]$$

where a, b, c are all coprime, with $c > 1$. Then there are integers u, v, w with $av + bu + cw = 1$, then $au - 19bv = cq + r$ for some quotient q and remainder r with $N(r) \leq \frac{c}{2}$ and let $s = u + v\sqrt{-19}$ and $t = q - w\sqrt{-19}$. Then we find that

$$0 < N(\frac{x}{y}s - t) \leq \frac{1}{4} + \frac{19}{c^2}$$

Then $s = 1, t = \frac{(a-1)+b\sqrt{-19}}{2} \in A$ satisfy $0 < N(\frac{x}{y}s - t)$

Now, suppose that $c = 3$, then $3 \nmid (a^2 + 19b^2)$. Then $a^2 + 19b^2 = 3q + r$ with $r = 1$ or $r = 2$. Then $s = a - b\sqrt{-19}, t = q$ satisfy $0 < N(\frac{x}{y}s - t)$. Finally, for $c = 4$, with a, b not both even, so that $a^2 + 19b^2$ is odd. Then $a^2 + 19b^2 = 4q + r$ so for $q, r \in \mathbb{Z}$ with $0 < r < 4$, then $s = a - b\sqrt{-19}, t = q$ satisfy $0 < N(\frac{x}{y}s - t)$. Now, if both a and b are odd, then $a^2 + 19b^2 \equiv 1 + 3 \pmod{8}$ so that $a^2 + 19b^2 = 8q + 4$ for some $q \in \mathbb{Z}$, then

$$s = \frac{a - b\sqrt{-19}}{2} \text{ and } t = q$$

satisfy $0 < N(\frac{x}{y}s - t)$.

1.7 Unique Factorization Domains.

Definition. Let A be an integral domain. A nonzero element $r \in A$ that is not an associate is called **irreducible** if whenever $r = ab$, then either a or b are units in A ; otherwise, we call r **reducible**.

Definition. Let A be an integral domain. An element $p \in A$ is called **prime** if the ideal (p) is a prime ideal. That is p is not a unit and whenever $p|ab$, then either $p|a$ or $p|b$. We call two elements $a, b \in A$ **associates** if $a = ub$ for some unit $u \in A$.

Lemma 1.7.1. *In an integral domain, a prime element is always irreducible.*

Proof. Let (p) be a nonzero prime ideal with $p = ab$, for some $a, b \in A$. Then $ab \in (p)$, so that either $a \in (p)$, or $b \in (p)$. Suppose that $a \in (p)$. Then $a = pr$ for some $r \in A$, so that $p = (pr)b = p(rb)$, so that $rb = 1$. This makes b a unit. Similarly, we see that a is a unit if $b \in (p)$. In either case, p is irreducible. ■

Example 1.11. (1) In the ring \mathbb{Z} of integers, those elements which are irreducible are precisely those which are prime, since the ideals $2\mathbb{Z}, 3\mathbb{Z}, \dots, p\mathbb{Z}, \dots$, for p a prime number are also the prime ideals of \mathbb{Z}

(2) Irreducible elements need not be prime. The element $3 \in \mathbb{Z}[\sqrt{-5}]$ is irreducible, as was shown in example 1.8, however it is not prime. Notice that $3|9 = (2 + \sqrt{-5})(2 - \sqrt{-5})$, but $3 \nmid (2 + \sqrt{-5})$ and $3 \nmid (2 - \sqrt{-5})$.

Lemma 1.7.2. *In a principle ideal domain, a nonzero element is prime if, and only if it is irreducible.*

Proof. Let A be a PID, and suppose that p is irreducible. Let (m) be the principle ideal containing (p) , then $p = rm$, and by irreducibility, either r or m are units, in either case, we get that either $(p) = (m)$ or $(m) = (1)$. This makes (p) a maximal ideal, and hence a prime ideal. ■

Example 1.12. (1) Since 3 is not prime in $\mathbb{Z}[\sqrt{-5}]$, then (3) is not a prime ideal in this ring. Therefore $\mathbb{Z}[\sqrt{-5}]$ cannot be a PID.

(2) Notice that since \mathbb{Z} is a PID, then the fact that irreducible and prime elements coincide is guaranteed by lemma 1.7.2.

Definition. We call an integral domain A a **unique factorization domain (UFD)** if for every nonzero element $r \in A$ which is not a unit, the following are true.

- (1) r can be written as the product of, not necessarily distinct, irreducible elements. We call this product the **factorization** of r .
- (2) The factorization of r is unique up to associates.

Example 1.13. (1) All fields are unique factorization domains.

- (2) Polynomial rings are unique factorization domains whenever the ground ring A is a unique factorization domain.
- (3) The subring $\mathbb{Z}[2i]$ of $\mathbb{Z}[i]$ is an integral domain, but it is not a UFD. Notice that both 2 and $2i$ are irreducible in $\mathbb{Z}[2i]$, but that $4 = 2 \cdot 2 = (2i) \cdot (-2i)$.
- (4) $\mathbb{Z}[\sqrt{-5}]$ is another example of an integral domain that is not a UFD.

Lemma 1.7.3. *In a unique factorization domain A , a nonzero element is prime if, and only if it is irreducible.*

Proof. Since prime elements are irreducible, it remains to show that irreducible elements are prime. Let p be irreducible and suppose that $p|ab$, for $a, b \in A$. Then $ab = pc$ for some $c \in A$. Writing ab as a product of irreducibles, since A is a UFD, p must be associate to one of the irreducibles in the factorization of a , or to one in the factorization of b . In either case, we get that $p|a$ or $p|b$, and hence p is prime. ■

Lemma 1.7.4. *Let $a, b \in A$ nonzero elements of a unique factorization domain A . If $a = up_1^{e_1} \dots p_n^{e_n}$ and $b = vp_1^{f_1} \dots p_n^{f_n}$, where $u, v \in A$ are units, then the element*

$$d = p_1^{\min\{e_1, f_1\}} \dots p_n^{\min\{e_n, f_n\}}$$

is the greatest common divisor of a and b .

Proof. Notice that by definition of d , that $d|a$ and $d|b$. Now, let c be a common divisor of a and b with the unique prime factorization $c = q_1^{g_1} \dots q_m^{g_m}$. Since $q_i|c$ for each $1 \leq i \leq m$, then $q_i|p_j$ for each prime factor in the factorizations of a and b . Since both q_i and p_j are irreducible, they are associates. That implies that the primes of c are the primes of a and b . Moreover notice that since each $g_i \leq e_i, f_i$, that $c|d$, and so $d = (a, b)$. ■

Definition. Let A be a principle ideal domain. Let $\{a_n\}$ a sequence of elements of A . We call the increasing sequence of ideals $\{(a_n)\}$ an **infinite ascending chain** of ideals in A and write

$$(a_1) \subseteq (a_2) \subseteq \dots \subseteq (a_n) \subseteq \dots \subseteq A$$

We say that the infinite ascending chain $\{(a_n)\}$ **stabilizes** if for some $k \geq n$, we have $(a_n) = (a_k)$.

Lemma 1.7.5. *In any principle ideal domain, infinite ascending chains of ideals stabilize.*

Proof. Let $\mathfrak{a}_1 \subseteq \mathfrak{a}_2 \subseteq \dots \subseteq A$ an infinite ascending chain of ideals and let $\mathfrak{a} = \bigcup \mathfrak{a}_k$. Then \mathfrak{a} is an ideal in A , and since A is a PID, $\mathfrak{a} = (a)$ for some $a \in A$. This makes $a \in \mathfrak{a}_n$ for some n , and hence $\mathfrak{a}_n \subseteq \mathfrak{a}$. This makes $\mathfrak{a}_n = \mathfrak{a}$ for some $n \geq 1$, and hence this chain stabilizes. ■

Theorem 1.7.6. *Every principle ideal domain is a unique factorization domain.*

Proof. Let A be a PID, and $r \in A$ a nonzero element which is not a unit. If r is irreducible, we are done. Otherwise, we have $r = r_1 r_2$ for some $r_1, r_2 \in A$. Now, if both r_1 and r_2 are irreducible, we are done. Suppose then, without loss of generality, that r_1 is reducible. Then

$r_1 = r_{11}r_{12}$, and if both r_{11} and r_{12} are irreducible, we are done. Suppose then that r_{11} is reducible; continuing this process, we arrive at an infinite ascending chain of ideals

$$(r) \subseteq (r_1) \subseteq (r_{11}) \subseteq \cdots \subseteq A$$

and since A is a PID, this chain stabilizes. Thus r can be factored into irreducible elements; since this process terminates.

Now, by induction on n , for $n = 0$, we notice that r is a unit, and we are done. Suppose, then for $n \geq 1$, that $r = p_1 \cdots p_n = q_1 \cdots q_m$ for some $m \geq n$, and where each p_i and q_j are (not necessarily distinct) irreducibles for all $1 \leq i \leq n$ and $1 \leq j \leq m$. Notice that $p_1 | q_1 \cdots q_m$, and so $p_1 | q_j$ for some j . This makes p_1 and q_j associates; i.e. $q_j = p_1 u$, with $u \in A$ a unit. Cancelling the p_1 from both sides of the equation, we get $p_2 \cdots p_n = q_1 \cdots q_{j-1} q_{j+1} \cdots q_m$. Repeating this process, we get a 1–1 correspondence between associates, and hence the factorization of r is unique up to associates. Therefore A is a UFD. ■

Corollary. *Every Euclidean domain is a unique factorization domain.*

Proof. Notice that Euclidean domains are PIDs by lemma ??.

Corollary (The Fundamental Theorem of Arithmetic). *\mathbb{Z} is a unique factorization domain.*

Proof. Notice that \mathbb{Z} is a Euclidean domain.

Corollary. *There exists a multiplicative Dedekind-Hasse norm on A .*

Proof. If A is a PID, then the theorem tells us it is a UFD. Define the norm $N : A \rightarrow \mathbb{N}$ by taking $0 \rightarrow 0$, $u \rightarrow 1$ if u is a unit, and $a \rightarrow 2^n$ where $a = p_1 \cdots p_n$, where each p_i is irreducible. Notice that for every $a, b \in A$, $N(ab) = N(a)N(b)$. Now, suppose further that $a, b \neq 0$ and consider the ideal $(a, b) = (r)$, for some $r \in A$. UIf $a \notin (b)$, neither is r , and hence $b \nmid r$. Now, since $b = xr$, $x \in A$, then x cannot be a unit in A , so that $N(b) = N(xr) = N(x)N(r) > N(r)$. This completes the proof. ■

1.8 The Nilradical and Jacobson Radical

Theorem 1.8.1 (The Binomial Theorem). *Let A be a commutative ring with identity. Then for all $x, y \in A$, and $n \in \mathbb{Z}^+$*

$$(x + y)^n = \sum_{r+s=n} \binom{n}{r} x^r y^s$$

Lemma 1.8.2. *Let A be a commutative ring with identity, and \mathfrak{N} the set of all nilpotent elements of A . Then \mathfrak{N} is an ideal of A .*

Proof. If $x \in \mathfrak{N}$, then there is an $n \in \mathbb{Z}^+$ for which $x^n = 0$, notice that this also implies that $(-x)^n = 0$, so that $-x \in \mathfrak{N}$. Now, let $x, y \in \mathfrak{N}$. Then for some $m, n \in \mathbb{Z}^+$, we have $x^m = 0$ and $y^n = 0$. Consider then $(x + y)^{m+n-1}$. By the binomial theorem, we have

$$(x + y)^{m+n-1} = \sum_{r+s=m+n-1} \binom{m+n-1}{r} x^r y^s$$

Now, since $r + s = m + n - 1$, notice that either $r < m$, or $s < n$, but not both. This makes $x^r y^s = 0$ for all r, s , so that $(x + y)^{m+n-1} = 0$. This makes \mathfrak{R} a subgroup of A . Lastly, notice that if $a \in A$, and $x \in \mathfrak{R}$, then for some $n \in \mathbb{Z}^+$, $ax^n = (ax)^n = 0$, which makes \mathfrak{R} an ideal. ■

Corollary. A/\mathfrak{R} has no nonzero nilpotent elements.

Proof. Let $x \in A$ be nilpotent, then $x \in \mathfrak{R}$, so that $x + \mathfrak{R} = \mathfrak{R}$ in A/\mathfrak{R} . Therefore, the only nilpotent element of A/\mathfrak{R} is \mathfrak{R} itself. ■

Definition. We define the **nilradical** of a commutative ring A , with identity, to be the ideal, $\text{Nil } A$, consisting of all nilpotent elements of A .

Lemma 1.8.3. *Let A be a commutative ring with identity. Then $\text{Nil } A$ is the intersection of all prime ideals of A ; i.e.*

$$\text{Nil } A = \bigcap_{\mathfrak{p} \subseteq A} \mathfrak{p} \text{ where } \mathfrak{p} \text{ is a prime ideal of } A$$

Proof. Let \mathfrak{R} be the intersection of all prime ideals of A . Suppose that $x \in A$ is nilpotent. Then $x^n = 0$ for some $n \in \mathbb{Z}^+$, so that $x^n \in \mathfrak{p}$. Since \mathfrak{p} is prime, $x \in \mathfrak{p}$, which puts $x \in \mathfrak{R}$.

Conversely, suppose that $x \in A$ is not nilpotent, and let Σ be the set of all ideals \mathfrak{a} for which $x \notin \mathfrak{a}$. Notice that since 0 is nilpotent in A , $0 \in \Sigma$, so that Σ is nonempty. Therefore, by Zorn's lemma, Σ has a maximal element \mathfrak{p} . We claim that this \mathfrak{p} is prime. Suppose that $a, b \notin \mathfrak{p}$, then $\mathfrak{p} \subseteq \mathfrak{p} + (a)$ and $\mathfrak{p} \not\subseteq \mathfrak{p} + (b)$. So $\mathfrak{p} + (a), \mathfrak{p} + (b) \notin \Sigma$, by the maximality of \mathfrak{p} . This puts $x^m \in \mathfrak{p} + (a)$ and $x^n \in \mathfrak{p} + (b)$, for some $n, m \in \mathbb{Z}^+$. Thus $x^{m+n} \in \mathfrak{p} + (ab) \not\subseteq \Sigma$. Therefore, $ab \notin \mathfrak{p}$, which makes \mathfrak{p} a prime ideal for which $x \notin \mathfrak{p}$; i.e. $x \notin \mathfrak{R}$. ■

Definition. We define the **Jacobson radical** of a commutative ring A , with identity, to be the intersection of all maximal ideals of A . We denote it by $\text{Jac } A$.

Lemma 1.8.4. *Let A be a commutative ring with identity. Then $x \in \text{Jac } A$ if, and only if $1 - xy$ is a unit in A for some $y \in A$.*

Proof. Suppose that $x \in \text{Jac } A$, but that $1 - xy$ is not a unit of A . Then by lemma 1.4.2, we have $(1 - xy) \subseteq \mathfrak{m}$ for some maximal ideal \mathfrak{m} of A ; hence $1 - xy \in \mathfrak{m}$. However, since $x \in \text{Jac } A$, then $xy \in \mathfrak{m}$, which puts $1 \in \mathfrak{m}$, so that $\mathfrak{m} = (1)$ which contradicts that \mathfrak{m} is maximal. Therefore, $1 - xy$ must be a unit.

Conversely, suppose that $x \in \mathfrak{m}$ for some maximal ideal \mathfrak{m} of A . Then $(\mathfrak{m}, x) = (1)$ so that $u + xy = 1$ for some $u \in \mathfrak{m}$ and $y \in A$. This makes $1 - xy \in \mathfrak{m}$ so that $1 - \mathfrak{m}$ is not a unit. ■

1.9 Operations on Ideals

We observe some additional properties, of ideals, namely, concerning operations on ideals. For this section, assume we are working over a commutative ring A with identity, unless otherwise specified.

Lemma 1.9.1. *Let A be a commutative ring with identity, and let \mathfrak{a} , and \mathfrak{b} ideals of A . Then the following are true*

- (1) $\mathfrak{a} + \mathfrak{b}$ is the smallest ideal of A containing both \mathfrak{a} and \mathfrak{b} .
- (2) $\mathfrak{a}\mathfrak{b} = \{\sum x_i y_i : x_i \in \mathfrak{a} \text{ and } y_i \in \mathfrak{b}\}$.
- (3) $\mathfrak{a}\mathfrak{b} \subseteq \mathfrak{a} \cap \mathfrak{b}$, and $\mathfrak{a} \cap \mathfrak{b}$ is the largest ideal contained in both \mathfrak{a} and \mathfrak{b} .

Lemma 1.9.2. *Sums, intersections, and products of ideals in a commutative ring with identity are commutative, and associative. Moreover, the product of ideals distributes over the sum of ideals. That is, if \mathfrak{a} , \mathfrak{b} , and \mathfrak{c} are ideals, then*

$$\mathfrak{a}(\mathfrak{b} + \mathfrak{c}) = \mathfrak{a}\mathfrak{b} + \mathfrak{a}\mathfrak{c}$$

Lemma 1.9.3. *For any ideals \mathfrak{a} , \mathfrak{b} , and \mathfrak{c}*

$$\mathfrak{a} \cap (\mathfrak{b} + \mathfrak{c}) = \mathfrak{a} \cap \mathfrak{b} + \mathfrak{a} \cap \mathfrak{c} \text{ if } \mathfrak{b} \subseteq \mathfrak{a} \text{ or } \mathfrak{c} \subseteq \mathfrak{a}$$

Definition. We call two ideals \mathfrak{a} and \mathfrak{b} **coprime**, or **comaximal** if $\mathfrak{a} + \mathfrak{b} = (1)$.

Lemma 1.9.4. *The following are true for anyu ideals \mathfrak{a} and \mathfrak{b} .*

- (1) if \mathfrak{a} and \mathfrak{b} are coprime, then $\mathfrak{a} \cap \mathfrak{b} = \mathfrak{a}\mathfrak{b}$.
- (2) \mathfrak{a} and \mathfrak{b} are coprime if, and only if there exists an $x \in \mathfrak{a}$ and a $y \in \mathfrak{b}$ for which $x + y = 1$.

Corollary. *If $m, n \in \mathbb{Z}^+$ are coprime then their ideals $n\mathbb{Z}$ and $m\mathbb{Z}$ are coprime.*

Definition. Let $\{A_\alpha\}$ be a (not necessarily countable) collection of commutative rings with identity. We define the **direct product** of $\{A_\alpha\}$ to be the set

$$A = \prod_{\alpha} A_{\alpha}$$

Lemma 1.9.5. *Let $\{A_\alpha\}$ be a collection of commutative rings with identity. Then the direct product of $\{A_\alpha\}$ forms a commutative ring with identity under componentwise addition and componentwise multiplication.*

Lemma 1.9.6. *Let A be a commutative ring, and $\mathfrak{a}_1, \dots, \mathfrak{a}_n$ ideals of A . Define the ring homomorphism $\phi : A \rightarrow \prod A/\mathfrak{a}_i$ by*

$$\phi : x \rightarrow (x + \mathfrak{a}_1, \dots, x + \mathfrak{a}_n)$$

Then the following are true.

- (1) If \mathfrak{a}_i and \mathfrak{a}_j are coprime whenever $i \neq j$, then

$$\prod \mathfrak{a}_i = \bigcap \mathfrak{a}_i$$

- (2) ϕ is onto if, and only if \mathfrak{a}_i and \mathfrak{a}_j are coprime whenever $i \neq j$.

(3) ϕ is 1-1 if, and only if

$$\bigcap \mathfrak{a}_i = (0)$$

Proof. By induction on n it was shown that for $n = 2$ that if $\mathfrak{a}_1, \mathfrak{a}_2$ are coprime, then $\mathfrak{a}_1 \mathfrak{a}_2 = \mathfrak{a}_1 \cap \mathfrak{a}_2$. Now, suppose that

$$\mathfrak{b} = \prod_{i=1}^n \mathfrak{a}_i = \bigcap_{i=1}^n \mathfrak{a}_i$$

for all $n \geq 2$ and consider the case for $n + 1$. Since $\mathfrak{a}_i + \mathfrak{a}_{n+1} = (1)$ (they are coprime by hypothesis), we have $x_i + y_i = 1$ where $x_i \in \mathfrak{a}_i$ and $y_i \in \mathfrak{a}_{n+1}$. Hence notice that

$$\prod_{i=1}^n x_i = \prod_{i=1}^n (1 - y_i) \equiv 1 \pmod{\mathfrak{a}_{n+1}}$$

so that $\mathfrak{b} + \mathfrak{a}_{n+1} = (1)$. Hence $\mathfrak{b} \mathfrak{a}_{n+1} = \mathfrak{b} \cap \mathfrak{a}_{n+1}$ which completes the proof.

Suppose now, that ϕ is onto. Then there exists an $x \in A$ such that $\phi(x) = (1, 0, \dots, 0)$ so that $x \equiv 1 \pmod{\mathfrak{a}_1}$ and $x \equiv 0 \pmod{\mathfrak{a}_i}$ for $i > 1$. Hence $1 = (1 - x) + x \in \mathfrak{a}_1 + \mathfrak{a}_i$ for all $i > 1$. This makes \mathfrak{a}_1 and \mathfrak{a}_i coprime. We can repeat this argument for any index $j \neq i$. Conversely suppose that \mathfrak{a}_1 and \mathfrak{a}_i are coprime. Then $\mathfrak{a}_1 + \mathfrak{a}_i = (1)$ for all $i > 1$ and we have $u_i + v_i = 1$ for some $u_i \in \mathfrak{a}_1$ and $v_i \in \mathfrak{a}_i$. Take then $x = \prod v_i$. Then

$$x = \prod (1 - u_i) \equiv 1 \pmod{\mathfrak{a}_1} \text{ and } x \equiv 0 \pmod{\mathfrak{a}_i}$$

thus $\phi(x) = (1, 0, \dots, 0)$. repeating for each index $j \neq i$, we get that ϕ is onto. Finally, observe that

$$\ker \phi = \{x \in A : (x + \mathfrak{a}_1, \dots, x + \mathfrak{a}_n) = (\mathfrak{a}_1, \dots, \mathfrak{a}_n)\} = \bigcap_{i=1}^n \mathfrak{a}_i$$

Which gives us the equivalent condition for ϕ to be 1-1. ■

Lemma 1.9.7. *The following are true for any commutative ring with identity.*

(1) *If $\mathfrak{p}_1, \dots, \mathfrak{p}_n$ are prime ideals, containing an ideal \mathfrak{a} , then $\mathfrak{a} \subseteq \mathfrak{p}_i$ for some $1 \leq i \leq n$.*

(2) *If $\mathfrak{a}_1, \dots, \mathfrak{a}_n$ are ideals, and \mathfrak{p} is a prime ideal containing $\bigcap \mathfrak{a}_i$, then $\mathfrak{a}_i \subseteq \mathfrak{p}$ for some $1 \leq i \leq n$.*

Proof. For the first assertion, the result is vacuously true for $n = 1$. Now suppose the result is true for all $n \geq 1$. Then for every $1 \leq i \leq n$, there is an $x_i \in \mathfrak{a}$ such that $x_i \in \mathfrak{p}_j$ whenever $i \neq j$. Now, if $x_i \notin \mathfrak{p}_i$, we are done. Otherwise, $x_i \in \mathfrak{p}_i$, and consider

$$y = \sum_{i=1}^n x_1 x_2 \dots x_{i-1} x_{i+1} \dots x_n$$

Then $y \in \mathfrak{a}$ but $y \notin \mathfrak{p}_i$, which puts $\mathfrak{a} \not\subseteq \mathfrak{p}_i$, for all $1 \leq i \leq n$, hence $\mathfrak{a} \subseteq \mathfrak{p}_{n+1}$ and we are done.

For the second assertion, suppose that $\mathfrak{a}_i \not\subseteq \mathfrak{p}$ for all $1 \leq i \leq n$. Then let $x_i \in \mathfrak{a}$ such that $x_i \notin \mathfrak{p}$. Then we have

$$\prod \xi_i \in \mathfrak{a}_i$$

but $\prod x_i \notin \mathfrak{p}$, hence $\mathfrak{a}_i \not\subseteq \mathfrak{p}$. ■

Corollary. If $\mathfrak{p} = \bigcap \mathfrak{a}_i$, then $\mathfrak{p} = \mathfrak{a}_i$ for some $1 \leq i \leq n$.

Definition. Let \mathfrak{a} and \mathfrak{b} be ideals. We define the **ideal quotient** of \mathfrak{a} and \mathfrak{b} to be the set

$$(\mathfrak{a} : \mathfrak{b}) = \{x \in A : x\mathfrak{b} \subseteq \mathfrak{a}\}$$

We define the **annihilator** of \mathfrak{b} to be $(0 : \mathfrak{b})$ and denote it $\text{Ann } \mathfrak{b}$.

Lemma 1.9.8. *Ideal quotients of ideals are ideals.*

Proof. Let \mathfrak{a} and \mathfrak{b} be ideals. Then if $x \in (\mathfrak{a} : \mathfrak{b})$, we have $x\mathfrak{b} \subseteq \mathfrak{a}$. Now, let $a \in A$. Then $a(x\mathfrak{b}) = (ax)\mathfrak{b} \subseteq \mathfrak{a}$ so that $ax \in (\mathfrak{a} : \mathfrak{b})$. Notice also that since $x\mathfrak{b} \subseteq \mathfrak{a}$, then $-x\mathfrak{b} \subseteq \mathfrak{a}$. Now, let $x, y \in (\mathfrak{a} : \mathfrak{b})$. Then $x\mathfrak{b} \subseteq \mathfrak{a}$ and $y\mathfrak{b} \subseteq \mathfrak{a}$, thus $x\mathfrak{b} + y\mathfrak{b} = (x + y)\mathfrak{b} \subseteq \mathfrak{a}$ so that $x + y \in (\mathfrak{a} : \mathfrak{b})$. ■

Corollary. $\text{Ann } \mathfrak{b}$ is an ideal. Moreover, the set of zero divisors in the underlying ring is given by

$$D = \bigcup_{x \neq 0} \text{Ann}(x)$$

Example 1.14. Let $m, n \in \mathbb{Z}$, where $m = \prod p^{\mu_p}$ and $n = \prod p^{\nu_p}$. Then $(m\mathbb{Z} : n\mathbb{Z}) = q\mathbb{Z}$ where

$$q = \prod p^{\gamma_p} \text{ and } \gamma_p = \max\{\mu_p - \nu_p, 0\} = \mu_p - \min\{\mu_p, \nu_p\}$$

Lemma 1.9.9. *The following are true for any ideals \mathfrak{a} , \mathfrak{b} , and \mathfrak{c} .*

- (1) $\mathfrak{a} \subseteq (\mathfrak{a} : \mathfrak{b})$.
- (2) $(\mathfrak{a} : \mathfrak{b})\mathfrak{b} \subseteq \mathfrak{a}$.
- (3) $((\mathfrak{a} : \mathfrak{b}) : \mathfrak{c}) = (\mathfrak{a} : \mathfrak{b}\mathfrak{c}) = ((\mathfrak{a} : \mathfrak{c}) : \mathfrak{b})$.
- (4) If $\{\mathfrak{a}_i\}$ is a collection of ideals, then

$$\left(\bigcap \mathfrak{a}_i : \mathfrak{b}\right) = \bigcap (\mathfrak{a}_i : \mathfrak{b})$$

- (5) If $\{\mathfrak{b}_i\}$ is a collection of ideals, then

$$\left(\mathfrak{a} : \sum \mathfrak{b}_i\right) = \bigcap (\mathfrak{a} : \mathfrak{b}_i)$$

Proof. Left as an exercise. ■

Definition. For every ideal \mathfrak{a} of a commutative ring A , with identity, we define the **radical** of \mathfrak{a} to be the set

$$\text{rad } \mathfrak{a} = \{x \in A : x^n \in \mathfrak{a} \text{ for some } n \in \mathbb{Z}^+\}$$

Lemma 1.9.10. *For any ideal \mathfrak{a} , $\text{rad } \mathfrak{a}$ is an ideal of A .*

Proof. Consider the natural map $\phi : A \rightarrow A/\mathfrak{a}$ given by $x \rightarrow x + \mathfrak{a}$. Then notice that

$$\text{rad } \mathfrak{a} = \phi^{-1}(\text{Nil } A/\mathfrak{a})$$

■

Lemma 1.9.11. *The following are true for any ideals \mathfrak{a} and \mathfrak{b} .*

- (1) $\mathfrak{a} \subseteq \text{rad } \mathfrak{a}$.
- (2) $\text{rad}(\text{rad } \mathfrak{a}) = \text{rad } \mathfrak{a}$.
- (3) $\text{rad } \mathfrak{a}\mathfrak{b} = \text{rad}(\mathfrak{a} \cap \mathfrak{b}) = \text{rad } \mathfrak{a} \cap \text{rad } \mathfrak{b}$.
- (4) $\text{rad } \mathfrak{a} = (1)$ if, and only if $\mathfrak{a} = (1)$.
- (5) $\text{rad } \mathfrak{a} + \mathfrak{b} = \text{rad}(\text{rad } \mathfrak{a} + \text{rad } \mathfrak{b})$.
- (6) If \mathfrak{p} is a prime ideal, then $\text{rad } \mathfrak{p}^n = \mathfrak{p}$ for any $n \in \mathbb{Z}^+$.

Lemma 1.9.12. *The radical of an ideal \mathfrak{a} is the intersection of all prime ideals containing \mathfrak{a} .*

Lemma 1.9.13. *The set of zerodivisors of a commutative ring with identity is*

$$D = \bigcup_{x \neq 0} \text{rad}(\text{Ann}(x))$$

Example 1.15. Let $m \in \mathbb{Z}$ and $p_i \in \mathbb{Z}^+$ for $1 \leq i \leq r$ distinct prime divisros of m . Then

$$\text{rad } m\mathbb{Z} = (p_1 \dots p_r)\mathbb{Z} = \bigcap_{i=1}^r p_i\mathbb{Z}$$

Lemma 1.9.14. *Let \mathfrak{a} and \mathfrak{b} be ideals such that $\text{rad } \mathfrak{a}$ and $\text{rad } \mathfrak{b}$ are coprime. Then \mathfrak{a} and \mathfrak{b} are coprime.*

Proof. We have

$$\text{rad}(\mathfrak{a} + \mathfrak{b}) = \text{rad}(\text{rad } \mathfrak{a} + \text{rad } \mathfrak{b}) = \text{rad}(1) = (1)$$

this makes $\mathfrak{a} + \mathfrak{b} = (1)$. ■

1.10 Extensions and Contractions of Ideals

For this section, A and B denote commutative rings with identity.

Definition. Let $\phi : A \rightarrow B$ be a ring homomorphism. We define the **extension** of the ideal \mathfrak{a} of A to be the ideal \mathfrak{a}^e generated by $B\phi(\mathfrak{a})$. That is, $\mathfrak{a}^e = B\phi(\mathfrak{a})$.

Lemma 1.10.1. *Let $f : A \rightarrow B$ a ring homomorphism and \mathfrak{a} an ideal of A . Then*

$$\mathfrak{a}^e = \left\{ \sum y_i f(x_i) : y_i \in B \text{ and } x_i \in \mathfrak{a} \right\}$$

Proof. This follows directly from the definition of \mathfrak{a}^e . ■

Definition. Let $\phi : A \rightarrow B$ a ring homomorphism. We define the **contraction** of the ideal \mathfrak{b} of B to be the preimage $\phi^{-1}(\mathfrak{b})$, and denote it \mathfrak{b}^c ; that is, $\mathfrak{b}^c = \phi^{-1}(\mathfrak{b})$.

Lemma 1.10.2. *Let $\phi : A \rightarrow B$ a ring homomorphism, and \mathfrak{b} an ideal of B . Then \mathfrak{b}^c is an ideal of A . Moreover, if \mathfrak{b} is prime in B , then \mathfrak{b}^c is prime in A .*

Proof. Let $x \in \mathfrak{b}^c$. Then $\phi(x) \in \mathfrak{b}$, so that $-\phi(x) = \phi(-x) \in \mathfrak{b}$, which puts $-x \in \mathfrak{b}^c$; similarly, we get $x + y \in \mathfrak{b}^c$ whenever $x, y \in \mathfrak{b}^c$. Lastly, notice that if $a \in A$, and $x \in \mathfrak{b}^c$, then $\phi(a)\phi(x) = \phi(ax) \in \mathfrak{b}$, so that \mathfrak{b}^c is an ideal.

Now, suppose that \mathfrak{b} is prime. Then since $\mathfrak{b} \neq (1_B)$, $\mathfrak{b}^c \neq (1_A)$. Now, let $ab \in \mathfrak{b}^c$. Then $\phi(ab) = \phi(a)\phi(b) \in \mathfrak{b}$. Since \mathfrak{b} is prime, this puts $\phi(a) \in \mathfrak{b}$ or $\phi(b) \in \mathfrak{b}$; that is, $a \in \mathfrak{b}^c$ or $b \in \mathfrak{b}^c$. Therefore, \mathfrak{b}^c must also be prime. ■

Example 1.16. Let $\phi : A \rightarrow B$ a ring homomorphism. We have that for any prime ideal \mathfrak{b} of B , \mathfrak{b}^c is prime. The same is not true for extensions. If \mathfrak{a} is prime in A , $\mathfrak{a}^e = B\phi(\mathfrak{a})$ need not be prime in B .

Lemma 1.10.3. *Let $\phi : A \rightarrow B$ be a ring homomorphism, with $f = \iota \circ \pi$, where π is onto and ι is 1-1. Then there exists a 1-1 correspondence between the ideals $\phi(A)$ and the ideals of A containing $\ker \phi$. Moreover, prime ideals correspond to prime ideals.*

$$\begin{array}{ccc} A & & \\ \pi \downarrow & \searrow \phi = \iota \circ \pi & \\ \phi(A) & \xrightarrow{\iota} & B \end{array}$$

Example 1.17. Consider the map $\mathbb{Z} \rightarrow \mathbb{Z}[i]$ where $i^2 = -1$. A prime ideal $(p) = p\mathbb{Z}$ may or may not be prime when extended to $\mathbb{Z}[i]$. Now, $\mathbb{Z}[i]$ is a PID, so that we have the following.

- (1) $(2)^e = ((1+i)^2)$ in $\mathbb{Z}[i]$; that is, it is the square of a prime ideal in $\mathbb{Z}[i]$.
- (2) If $p \equiv 1 \pmod{4}$, then $(p)^e$ is the product of two prime ideals in $\mathbb{Z}[i]$, and if $p \equiv 3 \pmod{4}$, $(p)^e$ is a prime ideal in $\mathbb{Z}[i]$.

Lemma 1.10.4. *Let $\phi : A \rightarrow B$ be a ring homomorphism. Then the following are true for ideals \mathfrak{a} and \mathfrak{b} of A and B , respectively.*

- (1) $\mathfrak{a} \subseteq \mathfrak{a}^{ec}$ and $\mathfrak{b}^{ce} \subseteq \mathfrak{b}$.
- (2) $\mathfrak{a}^e = \mathfrak{a}^{ece}$, and $\mathfrak{b}^c = \mathfrak{b}^{cec}$.
- (3) If C is the set of all contracted ideals in A , and E is the set of all extended ideals in B , then

$$C = \{\mathfrak{a} \subseteq A : \mathfrak{a} = \mathfrak{a}^{ec}\} \text{ and } E = \{\mathfrak{b} \subseteq B : \mathfrak{b} = \mathfrak{b}^{ce}\}$$

Moreover, there exists a 1-1 correspondence of C onto E given by the map $\mathfrak{a} \rightarrow \mathfrak{a}^e$.

Proof. First, consider \mathfrak{a} in A . Then $\mathfrak{a}^e = B\phi(\mathfrak{a})$, so that if $x \in \mathfrak{a}$, then $\phi(x) \in f(\mathfrak{a})$, that is $x \in \mathfrak{a}^{ec}$. Similarly, we get $\mathfrak{b}^{ce} \subseteq \mathfrak{b}$.

Now, for the second assertion, we have

$$(\mathfrak{b}^{ce})^c \subseteq \mathfrak{b}^c \subseteq (\mathfrak{b}^c)^{ec}$$

so that $\mathfrak{b}^c = \mathfrak{b}^{cec}$. Similarly, we get $\mathfrak{a}^e = \mathfrak{a}^{ece}$.

Finally, let $\mathfrak{a} \in C$. Then there is a \mathfrak{b} in B for which $\mathfrak{a} = \mathfrak{b}^c$. Then $\mathfrak{a}^e = \mathfrak{b}^{ce} = \mathfrak{b}^{cec} = \mathfrak{a}^{ec}$. Conversely, if $\mathfrak{a} = \mathfrak{a}^{ec}$, then \mathfrak{a} is the contraction of \mathfrak{a}^e . We use a similar argument to prove the result for E . \blacksquare

Lemma 1.10.5. *If $\mathfrak{a}_1, \mathfrak{a}_2$ are ideals of A and $\mathfrak{b}_1, \mathfrak{b}_2$ are ideals of B , then the following are true.*

$$(1) (\mathfrak{a}_1 + \mathfrak{a}_2)^e = \mathfrak{a}_1^e + \mathfrak{a}_2^e \text{ and } \mathfrak{b}_1^c + \mathfrak{b}_2^c \subseteq (\mathfrak{b}_1 + \mathfrak{b}_2)^c.$$

$$(2) (\mathfrak{a}_1 \cap \mathfrak{a}_2)^e \subseteq \mathfrak{a}_1^e \cap \mathfrak{a}_2^e \text{ and } \mathfrak{b}_1^c \cap \mathfrak{b}_2^c = (\mathfrak{b}_1 \cap \mathfrak{b}_2)^c.$$

$$(3) (\mathfrak{a}_1 \mathfrak{a}_2)^e = \mathfrak{a}_1^e \mathfrak{a}_2^e \text{ and } \mathfrak{b}_1^c \mathfrak{b}_2^c \subseteq (\mathfrak{b}_1 \mathfrak{b}_2)^c.$$

$$(4) (\mathfrak{a}_1 : \mathfrak{a}_2)^e \subseteq (\mathfrak{a}_1^e : \mathfrak{a}_2^e) \text{ and } (\mathfrak{b}_1 : \mathfrak{b}_2)^c = (\mathfrak{b}_1^c : \mathfrak{b}_2^c).$$

$$(5) (\text{rad } \mathfrak{a})^e \subseteq \text{rad } \mathfrak{a}^e \text{ and } (\text{rad } \mathfrak{b})^c = \text{rad } \mathfrak{b}^c.$$

1.11 Multivariate Polynomial Rings

Theorem 1.11.1. *Let \mathfrak{a} be an ideal of A and $\mathfrak{a}[x]$ the ideal of $A[x]$ generated by \mathfrak{a} . Then*

$$A[x]/\mathfrak{a}[x] \simeq A/\mathfrak{a}[x]$$

Moreover, if \mathfrak{a} is a prime ideal in A , then $\mathfrak{a}[x]$ is a prime ideal in $A[x]$.

Proof. Consider the map $\pi : A[x] \rightarrow A/\mathfrak{a}[x]$ given by $f \rightarrow f \bmod \mathfrak{a}$. That is, reduce f modulo \mathfrak{a} . Then π is a ring homomorphism with kernel $\ker \pi = \mathfrak{a}[x]$. By the first isomorphism theorem, we get

$$A[x]/\mathfrak{a}[x] \simeq A/\mathfrak{a}[x]$$

Now, let \mathfrak{p} be a prime ideal in A , Then we have that A/\mathfrak{p} is an integral domain, hence, so is $A/\mathfrak{p}[x]$, which makes $\mathfrak{p}[x]$ a prime ideal of $A[x]$. \blacksquare

Example 1.18. Consider the ideal $n\mathbb{Z}$ in \mathbb{Z} . By above, we have

$$\mathbb{Z}[x]/n\mathbb{Z}[x] \simeq \mathbb{Z}/n\mathbb{Z}[x]$$

with natural map reduction of polynomials modulo n . If n is composite, then the ring $\mathbb{Z}/n\mathbb{Z}[x]$ is not an integral domain. If $n = p$ a prime, then $\mathbb{Z}/p\mathbb{Z}[x]$ is an integral domain.

Definition. We define the **polynomial ring** in n variables x_1, \dots, x_n with **coefficients** in A inductively to be

$$A[x_1, \dots, x_n] = A[x_1, \dots, x_{n-1}][x_n]$$

and is the set of all **multivariate polynomials** of the form $f(x_1, \dots, x_n) = \sum ax_1^{d_1} \dots x_n^{d_n}$. We call the monic term $x_1^{d_1} \dots x_n^{d_n}$ of f a **monomial**. We define the **degree** of a monomial to be $\deg x_1^{d_1} \dots x_n^{d_n} = d_1 + \dots + d_n$ and we define the **degree** of f to be $\deg f = \max \{\deg x_1^{d_1} \dots x_n^{d_n}\}$ (i.e. the maximum degree of all monomials of f). If all the monomials of f have the same degree, we call f **homogeneous**, or, a **form**.

Lemma 1.11.2. *Let A be a ring. Then $A[x_1, \dots, x_n]$ is a ring.*

Example 1.19. (1) Consider the polynomial ring $\mathbb{Z}[x, y]$ in two variables x and y with integer coefficients. Then $p(x, y) = 2x^3 + xy - y^2$ and has $\deg p = 3$. The polynomial $q(x, y) = -3xy + 2y^2 + x^2y^3$ has $\deg q = 5$. The sum

$$p + q(x, y) = 2x^3 - 2xy + y^2 + x^2y^3 \text{ has degree } \deg p + q = 5$$

and the product

$$pq(x, y) = -6x^4y + 4x^3y^2 + 2x^5y^3 - 3x^2y^2 + 5xy^3 + x^3y^4 - 2y^4 - x^2y^5$$

had degree $\deg pq = 8$.

(2) The polynomial $p(x, y, z) = 4y^2z^5 - 3xy^3z + 2x^2y$ over $\mathbb{Z}[x, y, z]$ has degree $\deg p = 7$ and the polynomial $q(x, y, z) = 5x^2y^3z^4 - 9x^2z + 7x^2$ has degree $\deg q = 9$. The polynomials

$$p + q(x, y, z) = 5x^2y^3z^4 + 4y^2z^5 - 3xy^3z + 2x^2y - 9x^2z + 7x^2$$

and

$$pq(x, y, z) = 20x^2y^5z^9 - 15x^3y^6z^5 + 10x^4y^4z^4 - 36x^2y^2z^6 + 28x^2y^2z^5 + 27x^3y^3z^2 - 21x^3y^3z - 18x^4yz + 14x^4y$$

have degrees $\deg(p + q) = 9$ and $\deg pq = 16$, respectively.

(3) Consider the polynomials p and q of the above example over $\mathbb{Z}/3\mathbb{Z}$, i.e. as polynomials in $\mathbb{Z}/3\mathbb{Z}[x, y, z]$. Then we have

$$\begin{aligned} p(x, y, z) &= xy^2z^5 + 2x^2y \\ q(x, y, z) &= 2x^2y^3z^4 + x^2 \end{aligned}$$

which makes

$$p + q(x, y, z) = 2x^2y^3z^4 + y^2z^5 + 2x^2y + x^2$$

and

$$pq(x, y, z) = 2x^2y^5z^9 + 1x^4y^4z^4 + 1x^2y^2z^5 + 14x^4y$$

of degrees $\deg(p + q) = 9$ and $\deg pq = 16$, still.

Lemma 1.11.3. *Let A be a commutative ring, and π a permutation of the set $\{1, \dots, n\}$. Then $A[x_1, \dots, x_n] \simeq A[x_{\pi(1)}, \dots, x_{\pi(n)}]$. That is, multivariate polynomial rings are independent of the ordering of their variables.*

Proof. Define the map $\Pi : A[x_1, \dots, x_n] \rightarrow A[x_{\pi(1)}, \dots, x_{\pi(n)}]$ termwise by first sending $x_1 \dots x_n \rightarrow x_{\pi(1)} \dots x_{\pi(n)}$. Then notice that Π defines a ring homomorphism, and moreover, for any $f \in A[x_1, \dots, x_n]$, Π permutes the terms of f . So that Π dictates the required isomorphism. ■

Example 1.20. (1) Consider the ideals (x) and (x, y) in $\mathbb{Q}[x, y]$. We have that (x) is a prime ideal in $\mathbb{Q}[x, y]$, since $\mathbb{Q}[x, y] \simeq \mathbb{Q}[y, x] = \mathbb{Q}[y][x]$. Moreover, let $fg \in (x, y)$ so that $fg(x, y) = xyr(x, y)$ for some $r \in \mathbb{Q}[x, y]$. Then $xy|fg$ which makes $xy|f$ or $xy|g$, so that $f \in (x, y)$ or $g \in (x, y)$. This makes (x, y) a prime ideal. Notice, however, that $(x) \subseteq (x, y)$, so that (x) is not maximal. (x, y) , however is a maximal ideal in $\mathbb{Q}[x, y]$.

(2) Notice that (x, y) is a prime ideal in $\mathbb{Z}[x, y]$, since $\mathbb{Z}[x, y]$ is a subring of $\mathbb{Q}[x, y]$, and (x, y) is prime in $\mathbb{Q}[x, y]$. Similarly, $(2, x, y)$ is prime in $\mathbb{Z}[x, y]$. Notice however that $(x, y) \subseteq (2, x, y)$ so that (x, y) is not maximal in $\mathbb{Z}[x, y]$; $(2, x, y)$ is maximal in $\mathbb{Z}[x, y]$.

(3) Notice that (x, y) is not a principle ideal in $\mathbb{Q}[x, y]$. Suppose that it were, then $(x, y) = (f)$ for some $f(x, y) \in \mathbb{Q}[x, y]$. Then we have that $x \in (f)$ and $y \in (f)$ so that $f|x$ and $f|y$. That is, $x = f(x, y)r(x, y)$ and $y = f(x, y)q(x, y)$. Then $x + y = f(x, y)(r(x, y) + q(x, y))$. Notice also that $\deg f \leq 1$. Then if $\deg f = 0$, f is a unit, and we get $(f) = \mathbb{Q}[x, y]$. On the other hand, if $\deg f = 1$, and since $x + y = f(x, y)(r(x, y) + q(x, y))$, we have that

1.12 Noetherian Rings

Definition. Let A be a commutative ring with identity. We say a sequence of ideals $\{\mathfrak{a}_n\}$ is an **ascending chain** of ideals if $\mathfrak{a}_n \subseteq \mathfrak{a}_{n+1}$ for all $n \in \mathbb{Z}^+$. We say that the chain $\{\mathfrak{a}_n\}$ **stabalizes** if there exists some $k \geq n$, $\mathfrak{a}_k = \mathfrak{a}_n$.

Definition. Let A be a commutative ring with identity. We call A **Noetherian** if every ascending chain of ideals of A stabalizes. We say that A satisfies the **ascending chain condition** on ideals.

Lemma 1.12.1. *If \mathfrak{a} is an ideal of a Noetherian ring A , then the factor ring A/\mathfrak{a} is also Noetherian. In particular, the image of a Noetherian ring under any ring homomorphism is Noetherian.*

Proof. This follows by the isomorphism theorems for ring homomorphisms. ■

Theorem 1.12.2. *The following are equivalent for any ring A .*

- (1) A is Noetherian.
- (2) Every nonempty collection of ideals of A contains a maximal element under inclusion.

(3) Every ideal of A is finitely generated.

Proof. Let A be Noetherian, and let \mathcal{A} be a nonempty collection of ideals of A . Choose an ideal $\mathfrak{a}_1 \in \mathcal{A}$. If \mathfrak{a}_1 is maximal, we are done. If not, then there is an ideal $\mathfrak{a}_2 \in \mathcal{A}$ for which $\mathfrak{a}_1 \subsetneq \mathfrak{a}_2$. Now, if \mathfrak{a}_2 is maximal, we are done. Otherwise, proceeding inductively, if there are no maximal ideals of A in \mathcal{A} , then by the axiom of choice, construct the infinite strictly increasing chain

$$\mathfrak{a}_1 \subsetneq \mathfrak{a}_2 \subsetneq \dots$$

of ideal of A . This contradicts that A is Noetherian, so \mathcal{A} must contain a maximal element.

Now, suppose that any nonempty collection of ideals of A contains a maximal element. Let \mathcal{A} be the collection of all finitely generated ideals of A , and let \mathfrak{a} be any ideal of A . By hypothesis, \mathcal{A} has a maximal element \mathfrak{a}' . Now suppose that $\mathfrak{a} \neq \mathfrak{a}'$, and choose an $x \in \mathfrak{a} \setminus \mathfrak{a}'$, then the ideal generated by \mathfrak{a}' and x is finitely generated, and so is in \mathcal{A} ; but that contradicts the maximality of \mathfrak{a}' . Therefore we must have $\mathfrak{a} = \mathfrak{a}'$.

Finally, suppose every ideal of A is finitely generated, and let $\mathfrak{a} = (a_1, \dots, a_n)$. Let

$$\mathfrak{a}_1 \subsetneq \mathfrak{a}_2 \subsetneq \dots$$

an ascending chain of ideals of A for which

$$\mathfrak{a} = \bigcup_{n \in \mathbb{Z}^+} \mathfrak{a}_n$$

Since $a_i \in \mathfrak{a}$ for each $1 \leq i \leq n$, we have that $a_i \in \mathfrak{a}_{j_i}$ and $i \in \mathbb{Z}^+$. Now, let $m = \max\{j_1, \dots, j_n\}$ and consider the ideal \mathfrak{a}_m . Then $a_i \in \mathfrak{a}_m$ for each i , which makes $\mathfrak{a} \subseteq \mathfrak{a}_m$. That is, $\mathfrak{a}_n = \mathfrak{a}_m$ for some $n \geq m$; which makes A Noetherian. ■

Example 1.21. (1) Every principal ideal domain (PID) is Noetherian, since any collection of ideals has a maximal element. Moreover, lemma 1.7.5 states that PIDs satisfy the ascending chain condition.

(2) The rings \mathbb{Z} , $\mathbb{Z}[i]$, and $k[x]$ (where k is a field) are Noetherian.

(3) The multivariate polynomial ring $\mathbb{Z}[x_1, x_2, \dots]$ is not Noetherian, since the ideal (x_1, x_2, \dots) is not finitely generated.

Theorem 1.12.3 (Hilbert's Basis Theorem). *If A is a Noetherian ring, then so is the polynomial ring $A[x]$.*

Proof. Let \mathfrak{a} be an ideal of $A[x]$, and let L be the set of all leading coefficients of polynomials in \mathfrak{a} . Notice that since $0 \in \mathfrak{a}$, then $0 \in L$, so that L is nonempty. Moreover, let $f(x) = ax^d + \dots$ and $g(x) = bx^e + \dots$ polynomials in \mathfrak{a} of degree $\deg f = d$ and $\deg g = e$, with leading coefficients $a, b \in A$. Then for any $r \in A$, we have the coefficient $ra - b = 0$, or $ra - b$ is the leading coefficient of the polynomial $rx^d f - x^d g \in \mathfrak{a}$. In either case, we get $ra - b \in L$. This makes L an ideal of A . Now, since A is Noetherian L is finitely generated; let $L = (a_1, \dots, a_n)$. Then for every $1 \leq i \leq n$, let $f_i \in \mathfrak{a}$ the polynomial of degree $\deg f_i = e_i$ whose leading coefficient is a_i . Take, then $N = \max\{e_1, \dots, e_n\}$. Then for any $d \in \mathbb{Z}/N\mathbb{Z}$,

let L_d be the set of all leading coefficients of polynomials in \mathfrak{a} , of degree d , together with 0. Let $f_{di} \in \mathfrak{a}$ a polynomial of degree $\deg f_{di} = d$ with leading coefficient b_{di} . We wish to show that

$$\mathfrak{a} = (f_1, \dots, f_n) \cup (f_{d1}, \dots, f_{dn})$$

Let $\mathfrak{a}' = (f_1, \dots, f_n) \cup (f_{d1}, \dots, f_{dn})$. By construction, since the generators were chosen from \mathfrak{a} , $\mathfrak{a}' \subseteq \mathfrak{a}$. Now, if $\mathfrak{a} \neq \mathfrak{a}'$. Then there is a nonzero polynomial $f \in \mathfrak{a}$ of minimum degree not contained in \mathfrak{a}' (i.e. $f \notin \mathfrak{a}'$). Let $\deg f = d$, and let a be the leading coefficient of f . Suppose that $d \geq N$. Since $a \in L$, a is an A -linear combination of the generators of L ; i.e.

$$a = r_1 a_1 + \dots + r_n a_n$$

where $r_1, \dots, r_n \in A$. Let

$$g = r_1 x^{d-e_1} f_1 + \dots + r_n x^{d-e_n} f_n$$

then $g \in \mathfrak{a}'$ and has degree $\deg g = d$ and leading coefficient a . Hence $f - g \in \mathfrak{a}'$ is of smaller degree, and by the minimality of f , $f - g = 0$, which makes $f = g \in \mathfrak{a}'$; a contradiction. Therefore $\mathfrak{a} = \mathfrak{a}'$

Now, if $d < N$, then $a \in L_d$, and so is an A -linear combination of generators of L_d ; that is

$$a = r_1 b_{d1} + \dots + r_n b_{dn}$$

where $r_1, \dots, r_n \in A$. Then let

$$g = r_1 f_{d1} + \dots + r_n f_{dn}$$

then $g \in \mathfrak{a}'$ is a polynomial of degree $\deg g = d$ and leading coefficient a ; which gives us the above contradiction.

Therefore, $\mathfrak{a} = \mathfrak{a}'$, and since \mathfrak{a}' is finitely generated, $A[x]$ is Noetherian. ■

Corollary. *Let k be a field. Then the polynomial ring in n variables $k[x_1, \dots, x_n]$ is Noetherian.*

Definition. Let k be a field. We call a ring A a **k -algebra** if k is contained in the center of A (i.e. $k \subseteq Z(A)$), and $1_k = 1_A$. We call A a **finitely generated k -algebra** if A is generated by k together with a finite set $\{r_1, \dots, r_n\}$ of elements of A .

Definition. Let k be a field and A and S k -algebras. We call a map $\phi : A \rightarrow S$ a **k -algebra homomorphism** if ϕ is a ring homomorphism, and ϕ is the identity on k .

Lemma 1.12.4. *Let k be a field. Then a ring A is a finitely generated k -algebra if, and only if there exists a k -algebra homomorphism $\phi : k[x_1, \dots, x_n] \rightarrow A$ taking $k[x_1, \dots, x_n]$ onto A .*

Proof. If A is generated by elements r_1, \dots, r_n as a k -algebra, then define the map $\phi : k[x_1, \dots, x_n] \rightarrow A$ by taking $x_i \rightarrow r_i$, for all $1 \leq i \leq n$, and $\phi(a) = a$ for all $a \in k$. Then ϕ extends to a ring homomorphism of $k[x_1, \dots, x_n]$ onto A .

Conversly, let ϕ be a k -algebra homomorphism of $k[x_1, \dots, x_n]$ onto A , such that the images $\phi(x_1), \dots, \phi(x_n)$ generate A as a k -algebra. Then A is finitely generated, and since $k[x_1, \dots, x_n]$ is Noetherian by the corollary to Hilbert's basis theorem, A is a quotient of a Noetherian ring, and hence A is Noetherian. This makes A a finitely generated k -algebra. ■

Example 1.22. Let A be a k -algebra, for some field k , viewed as a finite dimensional vector space over k . In particular, let $A = k[x]/(f(x))$, where $f(x)$ is a nonzero polynomial over k . Then A is a finitely generated k -algebra, since it has a finite basis, and that basis serves as a generator for A as a k -algebra. Thus, we have the ideals of A are k -subspaces. Moreover, any ascending chain of ideals of A has at most $\dim_k A - 1$ distinct terms, which means that A satisfies the ascending chain condition.

Bibliography

- [1] D. Dummit, *Abstract algebra*. Hoboken, NJ: John Wiley & Sons, Inc, 2004.
- [2] I. N. Herstein, *Topics in algebra*. New York: Wiley, 1975.