

Contents

1	Functional Analysis	3
1.1	Measure Theory	3
1.2	Integrals	4
1.3	Continuous Linear Operators	7
1.4	Hanh-Banach Theorems	9
1.5	Duality	10
1.6	Banach Spaces	12
1.7	L^p Spaces	14
1.8	Hilbert Spaces	15
1.9	Distributions	18
1.10	Sobolev Spaces of Integer Power	21
2	Harmonic Analysis	29
2.1	Schwartz Space	29
2.2	Fourier Transform	30
2.3	Fourier Operators	31
2.4	Sobolev–Fourier Spaces	33
2.5	Interpolation	36
2.6	Hardy–Littlewood Maximal Function	39
2.7	Calderón–Zygmund Theory	39
2.8	Multiplier Theory	40
2.9	Besov Spaces	41
2.10	Littlewood–Payley Theory	43
2.11	Oscillatory Integrals	43
3	PDE Tools	45
3.1	General Results	45
3.2	Fixed Point Theorems	47
3.3	Variational Formulation	48
3.4	Harmonic Functions	51
4	Differential Equations	51
4.1	Laplace Equation	51
4.2	Heat Equation	52
4.3	Wave Equation	52
4.4	Transport Equation	53
4.5	Disspersive PDEs	54
4.6	Fluid PDEs	55
4.7	Solid Mechanics	55
5	Optimization	57
5.1	General Notions of Optimization	57
5.2	Algorithms	61
6	Probability	61
6.1	Basic Definitions and Results	61
6.2	Limit Theorems	67
6.3	Gaussian and Chi-squared	68
6.4	Matringales	71
6.5	Stochastic Calculus	71

0.1 Definition [Notation]

Let $x, y \in \mathbb{R}^d, \alpha = (\alpha_1, \dots, \alpha_d) \in \mathbb{N}^d$

$$x \cdot y = (x | y)_{\ell^2} = \sum_{j=1}^d x_j y_j, \quad |x| = \|x\|_{\ell^2}$$

$$x^\alpha = x_1^{\alpha_1} \dots x_d^{\alpha_d}, \quad \langle x \rangle = (1 + |x|^2)^{\frac{1}{2}}$$

$$(f | g) = \int f g$$

$$(f(x, y) | g(x, y))_x = \int f(t, y) g(t, y) dt$$

$$(f(x) | g(x))_{L^2} = \int f(x) \bar{g}(x) dx$$

$$\langle T | \varphi \rangle_{X^*, X} = T(\varphi)$$

0.2 Definition [Modified Vinogradov Notation]

Two quantities are related as

$$A \lesssim B$$

if there exists c independent of A, B ; such that $A \leq cB$.

$A \sim B$ means $A \lesssim B$ and $B \lesssim A$

0.3 Definition [Landau Symbols]

Let $f, g : \mathbb{R} \rightarrow \mathbb{R}$ be functions and let $x_0 \in \mathbb{R} \cup \{\pm\infty\}$ with $g(x) \neq 0$ in a punctured neighborhood of x_0 .

1) $f(x) = o(g(x))$ as $x \rightarrow x_0$ if

$$\lim_{x \rightarrow x_0} \frac{f(x)}{g(x)} = 0.$$

2) $f(x) = \mathcal{O}(g(x))$ as $x \rightarrow x_0$ if there exist constants $C > 0$ and $\delta > 0$ such that

$$|f(x)| \leq C |g(x)| \quad \text{for all } x \text{ with } 0 < |x - x_0| < \delta.$$

Alternatively,

$$|f(x)| \lesssim |g(x)|, \quad x \in B(x_0, \delta)$$

1.0 Functional Analysis

1.1 Measure Theory

1.1 Definition [Push-Forward Measure]

Let $(X, \mathcal{A}, \mu)$ be a measure space. Let $(Y, \mathcal{G})$ be a measurable space and

$$T : X \rightarrow Y$$

be a measurable map. Then T induces a measure on $(Y, \mathcal{G})$ from μ .

$$\mu_T(B) := \mu(T^{-1}(B)), \quad \forall B \in \mathcal{G}$$

There is also a notation $\mu_T = T_{\#}\mu$

1.2 Definition [Absolute Continuity]

Let $(X, \mathcal{A})$ be a measurable space and μ, ν be two measures on it. We say that μ is absolutely continuous w.r.t. ν

$$\mu \ll \nu$$

iff

$$\nu(A) = 0 \implies \mu(A) = 0, \quad \forall A \in \mathcal{A}$$

Meaning that null sets of ν must also be null sets of μ .

μ can still have other null sets, thus μ contains less information than ν .

1.3 Theorem [Radon-Nikodym]

Let $(X, \mathcal{A})$ be a measurable space and μ, ν be two measures on it with

$$\mu \ll \nu$$

Then, there exists a measurable $f : X \rightarrow [0, \infty)$ such that

$$\int_A d\mu = \int_A f d\nu$$

Also written

$$d\mu = f d\nu,$$

motivating the terminology f : Radon-Nikodym derivative of μ w.r.t. ν

$$f = \frac{d\mu}{d\nu}$$

In probability, with $\nu = \text{Leb}_{\mathbb{R}}$, such f is usually called pdf (probability density function). Intuitively, we rescale ν with f in some way to get the same measure. The absolute continuity is necessary because we can scale down to zero by f but can't retrieve back the information.

1.4 Theorem [Transfer Theorem]

Let $(X, \mathcal{A}, \mu), (Y, \mathcal{G}, \mu_T), (\mathbb{R}, \mathcal{B}(\mathbb{R}))$ be measure spaces. Let

$$T : (X, \mathcal{A}) \rightarrow (Y, \mathcal{G}), \quad g : (Y, \mathcal{G}) \rightarrow (\mathbb{R}, \mathcal{B}(\mathbb{R}))$$

Then, $g(T) \in L^1(X, \mathcal{A}, \mu)$ iff $g \in L^1(Y, \mathcal{G}, \mu_T)$, and then we have

$$\int_X g(T(x)) d\mu(x) = \int_Y g(y) d\mu_T(y)$$

Usually $Y = \mathbb{R}^d, \mathcal{G} = \mathcal{B}(\mathbb{R}^d)$

(Doesn't require Radon-Nikodym)

1.2 Integrals

Main Integral Theorems

1.5 Theorem [(Fubini-)Tonelli]

Let $(X, \mathcal{A}, \mu)$ and $(Y, \mathcal{B}, \nu)$ be σ -finite measure spaces. Let $f : X \times Y \rightarrow [0, \infty]$ be $\mu \otimes \nu$ measurable.

Then,

$$\int_{X \times Y} f d(\mu \otimes \nu) = \int_X \left[\int_Y f(x, y) d\nu(y) \right] d\mu(x) = \int_Y \left[\int_X f(x, y) d\mu(x) \right] d\nu(y).$$

1.6 Theorem [Fubini]

Let $(X, \mathcal{A}, \mu)$ and $(Y, \mathcal{B}, \nu)$ be σ -finite measure spaces, and let $f : X \times Y \rightarrow \mathbb{R}$ be $L^1(X \times Y, \mathcal{A} \otimes \mathcal{B}, \mu \otimes \nu)$, i.e.

$$\int_{X \times Y} |f(x, y)| \, d(\mu \otimes \nu)(x, y) < \infty.$$

Then,

$$\int_{X \times Y} f \, d(\mu \otimes \nu) = \int_X \left[\int_Y f(x, y) \, d\nu(y) \right] d\mu(x) = \int_Y \left[\int_X f(x, y) \, d\mu(x) \right] d\nu(y),$$

and the iterated integrals exist and are finite.

1.7 Theorem [Dominated Convergence]

Let $(f_n)_{n \in \mathbb{N}}$ be a sequence of measurable functions on $(X, \mathcal{A}, \mu)$, such that

i) $\exists f : X \rightarrow \mathbb{R}$ μ -measurable with

$$f_n \xrightarrow[n \rightarrow \infty]{a.e.} f$$

ii) $\exists g \in L^1(X, \mathcal{A}, \mu)$ with

$$g \geq 0, \quad |f_n| \leq g, \forall n$$

Then, $f_n, f \in L^1(X, \mathcal{A}, \mu)$ with

$$\int_X |f_n - f| \, d\mu \xrightarrow[n \rightarrow \infty]{} 0$$

which implies

$$\lim_{n \rightarrow \infty} \int_X f_n \, d\mu = \int_X \lim_{n \rightarrow \infty} f_n \, d\mu = \int_X f \, d\mu$$

1.8 Theorem [Monotone Convergence]

Let $(f_n)_{n \in \mathbb{N}}$ be a sequence of measurable functions on $(X, \mathcal{A}, \mu)$, such that

i) $f_n(x) \nearrow f(x)$ almost everywhere on X .

Then, $f : X \rightarrow [0, \infty]$ is μ -measurable and

$$\lim_{n \rightarrow \infty} \int_X f_n \, d\mu = \int_X \lim_{n \rightarrow \infty} f_n \, d\mu = \int_X f \, d\mu.$$

1.9 Theorem [Fatou's Lemma]

Let $(f_n)_{n \in \mathbb{N}}$ be a sequence of measurable functions on $(X, \mathcal{A}, \mu)$.

Define $f(x) := \liminf_{n \rightarrow \infty} f_n(x)$, then $f : X \rightarrow [0, \infty]$ is measurable. **Then,**

$$\int_X f \, d\mu = \int_X \liminf_{n \rightarrow \infty} f_n \, d\mu \leq \liminf_{n \rightarrow \infty} \int_X f_n \, d\mu.$$

Integrals depending on a parameter

1.10 Theorem [Continuity Under the Integral Sign]

Let (T, d) be a metric space (of parameters), $t^* \in T$, and $f : T \times X \rightarrow \mathbb{C}$ with

- i) for every $t \in T$ fixed, $x \mapsto f(t, x) \in L^1(X, \mathcal{A}, \mu)$
- ii) for a.e. $x \in X$ fixed, $t \mapsto f(t, x)$ is continuous at t^*
- iii) $\exists g : X \rightarrow \mathbb{R}_+$ in $L^1(X, \mathcal{A}, \mu)$, such that for every $t \in T$

$$|f(t, x)| \leq g(x), \quad \text{for a.e. } x$$

Then,

$$F(t) := \int_X f(t, x) \, d\mu(x)$$

is continuous at t^*

1.11 Theorem [Differentiability Under the Integral Sign]

Let $I \subset \mathbb{R}$ be an interval, and $f : I \times X \rightarrow \mathbb{C}$ with

- i) for every $t \in I$ fixed, $x \mapsto f(t, x) \in L^1(X, \mathcal{A}, \mu)$

- ii) for a.e. $x \in X$ fixed, $t \mapsto f(t, x)$ is differentiable on I
- iii) $\exists g : X \rightarrow \mathbb{R}_+$ in $L^1(X, \mathcal{A}, \mu)$, such that for every $t \in T$

$$|\partial_t f(t, x)| \leq g(x), \quad \text{for a.e. } x$$

Then,

$$F(t) := \int_X f(t, x) \, d\mu(x)$$

is differentiable on I , and for all $t \in I$, its derivative is

$$\partial_t F(t) = \int_X \partial_t f(t, x) \, d\mu(x)$$

1.3 Continuous Linear Operators

1.12 Theorem [Vector Spaces]

Let $(E, \| \cdot \|, \mathbb{K})$ be a finite dimensional vector space.

- all norms on E are equivalent
- $(E, \| \cdot \|)$ is Banach
- closed and bounded is equivalent to compact

Let E be a vector space.

- any finite dimensional subset of E is closed
- $\text{span}(x_1, x_2, \dots)$ is closed if it spans a finite dimensional subspace, otherwise, it might be open

1.13 Theorem [Characterization of Continuous Linear Maps]

Let E, F be normed, $T : E \rightarrow F$ be a linear map. The following are equivalent

- 1) T is continuous at 0
- 2) T is continuous on E

3) T is uniformly continuous on E

4) T is Lipschitz on E

5) T is bounded in the following sense

$$(\exists M > 0 : \forall x \in E) : \|T(x)\|_F \leq M \|x\|_E$$

The space of all continuous linear operators between E, F will be denoted $\mathcal{L}(E, F)$

1.14 Definition

$\mathcal{L}(E, F)$ can be equipped with

$$\|T\|_{\mathcal{L}(E, F)} = \sup_{\|x\|_E=1} \|T(x)\|_F$$

The operator norm will be denoted either by $\|\cdot\|_{\mathcal{L}(E, F)}$ or $\|\cdot\|_{E \rightarrow F}$

1.15 Theorem

Let F Banach, then $(\mathcal{L}(E, F), \|\cdot\|_{\mathcal{L}(E, F)})$ is Banach.

1.16 Proposition

Let E be a finite dimensional normed vector space, F normed. **Then**, any linear map $T : E \rightarrow F$ is continuous.

1.17 Proposition

Let $A \in \mathcal{L}(D, E), B \in \mathcal{L}(E, F)$, then, $B \circ A \in \mathcal{L}(D, F)$ with

$$\|B \circ A\|_{\mathcal{L}(D, F)} \leq \|A\|_{\mathcal{L}(D, E)} \|B\|_{\mathcal{L}(E, F)}$$

1.18 Theorem [Continuous Linear Extension]

Let E be normed, F Banach, D dense in E . Let $T_0 \in \mathcal{L}(D, F)$. **Then**, there exists

the unique extension $T \in \mathcal{L}(E, F)$ with

$$T|_D = T_0, \quad \|T\|_{\mathcal{L}(E, F)} = \|T_0\|_{\mathcal{L}(D, F)}$$

1.4 Hahn-Banach Theorems

1.19 Theorem [Hahn-Banach Real Analytic Form]

Let E be a vector space over $\mathbb{R}$. Let $G \leq E$ be a **linear subspace**, $p : E \rightarrow \mathbb{R}$ be a **sublinear function**

$$\begin{aligned} p(\lambda x) &= \lambda p(x), & \forall x \in E, \forall \lambda > 0 \\ p(x + y) &\leq p(x) + p(y), & \forall x, y \in E \end{aligned}$$

Let $\ell_0 : G \rightarrow \mathbb{R}$ be a **linear functional** with

$$\ell_0(x) \leq p(x), \forall x \in G$$

Then, there exists a **linear functional** $\ell : E \rightarrow \mathbb{R}$ with

$$\ell|_G = \ell_0, \quad \ell \leq p$$

Proved by Zorn's lemma, non-constructive.

1.20 Theorem [Hahn-Banach for Normed Spaces]

Let $(E, \|\cdot\|)$ be a normed vector space over $\mathbb{K}$. Let $G \leq E$ be a **linear subspace**, $\ell_0 \in G^*$. **Then**, there exists a **linear functional** $\ell \in (E^*, \|\cdot\|_{E^*})$ with

$$\ell|_G = \ell_0, \quad \|\ell\|_{E^*} = \|\ell_0\|_{G^*}$$

1.21 Theorem [Hahn-Banach for Vanishing Forms]

Let $(E, \|\cdot\|)$ be a normed vector space over $\mathbb{K}$. Let $G < E$ be a **proper linear**

subspace. Then, there exists a linear functional $\ell \in (E^, \|\cdot\|_{E^*})$ with*

$$\ell|_G \equiv 0, \quad \ell \not\equiv 0$$

1.22 Theorem [Hahn-Banach Geometric Form]

*Let $A, B \subset E$ be two disjoint, non-empty, subsets. Let one of them be open. Then, there exists a **closed hyperplane** that separates A and B .*

Where a hyperplane is an affine subspace of codimension 1

$$H = [\ell = \alpha] = \{x \in E ; \ell(x) = \alpha\}$$

is closed iff $\ell : E \rightarrow \mathbb{R}$ is continuous.

1.23 Theorem [Hahn-Banach 2nd Geometric Form]

Let $A \subset E, B \subset E$ be two disjoint, non-empty, subsets. Let A be closed and B compact. Then, there exists a closed hyperplane that strictly separates A and B .

1.5 Duality

1.24 Definition [Linear Forms, Dual Space]

Let $(E, \mathbb{K}, \|\cdot\|)$ be a normed space, then $\mathcal{L}(E, \mathbb{K}) = E^*$, a space of continuous linear forms, is called a dual space.

1.25 Definition [Weak and Weak* Topologies]

Let E be a normed space and E^* its dual.

The **weak topology** $\sigma(E, E^*)$ on E is the coarsest topology such that all (continuous in strong norm topology) functionals $\ell \in E^*$ remain continuous.

We say that $(x_n) \subset E$ converges weakly to $x \in E$

$$x_n \xrightarrow{E} x$$

iff

$$\langle \ell | x_n \rangle_{E^*, E} \xrightarrow[n \rightarrow \infty]{} \langle \ell | x \rangle_{E^*, E} \quad (\forall \ell \in E^*)$$

The **weak* topology** $\sigma(E^*, E)$ on E^* is the coarsest topology such that all evaluation maps

$$x \mapsto \ell(x) = \langle \ell | x \rangle, \quad x \in E,$$

remain continuous.

We say that $(\ell_n) \subset E^*$ converges weak* to $\ell \in E^*$

$$\ell_n \xrightarrow{E^*} \ell$$

iff

$$\langle \ell_n | x \rangle_{E^*, E} \xrightarrow[n \rightarrow \infty]{} \langle \ell | x \rangle_{E^*, E} \quad (\forall x \in E)$$

1.26 Theorem [Dual of Banach]

Let E be Banach, then E^* is Banach.

1.27 Definition [Reflexivity]

Let E be normed and E^* be its dual. Let $E^{**} = (E^*)^*$, and denote the canonical embedding

$$J : E \rightarrow E^{**}, \quad J_\ell(x) = \langle \ell | x \rangle_{E^*, E}$$

J is a (linear) isometric injection.

If J is also a surjection, then E is called a reflexive space.

1.28 Definition [Transpose]

Let E be a normed space, $L \in \mathcal{L}(E, E)$. Then, we can define the transpose $L^\top \in \mathcal{L}(E^*, E^*)$ by the relation

$$\langle L^\top \ell | x \rangle_{E^*, E} := \langle \ell | Lx \rangle_{E^*, E}$$

If $L = L^\top$, an operator is called self-transpose.

1.29 Theorem [Fencel–Moreau]

Assume that $\varphi : E \rightarrow (-\infty, \infty]$ is convex, l.s.c., and $\varphi \not\equiv \infty$. **Then**, $\varphi^{**} = \varphi$.
Where $\varphi^* : E^* \rightarrow (-\infty, \infty]$, a Legendre–Fenchel transform of φ , is given by

$$\varphi^*(f) = \sup_{x \in E} \{\langle f | x \rangle - \varphi(x)\}, \quad f \in E^*$$

1.6 Banach Spaces

1.30 Theorem [Baire]

Let X be a complete metric space and let $(X_n)_{n \in \mathbb{N}}$ be a sequence of closed subsets in X with

$$\text{Int}(X_n) = \emptyset, \quad \forall n \in \mathbb{N}$$

Then,

$$\text{Int} \left(\bigcup_{n \in \mathbb{N}} X_n \right) = \emptyset$$

1.31 Theorem [Characterization via Absolute Convergence]

Let $(X, \| \cdot \|)$ be a normed vector space. The following are equivalent:

- i) X is a Banach
- ii) Every absolutely convergent series in X is convergent, i.e.,

$$\sum_{n=1}^{\infty} \|x_n\| < \infty \implies \sum_{n=1}^{\infty} x_n \text{ converges in } X.$$

1.32 Theorem [Banach–Steinhaus Uniform Boundedness Principle]

Let E, F be Banach and let $(T_i)_{i \in I}$ be a family in $\mathcal{L}(E, F)$ with

$$\sup_{i \in I} \|T_i x\|_F < \infty, \quad \forall x \in E.$$

Then,

$$\sup_{i \in I} \|T_i\|_{\mathcal{L}(E, F)} < \infty$$

Alternatively, there exists c

$$\|T_i x\| \leq c \|x\|, \quad \forall x \in E, \forall i \in I.$$

1.33 Theorem [Open Mapping]

Let E, F be Banach and let $T \in \mathcal{L}(E, F)$ surjective. **Then**, there exists $c > 0$, such that

$$T(B_E(0, 1)) \supset B_F(0, c).$$

In other words, T is an open mapping.

1.34 Theorem [Closed Graph]

Let E, F be Banach and let T be linear. If the graph $\Gamma(T)$ is closed in $E \times F$, **then**, T is continuous.

1.35 Theorem [Kakutani]

Let E be Banach. E is reflexive **iff** $\overline{B_E}$ is compact in the weak topology $\sigma(E, E^*)$

1.36 Theorem [Riesz]

Let $(E, \|\cdot\|)$ be normed. $\overline{B_E}$ is compact **iff** E is finite dimensional.

1.37 Theorem [Ascoli–Arzelà]

Let K be a compact metric space and let $\mathcal{H}$ be a bounded subset of $C(K; K)$. If $\mathcal{H}$ is uniformly equicontinuous, that is,

$$\left(\forall f \in \mathcal{H} : \forall \varepsilon > 0 : \exists \delta > 0 \right) \quad d(x, y) < \delta \implies |f(x) - f(y)| < \varepsilon$$

Then, the closure of $\mathcal{H}$ in $C(K; K)$ is compact

1.38 Theorem [Banach–Alaoglu]

Let E be normed. Then, the closed unit ball $\overline{B_{E^*}}$ is compact in E^* , w.r.t. the weak* topology $\sigma(E^*, E)$

1.39 Theorem [Banach–Picard Contractive Mapping Theorem]

Contractive functions on complete metric spaces uniquely fix a point.

Let (X, d) be a non-empty complete metric space, $K < 1$,

$f : X \rightarrow X$ be a continuous mapping with property

$$d(f(x), f(y)) \leq Kd(x, y), \quad \forall x, y \in X.$$

Then, there exists a unique x_* , such that $f(x_*) = x_*$

Such f is called a **strict contraction** and x_* is called a **fixed-point**.

1.7 L^p Spaces

1.40 Theorem [Fischer–Riesz]

L^p spaces are Banach for $p \in [1, \infty]$

1.41 Theorem [Kolmogorov–M. Riesz–Fréchet]

Let $\mathcal{F} \subset L^p(\mathbb{R}^d)$ be bounded, with $p \in [1, \infty)$. If

$$\lim_{|h| \rightarrow 0} \|\tau_h f - f\|_{L^p(\mathbb{R}^d)} = 0, \quad \text{uniformly in } f \in \mathcal{F}$$

then, the closure of restriction $\overline{\mathcal{F}|_{\Omega}}^{\|\cdot\|_{L^p(\Omega)}}$ is compact for any measurable $\Omega \subset \mathbb{R}^d$

1.42 Proposition [Reflexivity and Separability of L^p spaces]

For $p \in (1, \infty)$, the space $L^p(\Omega)$ is reflexive and separable. For L^1 is separable but not reflexive. L^∞ is neither separable nor reflexive.

1.43 Theorem [Duality of L^p]

1.44 Definition [Convolution]

1.45 Definition [Good Kernels]

A sequence $(\delta_n)_n \subset L^1(\mathbb{R}^d)$ is called a **family of good kernels** or an **approximation of identity** iff

1.8 Hilbert Spaces

Geometry, Orthogonality, Projections

1.46 Theorem [Cauchy–Schwarz Inequality]

Let $(H, (\cdot | \cdot))$ be an inner product space. Then, $\forall x, y \in H$

$$| (x | y) | \leq \|x\| \|y\| = \sqrt{(x | x)} \sqrt{(y | y)}$$

1.47 Definition [Orthogonality]

Let $(H, (\cdot | \cdot))$ be an inner product space.

- $x, y \in H$ are called **orthogonal** iff $(x | y) = 0$
- $A \subset H$, $A^\perp = \{y \in H : \forall x \in A ; (x | y) = 0\}$

1.48 Theorem [Parallelogram Identity]

Let $(H, (\cdot | \cdot))$ be an inner product space. Then, $\forall x, y \in H$,

$$\|x + y\|^2 + \|x - y\|^2 = 2\|x\|^2 + 2\|y\|^2.$$

1.49 Theorem [Polarization Identity]

Let $(H, (\cdot | \cdot))$ be an inner product space over $\mathbb{R}$ or $\mathbb{C}$. For all $x, y \in H$, the inner

product can be recovered from the norm by

$$(x | y) = \begin{cases} \frac{\|x + y\|^2 - \|x - y\|^2}{4}, & \text{if } H \text{ is real} \\ \frac{\|x + y\|^2 - \|x - y\|^2 + i\|x + iy\|^2 - i\|x - iy\|^2}{4}, & \text{if } H \text{ is complex} \end{cases}$$

1.50 Theorem [Pythagoras Identity]

Let $(H, (|))$ be an inner product space. Then for all orthogonal systems $x_1, \dots, x_n$

$$\|x_1 + \dots + x_n\|^2 = \|x_1\|^2 + \dots + \|x_n\|^2$$

1.51 Theorem [Fréchet–von Neumann–Jordan]

Let $(E, \| \cdot \|)$ be normed. This norm is Hilbert norm, i.e. rising from some inner product, *iff* it satisfies the parallelogram identity.

Moreover, in this case, it is explicitly derived by polarization identity.

1.52 Theorem [Closed Subspaces, Characterization]

1.53 Proposition

Let $K \leq H$, then K is dense in H iff $K^\perp = \{0\}$

Hilbert Space Theorems

1.54 Theorem [Riesz—Fréchet Representation]

Let H Hilbert. Given any $\varphi \in H^*$, there exists a unique $f_\varphi \in H$, such that

$$\langle \varphi | u \rangle_{H^*, H} = \varphi(u) = (f_\varphi | u)_H, \quad \forall u \in H$$

Moreover,

$$\|f_\varphi\|_H = \|\varphi\|_{H^*}$$

1.55 Theorem [Stampacchia]

Let $a(u, v)$ be a continuous coercive bilinear form on H Hilbert. Let $K \subset H$ be a non-empty closed and convex subset. **Then**, given any $\varphi \in H^*$, there exists a unique $u \in K$, such that

$$\varphi(v - u) \leq a(u, v - u), \quad \forall v \in K$$

1.56 Theorem [Lax–Milgram]

Let $a(u, v)$ be a *continuous coercive (elliptic) bilinear form* on H Hilbert.

$$|a(u, v)| \leq C \|u\|_H \|v\|_H, \quad a(v, v) \geq \alpha \|v\|_H^2$$

Then, given any $\varphi \in H^*$, there exists a unique $u \in H$, such that

$$a(u, v) = \ell(v), \quad \forall v \in H.$$

and it holds that

$$\|u\|_H \leq \frac{1}{\alpha} \|\ell\|_{H^*}$$

Moreover, if a is *symmetric*, then u is characterized by

$$u = \arg \min_{\ell \in H} \left\{ \frac{1}{2} a(v, v) - \ell(v) \right\}$$

1.57 Theorem [Lax–Milgram Operator Version]

Equivalent form for $a(u, v) = (Au | v)_H$ on H Hilbert.

Let $A : H \rightarrow H$ be a linear operator with

$$|(Au | v)_H| \leq C \|u\|_H \|v\|_H, \quad (Av | v)_H \geq \alpha \|v\|_H^2$$

Then, A is a bijection with a bounded inverse A^{-1} , satisfying

$$\|A^{-1}\| \leq \frac{1}{\alpha}$$

Self-Adjoint Operators

1.58 Definition [Adjoint]

Let H Hilbert, $A \in \mathcal{L}(H)$, its adjoint A^* is defined via

$$(A^*u | v)_H = (u | Av)_H$$

If $A = A^*$, an operator is called self-adjoint.

Hilbert Bases

1.9 Distributions

1.59 Definition [Bump Functions C_c^∞]

Let $\Omega \subseteq \mathbb{R}^d$ be open. The space of smooth functions $f : \Omega \rightarrow \mathbb{C}$ with compact support (bump functions / test functions) is denoted by $\mathcal{D}(\Omega) = C_c^\infty(\Omega; \mathbb{C})$ or $C_0^\infty(\Omega; \mathbb{C})$. One can also choose $C_c^\infty(\Omega; \mathbb{R})$.

1.60 Proposition [Density of Bump Functions]

1.61 Definition [Distributions]

Let $\Omega \subseteq \mathbb{R}^d$ be open. The space of **distributions** is the topological dual of the space of test functions:

$$\mathcal{D}^*(\Omega) = (\mathcal{D}(\Omega))^*$$

where $\mathcal{D}^*(\Omega)$ consists of all continuous linear functionals (distributions)

$$T : \mathcal{D}(\Omega) \rightarrow \mathbb{C}$$

Where continuity is with respect to the strong LF topology, from which we can derive the weak^{*} topology $\sigma(\mathcal{D}^*, \mathcal{D})$.

Distributions, also called **generalized functions**, will be then defined not pointwise, like functions, but w.r.t. to averages, given by the integrals against test functions. If we think of f as a distribution T_f ,

1.62 Example

Let $f \in L^1_{\text{loc}}(\Omega)$. Then f defines a distribution T_f as

$$T_f(\varphi) = \langle T_f | \varphi \rangle := (T_f | \varphi) = \int_{\Omega} f(x) \varphi(x) \, dx$$

1.63 Example

Let μ be a (signed) Borel measure on Ω , which is finite on compact subsets of Ω (Radon measure). Then

$$T_{\mu}(\varphi) = \langle T_{\mu} | \varphi \rangle_{D^*, D} = \int_{\Omega} \varphi(x) \, d\mu(x)$$

is a distribution. A concrete example of it is if we take $\Omega = \mathbb{R}$, $\mu = \delta_0 = \#_{\{0\}}$, which gives us the **Dirac Delta** (Dirac Mass)

$$\delta_0(\varphi) = \langle \delta_0 | \varphi \rangle_{D^*, D} = \int_{\mathbb{R}} \varphi(x) \, d\delta_0(x) = \varphi(0)$$

Observe that if $f \in C^\infty(\Omega)$, $\varphi \in \mathcal{D}(\Omega)$, we can always use integration by parts

$$\int (\partial_x^\alpha f) \varphi \, dx = (-1)^{|\alpha|} \int f (\partial_x^\alpha \varphi) \, dx,$$

which motivates the following definition.

1.64 Definition [Weak Derivative]

Let $f \in L^1_{\text{loc}}(\Omega)$, if there exists $g \in L^1_{\text{loc}}(\Omega)$, such that

$$\int f (\partial_x^\alpha \varphi) \, dx = (-1)^{|\alpha|} \int g \varphi \, dx, \quad \forall \varphi \in \mathcal{D}(\Omega).$$

Then, g is called a weak derivative of f of order α . Can also be defined for L^2 functions.

1.65 Definition [Distributional Derivative]

Let T be a distribution, then its **distributional derivative** of order α is defined via transpose

$$\langle \partial_x^\alpha T \mid \varphi \rangle := (-1)^{|\alpha|} \langle T \mid \partial_x^\alpha \varphi \rangle$$

1.66 Remark

Distributional, weak, classical derivatives are related as following. A weak derivative $\partial^\alpha f = g$ coincides with the classical derivative (if exists). The corresponding distributions T_f, T_g are thus related by the distributional derivative $\partial^\alpha T_f = T_g$

1.67 Example

Let $H : \mathbb{R} \rightarrow \mathbb{R}; x \mapsto \mathbf{1}_{\mathbb{R}_+}$. It has a distributional derivative Dirac δ_0 but has neither classical nor weak derivative.

1.68 Definition [Operations on Distributions]

Let $F \in \mathcal{D}^*, \varphi \in \mathcal{D}, \psi \in C^\infty$. Let $\tau_h f(x) = f(x - h)$ be a translation operator, let $f_a(x) = f(ax)$ be dilation and $f_L(x) = f(L(x))$ be a non-singular linear operator. Then, we define operators on S^* via transposes,

$$\begin{aligned} \langle \psi \cdot F \mid \varphi \rangle &:= \langle F \mid \psi \cdot \varphi \rangle \\ \langle \psi \star F \mid x \mapsto \varphi(-x) \rangle &:= \langle F \mid \psi \star \varphi \rangle \\ \langle \tau_h F \mid \varphi \rangle &:= \langle F \mid \tau_{-h} \varphi \rangle \\ \langle F_a \mid \varphi \rangle &:= a^{-d} \langle F \mid \varphi_{a^{-1}} \rangle \\ \langle F_L \mid \varphi \rangle &:= |\det L|^{-1} \langle F \mid \varphi_{L^{-1}} \rangle \end{aligned}$$

1.69 Definition [Convolution]

Let $\varphi \in \mathcal{D}(\mathbb{R}^d)$ and $T, S \in \mathcal{D}^*(\mathbb{R}^d)$

1.70 Definition [Convergence of Test Functions]

$(\varphi_n) \subset \mathcal{D}$ weakly converges to some $\varphi \in \mathcal{D}$

$$\varphi_n \xrightarrow[n \rightarrow \infty]{\sigma(\mathcal{D}, \mathcal{D}^*)} \varphi \qquad \varphi_n \xrightarrow{\mathcal{D}} \varphi$$

iff

$$\langle \Phi | \varphi_n \rangle \xrightarrow[n \rightarrow \infty]{} \langle \Phi | \varphi \rangle, \quad (\forall \Phi \in \mathcal{D}^*)$$

This is a very strong notion of convergence and we rarely care about the sequences of test functions and its strong and weak limits, so it is **never used**.

1.71 Definition [Convergence of Distributions]

$(T_n) \subset \mathcal{D}^*$ weakly^{*} converges to some $T \in \mathcal{D}^*$

$$T_n \xrightarrow[n \rightarrow \infty]{\sigma(\mathcal{D}^*, \mathcal{D})} T \quad T_n \xrightarrow{\mathcal{D}^*} T$$

iff

$$\langle T_n | \varphi \rangle \xrightarrow[n \rightarrow \infty]{} \langle T | \varphi \rangle, \quad (\forall \varphi \in \mathcal{D})$$

This is the weakest notion of convergence and is used very frequently.

Convergence in the sense of distributions is what we call when we take a sequence, e.g. $(u_n) \subset L^2$, and we look at the copy of this sequence in $\mathcal{D}^*$. As each u_n defines a distribution $T_{u_n} \in \mathcal{D}^*$, we can look at the weak^{*} limit of this sequence

$$T_{u_n} \xrightarrow{\mathcal{D}^*} T,$$

and then we can say (u_n) converges to $T \in \mathcal{D}^*$, in the sense of distributions. A posteriori, the limiting distribution T might also contain a copy in a more structured space, e.g. $u \in L^2$.

1.72 Proposition

We have the following chain of inclusions

$$D \subsetneq L^p \subsetneq L_{\text{loc}}^p \subsetneq L_{\text{loc}}^1 \subsetneq \mathcal{D}^*$$

1.10 Sobolev Spaces of Integer Power

1.73 Definition [Sobolev Spaces $W^{m,p}$]

Let $\Omega \subset \mathbb{R}^d$ be open. Let $m \in \mathbb{N}, p \in [1, \infty]$. Then, we define $W^{m,p}(\Omega)$ to be the of of

$L^p(\Omega)$ functions, whose distributional derivatives up to order m are in L^p

$$W^{m,p}(\Omega) = \{u \in L^p; \forall |\alpha| \leq m, \partial^\alpha u \in L^p\}$$

With the norm

$$\|u\|_{W^{m,p}} = \sum_{|\alpha| \leq m} \|\partial^\alpha u\|_{L^p}$$

and seminorm

$$[u]_{W^{m,p}} = \sup_{|\alpha|=m} \|\partial^\alpha u\|_{L^p}$$

Functional Inequalities, Embeddings

1.74 Theorem [Hölder Inequality]

Let $p, q, r \in [1, \infty]$, $f \in L^p(\Omega)$, $g \in L^q(\Omega)$ and

$$\frac{1}{p} + \frac{1}{q} = \frac{1}{r}$$

Then,

$$\|fg\|_{L^r} \leq \|f\|_{L^p} \|g\|_{L^q},$$

which implies that $fg \in L^r(\Omega)$.

(p, q) is called Hölder pair, p and q are each other's Hölder conjugates.

1.75 Theorem [Interpolation Inequality]

Let $p_0 < p < p_1$ with

$$\frac{1}{p} = \frac{\theta}{p_0} + \frac{1-\theta}{p_1}$$

Then,

$$\|f\|_p \leq \|f\|_{p_0}^\theta \|f\|_{p_1}^{1-\theta}$$

1.76 Proposition [Density of Test Functions]

Let Ω be Lipschitz, then, if we take the set of test functions, restricted to Ω and then

take its closure w.r.t. the norm of the respective Sobolev space, we will get

$$\overline{\mathcal{D}_{|\Omega}(\mathbb{R}^d)}^{\|\cdot\|_{W^{m,p}}} = W^{m,p}(\Omega).$$

1.77 Theorem [Poincaré Inequality]

Let Ω be Lipschitz and bounded in at least one direction, $p \in [1, \infty]$, $u \in W_0^{1,p}$. Then,

$$\|u\|_{L^p} \lesssim \|\nabla u\|_{L^p}$$

One consequence of that is it makes $\|\nabla u\|_{L^p}$ equivalent to the Sobolev norm on spaces, on which the Poincaré inequality holds.

1.78 Theorem [Gagliardo–Nirenberg–Sobolev Embeddings]

Let $\Omega \subset \mathbb{R}^d$ and let m, d, p, p^* satisfy

$$\frac{1}{p^*} := \frac{1}{p} - \frac{m}{d} > 0.$$

Then, we have a continuous embedding

$$W^{m,p}(\Omega) \hookrightarrow L^{p^*}$$

with

$$\|u\|_{L^{p^*}(\mathbb{R}^d)} \lesssim [u]_{W^{m,p}(\mathbb{R}^d)}$$

and generally

$$\|u\|_{L^{p^*}(\Omega)} \lesssim [u]_{W^{m,p}(\Omega)} + \|u\|_{L^p(\Omega)}$$

1.79 Theorem [Morrey]

Let $u \in W^{1,p}(\mathbb{R}^d)$ with $p > d$, then

$$W^{1,p}(\mathbb{R}^d) \hookrightarrow C^{0,\gamma}(\mathbb{R}^d), \quad \|u\|_{C^{0,\gamma}(\mathbb{R}^d)} \lesssim \|u\|_{W^{1,p}(\mathbb{R}^d)}.$$

with $\gamma = 1 - \frac{d}{p}$

1.80 Theorem [Rellich–Kondrashov]

Let Ω be bounded, Lipschitz, let

$$\frac{1}{p} - \frac{m}{d} < \frac{1}{q} \leq \frac{1}{p}$$

Then,

$$W^{m,p}(\Omega) \Subset L^q(\Omega)$$

In particular, $W^{m,p} \Subset L^p$

1.81 Theorem [Gagliardo–Nirenberg]

Let $j, m \in \mathbb{N}$ and $p, q, r \in [1, \infty]$ with $j < m$. If

$$\exists \theta \in [j/m, 1) : \frac{1}{p} - \frac{j}{d} = \theta \left(\frac{1}{r} - \frac{m}{d} \right) + \frac{1-\theta}{q}$$

Then,

$$[u]_{W^{j,p}(\mathbb{R}^d)} \lesssim [u]_{W^{m,r}(\mathbb{R}^d)}^\theta \|u\|_{L^q(\mathbb{R}^d)}^{1-\theta}$$

1.82 Proposition [Product Rule]

Let $u, v \in W^{m,p}(\mathbb{R}^d)$ with $mp > d$. Then,

$$\|uv\|_{L^\infty} \lesssim \|u\|_{W^{1,\infty}} \|v\|_{W^{m,p}} + \|u\|_{W^{m,p}} \|v\|_{W^{1,\infty}}$$

Compactness and Defects of Compactness

1.83 Theorem [Weak Compactness in Banach Spaces]

Let X be a reflexive Banach space. Let $(x_n)_n \subset X$ be bounded. Then, there exists a subsequence $(x_{n_k})_k$ that weakly converges to some $x \in X$

$$x_{n_k} \xrightarrow{X} x$$

$$\langle \ell | x_{n_k} \rangle_{X^*, X} \xrightarrow[k \rightarrow \infty]{} \langle \ell | x \rangle_{X^*, X}, \quad \forall \ell \in X^*$$

Moreover,

$$\|x\|_X \leq \liminf_k \|x_{n_k}\|_X$$

meaning that, in the weak limit, due to the defects of compactness, mass can escape.

1.84 Theorem [Weak* Compactness in Banach Spaces]

Let X be a separable Banach space. Let $(\ell_n)_n \subset X^*$ be bounded. Then, there exists a subsequence $(\ell_{n_k})_k$ that weakly* converges to some $\ell \in X^*$

$$\ell_{n_k} \xrightarrow{X^*} \ell$$

$$\langle \ell_{n_k} | x \rangle_{X^*, X} \xrightarrow{k \rightarrow \infty} \langle \ell | x \rangle_{X^*, X}, \quad \forall x \in X$$

Moreover,

$$\|\ell\|_{X^*} \leq \liminf_k \|\ell_{n_k}\|_{X^*}$$

meaning that, in the weak* limit, due to the defects of compactness, mass can escape.

1.85 Proposition

Let $(u_n)_n \subset W^{m,p}$ be bounded. Then, there exists an extraction $(u_{n_k})_n$ and some $u \in W^{m,p}$, such that $\forall |\alpha| \leq m$:

$$\partial^\alpha u_{n_k} \xrightarrow{L^2} \partial^\alpha u, \quad \|u\|_{W^{m,p}} \leq \liminf_{k \rightarrow \infty} \|u_{n_k}\|_{W^{m,p}}$$

1.86 Proposition

Let H be Hilbert, $(u_n)_n \subset H$. If $u_n \xrightarrow{H} u$, then,

$$\|u_n - u\|_H \xrightarrow{n \rightarrow \infty} 0 \iff \|u_n\|_H \xrightarrow{n \rightarrow \infty} \|u\|_H$$

1.87 Proposition [Brézis–Lieb Lemma]

Let $p \in [1, \infty)$ and $(u_n)_n \subset L^p(\Omega)$. If $u_n \xrightarrow[n \rightarrow \infty]{a.e.} u$, then,

$$\|u_n - u\|_{L^p} \xrightarrow{n \rightarrow \infty} 0 \iff \|u_n\|_{L^p} \xrightarrow{n \rightarrow \infty} \|u\|_{L^p}$$

Moreover, a stronger result holds

$$\int_{\Omega} \left| |u_n(x)|^p - |u(x)|^p - |u_n(x) - u(x)|^p \right| dx \xrightarrow{n \rightarrow \infty} 0$$

1.88 Proposition

Let $(u_n)_n \subset H^1(\Omega)$ be a bounded sequence. Then, (by Banach–Alaoglu) there exists an extraction $(u_{n_k})_k$ and some $u \in H^1$ such that

$$u_{n_k} \xrightarrow{H^1} u$$

If, additionally Ω bounded and Lipschitz, then (by Rellich–Kondrashev), $(u_n)_n$ is pre-compact in $L^2(\Omega)$

$$\|u_{n_k} - u\|_{L^2(\Omega)} \xrightarrow{k \rightarrow \infty} 0$$

1.89 Theorem [Lions Concentration-Compactness principle]

1.90 Proposition

Let $(u_n) \xrightarrow{L^p} u$, $(v_n) \xrightarrow{(L^p)^*} v$. Then we still can have a defect of compactness so that

$$u_n v_n \not\xrightarrow{\mathcal{D}^*} uv$$

This can happen due to the defects of compactness in the forms of: oscillations, vanishing, concentration, and splitting.

1.91 Theorem

Let $p \in [1, \infty]$, $(u_n) \xrightarrow[n \rightarrow \infty]{L^p} u$ and $v_n \xrightarrow{(L^p)^*} v$. Then,

$$\int u_n v_n \xrightarrow{n \rightarrow \infty} \int uv$$

Let $(u_n) \xrightarrow{L^p} u, (v_n) \xrightarrow{(L^p)^\star} v$ with (u_n) bounded in $W^{1,p}(\mathbb{R}^d)$, (v_n) bounded in $(L^p)^\star$. Then, for any φ bounded and compactly supported,

$$\int_{\mathbb{R}^d} u_n v_n \varphi \xrightarrow{n \rightarrow \infty} \int_{\mathbb{R}^d} uv \varphi$$

1.92 Theorem [Div-Curl Lemma]

Let $(u_n), (v_n) \subset (L^p(\mathbb{R}^d))^d$ be bounded sequences of vector fields with

- 1) $u_n \xrightarrow{L^p} u, v_n \xrightarrow{L^p} v$
- 2) $(\operatorname{div} u_n), (\operatorname{curl} v_n)$ are L^p bounded.

Then,

$$u_n v_n \xrightarrow{\mathcal{D}^\star} uv$$

The proof uses the Helmholtz decomposition in the Fourier space.

1.93 Proposition [Weak Continuity of the Determinant]

Let $(u_n) \subset (W^{1,d}(\mathbb{R}^d))^d$ be a bounded sequence of vector fields. If

$$u_n \xrightarrow{W^{1,d}} u$$

Then,

$$\det(\nabla u_n) \xrightarrow{\mathcal{D}^\star} \det(\nabla u)$$

1.94 Theorem [Compactness with a chain of spaces]

Let $E_1 \subset E_2 \subset E_3$ Banach spaces with E_1 compactly embedded into E_2 , let $p \in [1, \infty]$. If a bounded subset $A \subset L^p([0, T], E_1)$ has the equicontinuity property

$$\sup_{f \in A} \|f(x+h) - f(x)\|_{L_x^p([0, T-h], E_3)} \xrightarrow{h \rightarrow 0} 0$$

then, A is relatively compact in $L^p([0, T], E_2)$

1.95 Proposition [Aubin–Lions Lemma]

Let X Banach, let $u \in L^p([0, T]; X)$, define for $a > 0$ the right mean function $M_a \in C([0, T - a]; X)$

$$M_a(f)(t) = \frac{1}{a} \int_t^{t+a} f(s) \, ds$$

Then, for $p \leq q$

$$\|M_a(f)\|_{L^q([0, T-a])} \leq a^{1/q-1/p} \|f\|_{L^p([0, T])}$$

2.0 Harmonic Analysis

2.1 Schwartz Space

2.1 Definition [Schwartz Space $\mathcal{S}$]

Let $n \in \mathbb{N}, \alpha \in \mathbb{N}^d$. The Schwartz space is defined as the set

$$\mathcal{S}(\mathbb{R}^d) := \{\varphi \in C^\infty ; (\forall \alpha : \forall k) \quad [\varphi]_{\alpha,k} < \infty\}$$

where the seminorms are defined as

$$[\varphi]_{\alpha,k} := \sup_{x \in \mathbb{R}^d} (1 + |x|^2)^k |\partial^\alpha \varphi(x)|$$

This space is a Fréchet space with the Schwartz topology, however one rarely needs strong convergence results.

2.2 Definition [Tempered Distributions $\mathcal{S}^*$]

The space of tempered distributions is defined to be a topological dual of a Schwartz space

$$\mathcal{S}^* := (\mathcal{S})^*$$

It is the set of distributions that are continuous in the following sense

$$(\forall T \in \mathcal{S}^* : \exists n, A, C : \forall \varphi \in \mathcal{S}) \quad |T(\varphi)| \leq C \sup_{|\alpha| \leq A} [\varphi]_{\alpha,n}$$

We have some notions of strong topologies on both $\mathcal{S}, \mathcal{S}^*$ but in practice the derived $\sigma(\mathcal{S}, \mathcal{S}^*), \sigma(\mathcal{S}^*, \mathcal{S})$ topologies will be used.

2.3 Proposition

We have the following chain of inclusions over $p \in [1, \infty]$ on $\mathbb{R}^d$:

$$\mathcal{D} \subsetneq \mathcal{S} \subsetneq L^p \subsetneq \mathcal{S}^* \subsetneq \mathcal{D}^*$$

With non-examples:

- 1) $e^{-|x|^2} \in \mathcal{S}(\mathbb{R}^d)$, but not in $\mathcal{D}(\mathbb{R}^d)$

- 2) $(x \mapsto 1) \in L^\infty$, but not in $\mathcal{S}$
- 3) $\delta_0 \in \mathcal{S}^*$, but not in L^p
- 4) $e^x \in \mathcal{D}^*(\mathbb{R})$, but not in $\mathcal{S}^*(\mathbb{R})$

2.2 Fourier Transform

2.4 Definition [Fourier Transform]

Let $f \in \mathcal{S}(\mathbb{R}^d; \mathbb{C})$, the Fourier transform is a linear mapping

$$\begin{aligned} \mathcal{F} : \mathcal{S}(\mathbb{R}^d; \mathbb{C}) &\rightarrow \mathcal{S}(\mathbb{R}^d; \mathbb{C}) \\ f &\mapsto \widehat{f}(\xi) = \int_{\mathbb{R}^d} e^{-i x \cdot \xi} f(x) \, dx \end{aligned}$$

2.5 Theorem

The Fourier transform is invertible with the inverse

$$\mathcal{F}^{-1} = (2\pi)^{-d} \mathcal{F}^*$$

with

$$\mathcal{F}^*[f](\xi) = \int_{\mathbb{R}^d} e^{+i x \cdot \xi} f(x) \, dx$$

The Fourier transform is an isometric isomorphism on $\mathcal{S}(\mathbb{R}^d; \mathbb{C})$.

2.6 Theorem

Fourier transform is self-transposed and we can be thus defined as generally as on the space of tempered distributions

$$\begin{aligned} \mathcal{F} : \mathcal{S}^*(\mathbb{R}^d; \mathbb{C}) &\rightarrow \mathcal{S}^*(\mathbb{R}^d; \mathbb{C}) \\ T &\mapsto \mathcal{F}T \end{aligned}$$

via transposition

$$\langle \mathcal{F}T | f \rangle_{\mathcal{S}^*, \mathcal{S}} = \langle T | \mathcal{F}f \rangle_{\mathcal{S}^*, \mathcal{S}} \quad \forall f \in \mathcal{S}$$

2.7 Theorem [Fourier-Plancherel, Parseval]

Let $f \in \mathcal{S}(\mathbb{R}^d)$. Then,

$$(2\pi)^d (f | g)_{L^2} = \left(\widehat{f} | \widehat{g} \right)_{L^2}$$

and, in particular,

$$(2\pi)^d \|f\|_{L^2} = \|\widehat{f}\|_{L^2}$$

2.8 Theorem [Extension to L^2]

By the extension of linear maps, $\mathcal{F}$ can be extended to $L^2(\mathbb{R}^d; \mathbb{C})$, and by Fourier-Plancherel, $\mathcal{F}$ is an isometric isomorphism on L^2 . Note that if $(f \in L^2, f \notin L^1)$, there will be no explicit formula to compute $\widehat{f}$.

By density, $\mathcal{F}$ extends to L^p spaces for $p \in [1, \infty)$.

2.9 Remark

Fourier transform and its combinations appear in many fields and problems

Domain	Space	→	Fourier Domain (Space)
(x)	Position space	→	(ξ) : Spatial frequency space
(t)	Time domain	→	(τ) : Frequency space
(t, x)		→	(τ, ξ) : Space-time frequency space
(t, x)	Physical space	→	(t, ξ) : Momentum space (spatial Fourier space)
(t, x)		→	(τ, x) : Spectral space (temporal Fourier space)

2.3 Fourier Operators

2.10 Definition [Fourier Multiplier]

An operator T_m is a **Fourier (Multiplier) Operator** if there is a function $m(\xi)$ called a **Fourier Symbol**, such that it has a following form

$$T_m := m(\nabla/i) := \mathcal{F}^{-1} m(\xi) \mathcal{F}$$

$$\mathcal{F}[T_m f(x)] = \mathcal{F}[m(\nabla/i) f(x)] = m(\xi) \mathcal{F} f = m(\xi) \widehat{f}(\xi)$$

$\mathcal{F}^{-1} m(\xi) \mathcal{F} f$ should be understood as $\mathcal{F}^{-1} (m(\xi) \mathcal{F}[f])$.

2.11 Example [Fourier Operators]

Let $k \in \mathbb{N}^\times, s \in \mathbb{C}$

Derivative $\partial_x^k = \mathcal{F}^{-1}(i\xi)^k \mathcal{F}$

$$\mathcal{F}[\partial_x^k f(x)] = \mathcal{F}[\mathcal{F}^{-1}(i\xi)^k \mathcal{F}f(x)] = (i\xi)^k \widehat{f}(\xi)$$

Fractional (Riesz) derivative $D_x^s := |\partial_x|^s := \mathcal{F}^{-1}|\xi|^s \mathcal{F}$

$$\mathcal{F}[D_x^s f(x)] = \mathcal{F}[\mathcal{F}^{-1}|\xi|^s \mathcal{F}f(x)] = |\xi|^s \widehat{f}(\xi)$$

Hilbert transform $\mathcal{H} = \mathcal{F}^{-1}(-i \operatorname{sgn}(\xi)) \mathcal{F}$

$$\mathcal{F}[\mathcal{H}f(x)] = \mathcal{F}[\mathcal{F}^{-1}(-i \operatorname{sgn}(\xi)) \mathcal{F}f(x)] = -i \operatorname{sgn}(\xi) \widehat{f}(\xi)$$

The connection between ordinary derivative and Riesz derivative is

$$\partial_x^k = (D_x \mathcal{H})^k = (\mathcal{H} D_x)^k.$$

Riesz derivative is sometimes written in the form of a fractional Laplacian

$$|\partial_x|^2 = (-\Delta)^{s/2}.$$

Japanese bracket $\langle D_x \rangle^s = \mathcal{F}^{-1}(1 + |\xi|^2)^{s/2} \mathcal{F}$

$$\mathcal{F}[\langle D_x \rangle^s f(x)] = \mathcal{F}[\mathcal{F}^{-1}\langle \xi \rangle^s \mathcal{F}f(x)] = \langle \xi \rangle^s \widehat{f}(\xi)$$

2.12 Definition [Smooth Cut-off Function]

A function $\eta(x) \in C^\infty(\mathbb{R})$ is called a **smooth cut-off** function (or a **bump** function) supported on $[-1, 1]$ iff

$$\eta(x) \in [0, 1]; \quad \eta(x) = 1, \quad x \in \left[-\frac{1}{2}, \frac{1}{2}\right]; \quad \eta(x) = 0, \quad x \in \mathbb{R} \setminus [-1, 1]$$

2.4 Sobolev–Fourier Spaces

2.13 Definition [Sobolev–Fourier Spaces]

Let $s \in \mathbb{R}$, we define

$$H^s(\mathbb{R}^d) = \{u \in \mathcal{S}'(\mathbb{R}^d); \|u\|_{H^s} < \infty\}$$

with

$$\|u\|_{H^s} := \|\langle \xi \rangle^s \widehat{u}(\xi)\|_{L^2} = \left(\int_{\mathbb{R}^d} |\widehat{u}(\xi)|^2 (1 + |\xi|^2)^s d\xi \right)^{1/2}$$

2.14 Proposition

Fourier transform is an isometric isomorphism between Hilbert spaces

$$H^s(\mathbb{R}^d) \stackrel{\mathcal{F}}{\cong} L^2(\mathbb{R}^d, (1 + |\xi|^2)^s d\xi)$$

2.15 Proposition

Let $s \in \mathbb{R}$. Then, $\mathcal{S}(\mathbb{R}^d)$ is dense in $H^s(\mathbb{R}^d)$

2.16 Proposition [Duality]

Let $s \in \mathbb{R}$. Then, $(H^s(\mathbb{R}^d))^ = H^{-s}(\mathbb{R}^d)$*

2.17 Theorem [Embeddings]

Let $s > d/2$. Then, H^s is continuously embedded into the space of functions that tend to 0 at infinities.

Let $0 \leq s < d/2$. Then, H^s is continuously embedded into $L^{\frac{2d}{d-2s}}$ and

$$\|f\|_{L^p} \lesssim \|f\|_{\dot{H}^s}$$

Trace, Extension

2.18 Theorem [Trace, Lifting]

Let $\Omega \subset \mathbb{R}^d$ be a domain with Lipschitz boundary, let $u \in H^1(\Omega)$. Then, there exists a **Trace** operator in $\mathcal{L}(H^1(\Omega), H^{\frac{1}{2}}(\partial\Omega))$

$$\begin{aligned}\gamma_0 : H^1(\Omega) &\rightarrow H^{\frac{1}{2}}(\partial\Omega) \\ u &\mapsto \gamma_0 u = "u|_{\partial\Omega}"\end{aligned}$$

$\partial\Omega$ is a null set w.r.t. Lebesgue measure on $\mathbb{R}^d$, thus it's impossible to directly restrict an H^1 function to it. Instead, we consider $\varphi \in C^\infty(\overline{\Omega})$ and obtain an estimate for continuity of $\gamma_0 : C^\infty(\overline{\Omega}) \rightarrow H^{\frac{1}{2}}(\partial\Omega)$

$$\|\gamma_0(\varphi)\|_{H^{\frac{1}{2}}(\partial\Omega)} \lesssim \|\varphi\|_{H^1(\Omega)}, \quad \forall \varphi \in (C^\infty(\overline{\Omega}), \|\cdot\|_{H^1(\Omega)})$$

C^∞ is dense in H^1 and thus we can extend trace to the whole space H^1 .

Thus, we lose $1/2$ of derivative by reducing the dimension.

The results extends to $H^s(\mathbb{R}^d) \rightarrow H^{s-\frac{1}{2}}(\mathbb{R}^{d-1})$ for $s > 1/2$, and can be obtained by extension from the Schwartz class.

Moreover, trace admits an explicit right inverse L ,

$$\gamma_0 \circ L = \mathbf{1}_{H^{s-\frac{1}{2}}(\mathbb{R}^{d-1})}$$

2.19 Theorem [Extension by 0]

Let $u \in H^s(\mathbb{R}_+^*)$. The extension by 0 as a mapping $H^s(\mathbb{R}_+^*) \rightarrow H^s(\mathbb{R})$ is continuous iff $s < 1/2$. For $s = 1/2$ iff

$$\int_0^\infty \frac{|u(x)|^2}{x} dx < \infty$$

We denote by $H_{00}^{\frac{1}{2}}$ the space of functions that can be extended by 0, it's a complete space for the norm

$$\|u\|_{H_{00}^{1/2}} = \int_0^\infty \frac{|u(x)|^2}{x} dx + \|u\|_{H^{1/2}}$$

2.20 Definition [Zero-trace Sobolev Spaces]

Let $s > 0$, we define

$$H_0^s(\Omega) = \overline{\mathcal{D}(\Omega)}^{H^s(\Omega)}$$

2.21 Proposition

Let $s > 0$. Then,

$$(H^s(\Omega))^* = H_0^{-s}(\Omega)$$

Moreover, if $0 < s < 1/2$, then

$$H_0^s(\Omega) = H^s(\Omega)$$

if $s = 1/2$, then γ_0 is continuous but not surjective

$$H_0^{1/2}(\Omega) \subsetneq H^{1/2}(\Omega)$$

if $s > 1/2$, then the trace operator γ_0 is well defined and continuous and so we have the usual characterization

$$H_0^s(\Omega) = \{u \in H^s(\Omega) ; \gamma_0(u) = 0\}$$

Results

2.22 Proposition [Stein Derivative]

Let $s \in (0, 1)$, then the $H^s(\mathbb{R}^d)$ norm has this equivalent

$$\|u\|_{H^s} \sim \|u\|_{L^2} + \left(\iint_{(\mathbb{R}^d)^2} \frac{|u(x+h) - u(x)|^2}{|h|^{d+2s}} dx dh \right)^{1/2}$$

2.23 Theorem [Rellich]

Let $s, R > 0$. Denote by $H_R^s(\mathbb{R}^d)$ the space of $H^s(\mathbb{R}^d)$ functions which vanish outside of $B_{\mathbb{R}^d}(0, R)$. Then, for any bounded subset $A \subset H_R^s(\mathbb{R}^d)$

$$\overline{A}^{L^2(\mathbb{R}^d)} \text{ is compact in } L^2(\mathbb{R}^d)$$

2.5 Interpolation

2.24 Definition [Distribution Function]

Let $f \in L^p(\mathbb{R}^d)$, $\lambda := \text{Leb}_{\mathbb{R}^d}$ the distribution function of f is

$$m_f(t) := \lambda(|f| < t)$$

It's always well-defined by Chebyshev inequality.

2.25 Definition [Decreasing Rearrangement]

Let f be such that

$$m_f(t) \xrightarrow[t \rightarrow \infty]{} 0$$

Then, we can define $f^* : \mathbb{R}_+^* \rightarrow \mathbb{R}_+$ as

$$f^*(r) = \int_0^\infty \mathbf{1}_{r \in [0, m_f(t))} dt = \inf_{s \in \mathbb{R}_+^*} \{m_f(s) \leq r\}$$

2.26 Definition [Lorentz Spaces]

Let $p \in [1, \infty)$, weak L^p spaces are defined to be

$$L^{p,\infty}(\mathbb{R}^d) := \{f ; \sup_{t \in \mathbb{R}_+^*} t^p m_f(t) < \infty\}$$

Its generalization, Lorentz Spaces are defined to be

$$L^{p,q}(\mathbb{R}^d) := \{f ; \left\| f^*(t) t^{1/p} \right\|_{L^q(\mathbb{R}^d, dt/t)} < \infty\}$$

2.27 Definition [Sublinear Operator]

An operator T is called sublinear iff

$$|T(cf)| = |c| |T(f)|, \quad |T(f+g)| \leq |T(f)| + |T(g)|$$

2.28 Definition [Strong Type]

Let $p, q \in [1, \infty]$, a sublinear operator $T : L^p \rightarrow L^q$ is called **of (strong) type (p, q)**

iff

$$\|Tf\|_{L^q} \lesssim \|f\|_{L^p}$$

2.29 Definition [Weak Type]

Let $p, q \in [1, \infty]$, a sublinear operator $T : L^p \rightarrow L^q$ is called **of weak type** (p, q) iff

$$\|Tf\|_{L^{q,\infty}} \lesssim \|f\|_{L^p}$$

For $q = \infty$, notions of weak and strong type coincide.

2.30 Theorem [Hölder Interpolation Inequality (log-convexity of $\|\cdot\|_{L^p}$)]

Let $p_0, p_1 \in [1, \infty]$ and $\theta \in [0, 1]$ with

$$\frac{1}{p_\theta} = \frac{\theta}{p_0} + \frac{1-\theta}{p_1}, \quad \text{for some } \theta \in [0, 1]$$

Then,

$$\|f\|_{L^{p_\theta}} \lesssim \|f\|_{L^{p_0}}^\theta \|f\|_{L^{p_1}}^{1-\theta}$$

2.31 Proposition

Let $f \in L^{p_\theta}$, then it can be written as a sum

$$f = g + h$$

with $g \in L^{p_0}, h \in L^{p_1}$

2.32 Theorem [Marcinkiewicz Interpolation]

Let T be a **sublinear** operator and $p_0, p_1, q_0, q_1 \in [1, \infty]$, $p_0 \neq p_1, q_0 \neq q_1$. If T is of **weak type** (p_0, q_0) and of **weak type** (p_1, q_1) , then, T is of **strong type** (p_θ, q_θ) with

$$\left(\frac{1}{p_\theta}, \frac{1}{q_\theta}\right) = \left(\frac{\theta}{p_0} + \frac{1-\theta}{p_1}, \frac{\theta}{q_0} + \frac{1-\theta}{q_1}\right), \quad \forall \theta \in (0, 1)$$

and moreover,

$$\|T\|_{L^{p_\theta} \rightarrow L^{q_\theta}} \lesssim M_0^\theta M_1^{1-\theta}$$

$$\|Tf\|_{L^{q_0,\infty}} \leq M_0 \|f\|_{L^{p_0}}, \quad \|Tf\|_{L^{q_1,\infty}} \leq M_1 \|f\|_{L^{p_1}}$$

2.33 Theorem [Weak Marcinkiewicz Interpolation]

2.34 Lemma [Hadamard Three-Lines Theorem]

let f holomorphic on $0 < \operatorname{Re}(z) < 1$, bounded and continuous on $0 < \operatorname{Re}(z) \leq 1$ with

$$M_0 = \sup_t |f(1+it)|, \quad M_1 = \sup_t |f(it)|$$

Then,

$$|f(\theta + it)| \leq M_0^\theta M_1^{1-\theta}, \quad \forall \theta \in [0, 1]$$

2.35 Theorem [Reisz–Thorin Complex Interpolation]

Let T be a linear operator and $p_0, p_1, q_0, q_1 \in [1, \infty]$. If T is simultaneously of strong type (p_0, q_0) and of (p_1, q_1) , then, T is of strong type (p_θ, q_θ) with

$$\left(\frac{1}{p_\theta}, \frac{1}{q_\theta}\right) = \left(\frac{\theta}{p_0} + \frac{1-\theta}{p_1}, \frac{\theta}{q_0} + \frac{1-\theta}{q_1}\right), \quad \forall \theta \in [0, 1]$$

and moreover,

$$\|T\|_{L^{p_\theta} \rightarrow L^{q_\theta}} \lesssim \|T\|_{L^{p_0} \rightarrow L^{q_0}}^\theta \|T\|_{L^{p_1} \rightarrow L^{q_1}}^{1-\theta}$$

2.36 Theorem [Stein Interpolation]

2.37 Theorem [Hausdorff–Young]

Let $p \in [1, 2]$, the Fourier transform extends continuously to a map $L^p(\mathbb{R}^d) \rightarrow L^{p'}(\mathbb{R}^d)$ for

$$\frac{1}{p} + \frac{1}{p'} = 1$$

with

$$\|\mathcal{F}\|_{L^p \rightarrow L^{p'}} \leq (2\pi)^{d/p'}$$

2.6 Hardy–Littlewood Maximal Function

2.38 Definition [Hardy–Littlewood Maximal Function]

Let $f \in L^1_{\text{loc}}(\mathbb{R}^d)$,

$$\mathcal{M}f(x) := \sup_{t>0} \frac{1}{|B(x, t)|} \int_{B(x, t)} |f(y)| \, dy$$

$\mathcal{M}$ is subadditive and it is continuous for $L^\infty \rightarrow L^\infty$ and $L^1 \rightarrow L^{1,\infty}$

2.39 Theorem [Hardy–Littlewood–Sobolev Inequality]

Let $\alpha \in (0, d)$ and $1 < p < r < \infty$ with

$$1 + \frac{1}{r} = \frac{1}{p} + \frac{\alpha}{d}$$

Then,

$$\| |\cdot|^\alpha \star f \|_{L^r(\mathbb{R}^d)} \lesssim \|f\|_{L^p(\mathbb{R}^d)}$$

2.7 Calderón–Zygmund Theory

2.40 Proposition [Calderón–Zygmund Decomposition]

Let $f \in L^1(\mathbb{R}^d)$ and $\lambda > 0$. Then, there exists a (finite or) countable family of cubes $(Q_j)_{j \in J}$ such that

$$\lambda < \frac{1}{|Q_j|} \int_{Q_j} f(x) dx \leq 2^d \lambda$$

and

$$\|f\|_{L^\infty(\mathbb{R}^d \setminus (\cup Q_j))} \leq \lambda$$

2.41 Definition [Calderón–Zygmund Convolution Kernel]

Calderón–Zygmund Convolution Kernel $K \in C^1(\mathbb{R}^d \setminus \{0\}; \mathbb{C})$ is a function, such that

$$|\nabla K| \lesssim |x|^{-(d+1)}$$

It also defines an associated (singular) Calderón–Zygmund operator $T : f \mapsto K \star f$ that is continuous $L^2 \rightarrow L^2$

2.42 Theorem [Calderón–Zygmund]

Calderón–Zygmund operators are continuous $L^p \rightarrow L^p$ for $p \in (1, \infty)$.

2.8 Multiplier Theory

2.43 Theorem [Mikhlin–Hörmander]

Let $m : \mathbb{R}^d \rightarrow \mathbb{C}$ be a symbol with

$$(\forall \alpha : |\alpha| \leq \frac{d}{2} + 1 : \exists C_\alpha : \forall \xi \in \mathbb{R}^d) : \quad |\xi|^{|\alpha|} |\partial^\alpha m(\xi)| \leq C_\alpha$$

Then, the Fourier multiplier operator

$$\begin{aligned} T_m : \mathcal{S} &\rightarrow \mathcal{S} \\ f &\mapsto \mathcal{F}^{-1} \left(m(\xi) \widehat{f} \right) = \mathcal{F}^{-1} m(\xi) \mathcal{F} f \end{aligned}$$

continuously extends to a bounded operator on $L^p(\mathbb{R}^d) \rightarrow L^p(\mathbb{R}^d)$ for $p \in (1, \infty)$

2.44 Proposition

The following Calderon–Zygmund operators are continuous $L^p \rightarrow L^p$ for $p \in (1, \infty)$

$$R_j : u \mapsto \mathcal{F}^{-1} \left(\frac{\xi_j}{|\xi|} \widehat{u} \right)$$

and

$$\partial_i \partial_j \Delta^{-1} : u \mapsto \mathcal{F}^{-1} \frac{\xi_i \xi_j}{|\xi|^2} \widehat{u}$$

In particular, it means that, for those spaces, Laplacian can control all the derivatives

up to 2

$$\|u\|_{W^{2,p}} \sim \|u\|_{L^p} + \|\Delta u\|_{L^p}$$

2.45 Definition [Bessel Potential Spaces]

Let $s \geq 0$, $p \in (1, \infty)$,

$$H^{s,p}(\mathbb{R}^d) := \{u \in L^p ; \langle (-\Delta)^{s/2} \rangle u = \mathcal{F}^{-1}(\langle \xi \rangle^s \hat{u}) \in L^p\}$$

2.9 Besov Spaces

2.46 Definition [Dyadic Partition of Unity]

2.47 Definition [Besov Spaces]

Besov Spaces are defined as

$$B_{p,q}^s(\mathbb{R}^d) \{u \in \mathcal{S}^*(\mathbb{R}^d) ; \|u\|_{B_{p,q}^s} < \infty\}$$

with the norm

$$\|u\|_{B_{p,q}^s} := \|2^{sj} \Delta_j u\|_{l_j^q L_x^p} = \left(\sum_{j \in \mathbb{N}} (2^{sj} \|\Delta_j u\|_{L_x^p})^q \right)^{1/q}$$

Estimates

2.48 Proposition

Let $p \in [1, \infty]$ and $j \in \mathbb{N}$. Then,

$$\|\Delta_j u\|_{L^p} \lesssim \|u\|_{L^p}$$

2.49 Theorem [Bernstein Inequalities]

Let $\lambda > 0$ and $1 \leq p \leq q \leq \infty$ and $k \in \mathbb{N}$. If $\text{supp}(\widehat{u}) \subset B(0, \lambda)$, then

$$\sup_{|\alpha|=k} \|\partial^\alpha u\|_{L^q} \lesssim \lambda^{k+d(1/p-1/q)} \|u\|_{L^p}$$

If $\text{supp}(\widehat{u}) \subset \{\lambda/2 \leq |\xi| \leq 2\lambda\}$, then

$$\sup_{|\alpha|=k} \|\partial^\alpha u\|_{L^p} \sim \lambda^k \|u\|_{L^p}$$

2.50 Theorem

$B_{p,q}^s$ is Banach for all exponents. $\mathcal{S} \subset B_{p,q}^s$. $\mathcal{D}$ is dense in $B_{p,q}^s$ for $p, q < \infty$

2.51 Proposition [Besov Embeddings]

Let $s \in \mathbb{R}$; $1 \leq p_1 \leq p_2 \leq \infty$; $1 \leq q_1 \leq q_2 \leq \infty$. Then,

$$\|u\|_{B_{p_2,q_2}^{s-d(1/p_1-1/p_2)}} \lesssim \|u\|_{B_{p_1,q_1}^s}$$

2.52 Proposition [Besov Interpolation]

Let $s_1 < s_2$; $1 \leq p, q \leq \infty$ and $\theta \in [0, 1]$. Then,

$$\|u\|_{B_{p,q}^{\theta s_1 + (1-\theta)s_2}} \leq \|u\|_{B_{p,q}^{s_1}}^\theta \|u\|_{B_{p,q}^{s_2}}^{1-\theta}$$

Paraproducts

2.53 Definition [Paraproduct]

Let u be fixed, the map

$$v \mapsto T_v u := \sum_{k \geq 2} S_{k-2} u \Delta_k v$$

Is called paraproduct of v by u , and the decomposition

$$uv = T_v u + T_u v + R(u, v)$$

is called the Bony decomposition

2.54 Theorem

Let $u \in L^\infty, v \in B_{p,q}^s$, $p, q \in [1, \infty], s \in \mathbb{R}$. Then,

$$\|T_u v\|_{B_{p,q}^s} \lesssim \|u\|_{L^\infty} \|v\|_{B_{p,q}^s}$$

and if

$$s_1 + s_2 > 0, \quad 1/p = 1/p_1 + 1/p_2, \quad 1/q = 1/q_1 + 1/q_2$$

we have

$$\|R(u, v)\|_{B_{p,q}^{s_1+s_2}} \lesssim \|u\|_{B_{p_1,q_1}^{s_1}} \|v\|_{B_{p_2,q_2}^{s_2}}$$

2.55 Proposition [Product Rule for Besov Spaces]

Let $u, v \in B_{p,q}^s \cap L^\infty$, $s > 0$. Then,

$$\|uv\|_{B_{p,q}^s} \lesssim \|u\|_{L^\infty} \|v\|_{B_{p,q}^s} + \|v\|_{L^\infty} \|u\|_{B_{p,q}^s}$$

2.10 Littlewood–Payley Theory

2.11 Oscillatory Integrals

2.56 Definition [Oscillatory Integrals]

Oscillatory integral of the first kind is defined to be an integral of the form

$$I(\lambda) = \int e^{i\lambda\phi(x)} \psi(x) \, dx$$

for $\lambda > 0, \phi : \mathbb{R}^d \rightarrow \mathbb{R}, \psi : \mathbb{R}^d \rightarrow \mathbb{C}$

Oscillatory integral of the second kind is defined as

$$T_\lambda f(x) = \int e^{i\lambda\phi(x,y)} K(x, y) f(y) \, dy$$

for $\lambda > 0, \phi : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}, K : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{C}$ and $\psi : \mathbb{R}^d \rightarrow \mathbb{C}$.

In both cases, we are interested in the asymptotic regime, as $\lambda \rightarrow \infty$

2.57 Proposition

2.58 Proposition [Van der Korput Lemma]

2.59 Proposition [Stationary Phase]

2.60 Lemma [Morse Lemma]

3.0 PDE Tools

3.1 General Results

3.1 Lemma [Grönwall]

Let I be an interval, $u \in C(I; \mathbb{R}_+)$ and $\beta \in L^1_{loc}(I; \mathbb{R})$ be integrable. If there exists $C \geq 0$ such that, for all $t \in I$,

$$u(t) \leq C + \int_{t_0}^t \beta(s)u(s) \, ds,$$

or equivalently if u is differentiable and

$$u'(t) \leq \beta(t)u(t), \quad t \in I, \quad u(t_0) = C,$$

then, for all $t \in I$,

$$u(t) \leq C \exp\left(\int_{t_0}^t \beta(s) \, ds\right).$$

3.2 Theorem [Cauchy–Lipschitz–Picard]

Let E Banach, and $F : E \rightarrow E$ be K -Lipschitz. **Then**, given any initial data u_0 , there exists a unique solution $u \in C^1([0, \infty); E)$ of the initial value problem

$$\left\{ \begin{array}{l} \partial_t u(t) = F u(t) \quad t \in [0, \infty) \\ u(0) = u_0 \end{array} \right.$$

3.3 Definition [Unbounded Operator]

A linear operator $A : D(A) \subset H \rightarrow H$ is said to be

- 1) unbounded, if it is not necessarily bounded
- 2) monotone, if

$$(Av \mid v)_H \geq 0, \quad \forall v \in H$$

- 3) maximal monotone, if

$$I_H + A \text{ surjective}$$

3.4 Theorem [Hille–Yosida, Homogeneous]

Let H be Hilbert, let $A : D(A) \rightarrow E$ be a maximal monotone linear operator. **Then**, given any initial data $u_0 \in D(A)$, there exists a unique solution

$$u \in C^1([0, \infty]; H) \cap C([0, \infty]; D(A))$$

of

$$\begin{cases} \partial_t u + Au = 0 & t \in [0, \infty] \\ u(0) = u_0 \end{cases}$$

And moreover,

$$|u(t)| \leq |u_0|, \quad |\partial_t u(t)| = |Au(t)| \leq |Au_0|$$

3.5 Definition [Linear Propagators]

3.6 Theorem [Hille–Yosida, Inhomogeneous]

3.7 Theorem [Hille–Yosida, Nonlinear, Global]

3.8 Theorem [Hille–Yosida, Nonlinear, Local]

3.9 Theorem [Cauchy–Lipschitz–Picard on a real interval]

Let E be a Banach space, $U \subset \mathbb{R} \times E$ open, and $f : U \rightarrow E$. Fix $(t_0, x_0) \in U$ and strictly positive reals $\alpha_0, R_0, K > 0$. For $\alpha > 0$ denote $I_\alpha = [t_0 - \alpha, t_0 + \alpha]$ and $B(x_0, R_0)$. Assume that on $\mathcal{A} := I_{\alpha_0} \times B(x_0, R_0)$ the map f is continuous and Lipschitz in the second variable:

$$\|f(t, x) - f(t, y)\|_E \leq K\|x - y\|_E \quad \text{for all } (t, x), (t, y) \in \mathcal{A}.$$

Set

$$M_0 := \sup_{(t,x) \in \mathcal{A}} \|f(t, x)\|_E \quad (\text{finite by continuity on } \mathcal{A}),$$

and choose

$$\alpha := \min\left(\alpha_0, \frac{R_0}{M_0}, \frac{1}{K}\right).$$

Then there exists a unique $\gamma \in C^1(I_\alpha; E)$ with

$$\gamma(t) = x_0 + \int_{t_0}^t f(s, \gamma(s)) \, ds, \quad t \in I_\alpha,$$

hence $\dot{\gamma}(t) = f(t, \gamma(t))$ for all $t \in I_\alpha$ and $\gamma(t) \in B(x_0, R_0)$ for $t \in I_\alpha$.

3.2 Fixed Point Theorems

3.10 Theorem [Banach–Picard Fixed Point]

Contractive functions on complete metric spaces uniquely fix a point.

Let (X, d) be a non-empty complete metric space, $K < 1$,

$f : X \rightarrow X$ be a continuous mapping with property

$$d(f(x), f(y)) \leq K d(x, y), \quad \forall x, y \in X.$$

Then, there exists a unique $x_\star$, such that $f(x_\star) = x_\star$

*Such f is called a **strict contraction** and $x_\star$ is called a **fixed-point**.*

3.11 Theorem [Kato Fixed Point]

Let V, X be two Banach spaces with $V \hookrightarrow X$ continuously, V separable, reflexive,

$f : V \rightarrow V$, such that, for some $R > 0, k \in (0, 1)$ it holds that

$$(\forall u, v \in \overline{B_V}(0, R)) : \quad \|f(u)\|_V \leq R, \quad \|f(u) - f(v)\|_X \leq k \|u - v\|_X$$

Then, f has a unique fixed point in $\overline{B_V}(0, R)$

3.12 Theorem [Brouwer Fixed Point]

Let E be a **finite-dimensional** vector space, $C \subset E$ a compact, convex subset. If $T : C \rightarrow C$ is continuous, then T has a fixed point (not necessarily unique).

3.13 Theorem [Schauder Fixed Point]

Let E be a **normed** vector space, $C \subset E$ a compact, convex subset. If $T : C \rightarrow C$ is continuous, then T has a fixed point (not necessarily unique).

3.14 Lemma [Leray–Schauder fixed point]

Let E be banach, $T(x, \lambda) : E \times [0, 1] \rightarrow E$ a continuous map with

- 1) T is a compact mapping
- 2) $T(x, 0) \equiv 0$
- 3) $\exists R > 0 : \forall \lambda \in [0, 1] : T(x, \lambda) = x \implies \|x\| < R$

Then, $x \mapsto T(x, 1)$ has a fixed point (not necessarily unique).

3.3 Variational Formulation

3.15 Theorem [Green's Identity]

Let Ω with a Lipschitz boundary, let $u, v \in H^1(\Omega)$, and let $\gamma_0 : H^1(\Omega) \rightarrow H^{1/2}(\partial\Omega)$ be the trace operator and $\gamma_1 : H^1(\Omega) \rightarrow H^{-1/2}(\partial\Omega)$ the weak normal derivative, behaving as (actually defined more technically)

$$\gamma_0(u) = u|_{\partial\Omega}, \quad \gamma_1(u) = \partial_n u|_{\partial\Omega}$$

Then,

$$\int_{\Omega} v \Delta u \, dx = \int_{\partial\Omega} \gamma_0(v) \gamma_1(u) \, d\sigma - \int_{\Omega} \nabla u \cdot \nabla v$$

Let Ω be bounded, $u \in H^1(\Omega)$ and $f \in H^{-1}(\Omega)$. Then, the variational formulation of

the Poisson equation with Homogeneous boundary conditions

$$\left| \begin{array}{l} -\Delta u(x) = f(x), \quad x \in \Omega, \\ \gamma_0(u)(\sigma) = 0, \quad \sigma \in \partial\Omega \end{array} \right.$$

is the following, $\forall v \in H^1(\Omega)$,

$$a(u, v) = \ell(v)$$

Where, ℓ is a continuous linear form

$$\ell(v) = \langle f | v \rangle_{H^{-1}(\Omega), H^1(\Omega)} = \int_{\Omega} f v$$

and $a : H^1 \times H^1 \rightarrow \mathbb{R}$ is a continuous, coercive bilinear form

$$a(u, v) = \int_{\Omega} \nabla u \cdot \nabla v$$

By Lax-Milgram, it has a unique solution in $H^1(\Omega)$, with the norm controlled by

$$\|u\|_{H^1(\Omega)} \leq \frac{1}{\alpha} \|f\|_{H^{-1}(\Omega)}$$

where α is the coercivity (ellipticity) constant of the bilinear form.

For the more general case of inhomogeneous Robin boundary conditions (Dirichlet + Neumann), let $\Omega \subset \mathbb{R}^d$, $u \in H^1(\Omega)$ and $f \in H^{-1}(\Omega)$, $g \in H^{-1}(\partial\Omega)$ and n the outward normal vector on $\partial\Omega$ with $\gamma_1(u) = \partial_n u$

$$\left| \begin{array}{l} -\Delta u(x) + u = f(x), \quad x \in \Omega, \\ \gamma_1(u)(\sigma) + \gamma_0(u)(\sigma) = g(\sigma), \quad \sigma \in \partial\Omega \end{array} \right.$$

via

$$\int_{\Omega} \Delta u v = - \int_{\Omega} \nabla u \cdot \nabla v + \int_{\partial\Omega} \gamma_1(u) \gamma_0(v)$$

Energy Formulation

Given the variational formulation, we can define the energy functional

$$J(u) = \frac{1}{2} a(u, u) - \ell(u)$$

In the case of symmetric bilinear form a , the solution of the variational formulation is characterized as the unique ground state of the energy functional

$$u = \arg \min_{v \in H^1(\Omega)} J(v) = \arg \min_{v \in H^1(\Omega)} \frac{1}{2} a(v, v) - \ell(v)$$

Homogenization

$$\left| \begin{array}{l} - \left[a \left(\frac{x}{\varepsilon} \right) u'_\varepsilon(x) \right]' = f(x), \quad x \in (0, 1), \\ u_\varepsilon(0) = u_\varepsilon(1) = 0 \end{array} \right.$$

As $\varepsilon \rightarrow 0$, it goes to an effective equation

$$\left| \begin{array}{l} - [a_\star u'_\star(x)]' = f(x), \quad x \in (0, 1), \\ u_\star(0) = u_\star(1) = 0 \end{array} \right.$$

with

$$a_\star = \left\langle \frac{1}{a} \right\rangle^{-1} = \left(\int_0^1 \frac{1}{a(x)} dx \right)^{-1}$$

3.16 Proposition

Let $b \in L^\infty(\mathbb{R})$ with period 1 and $\phi \in L^1(\mathbb{R})$. Then,

$$\lim_{\varepsilon \rightarrow 0} \int_{\mathbb{R}} b \left(\frac{x}{\varepsilon} \right) \phi(x) dx = \langle b \rangle \int_{\mathbb{R}} \phi(x) dx$$

Some Things to Remember

3.17 Proposition

Let $u(x) : \mathbb{R}^d \rightarrow \mathbb{R}$ be radial of the form $f(r)$, $r = |x|_{\ell^2}$.

The Laplacian Δ then takes the scalar form

$$\Delta u(x) = f''(r) + \frac{d-1}{r} f'(r)$$

And the gradient has the form

$$\nabla u(x) = f'(r) \frac{1}{r} x$$

3.4 Harmonic Functions

4.0 Differential Equations

4.1 Laplace Equation

Let $u : \Omega \rightarrow \mathbb{R}$

$$\Delta u(x) = 0 \quad (\text{Laplace})$$

Nongomogeneous Laplace also known as Poisson equation

$$-\Delta u(x) = f(x) \quad (\text{Poisson})$$

Dirichlet Problem

Let $\Omega \subset \mathbb{R}^d$ with Lipschitz boundary.

$$\left| \begin{array}{ll} -\Delta u(x) = f(x), & x \in \Omega, \\ u(x) = 0, & x \in \partial\Omega \end{array} \right. \quad (1)$$

Intuition:

Laplace equation models the steady-state flows in equilibrium. Poisson equation adds a source potential.

Symmetries:

Solution:

Conservation Laws:

A variant of a Laplace equation is the related Helmholtz equation

$$-\Delta u(x) = k^2 u(x) \quad (\text{Helmholtz})$$

4.2 Heat Equation

Let $u : I \times \Omega \rightarrow \mathbb{R}$

$$\partial_t u = \Delta u \quad (\text{Heat})$$

If $\Omega = \mathbb{R}^d$, we have an initial value (Cauchy) problem for the heat equation on the whole space. Solving such a problem requires an initial condition.

$$\left| \begin{array}{ll} \partial_t u(t, x) = \Delta u(t, x) & (t, x) \in I \times \mathbb{R}^d \\ u(0, x) = u_0(x) & x \in \mathbb{R}^d \end{array} \right. \quad (\text{Free Heat})$$

If Ω is a regular bounded region, we need to specify a boundary condition.

$$\left| \begin{array}{ll} \partial_t u(t, x) = \Delta u(t, x) & (t, x) \in I \times \Omega \\ u(0, x) = u_0(x) & x \in \Omega \\ u(t, x) = g(t, x) & x \in \partial\Omega \end{array} \right. \quad (2)$$

Intuition:

Symmetries:

Solution:

Conservation Laws:

Nonhomogeneous heat equation

$$\partial_t u - \Delta u = f(t, x) \quad (3)$$

4.3 Wave Equation

Let $u : I \times \Omega \rightarrow \mathbb{R}$

$$\partial_t^2 u(t, x) = \Delta u(t, x) \quad (\text{Wave})$$

$$\left| \begin{array}{ll} \partial_t^2 u(t, x) = \Delta u(t, x) & (t, x) \in I \times \mathbb{R}^d \\ u(0, x) = u_0(x) & x \in \mathbb{R}^d \\ \partial_t u(0, x) = \dot{u}_0(x) & x \in \mathbb{R}^d \end{array} \right. \quad (\text{Free Wave})$$

$$\left| \begin{array}{ll} \partial_t^2 u(t, x) = \Delta u(t, x) & (t, x) \in I \times \Omega \\ u(0, x) = u_0(x) & x \in \Omega \\ \partial_t u(0, x) = \dot{u}_0(x) & \\ u(t, x) = g(t, x) & x \in \partial\Omega \end{array} \right. \quad (4)$$

Intuition:

Symmetries:

Solution:

Conservation Laws:

$$\left| \begin{array}{ll} (\square + m^2) u(t, x) = 0 & (t, x) \in I \times \mathbb{R}^d, \ m \geq 0 \\ u(0, x) = u_0(x) & x \in \mathbb{R}^d \\ \partial_t u(0, x) = \dot{u}_0(x) & \end{array} \right. \quad (\text{Free Klein–Gordon})$$

4.4 Transport Equation

$$\left| \begin{array}{ll} \partial_t u(t, x) + a(t, x) \cdot \nabla u(t, x) = 0 & (t, x) \in I \times \mathbb{R}^d \\ u(0, x) = u_0(x) & x \in \mathbb{R}^d \end{array} \right. \quad (\text{Transport})$$

$$\left| \begin{array}{ll} \partial_t u(t, x) + \nabla \cdot F(u(t, x)) = 0 & (t, x) \in I \times \mathbb{R}^d \\ u(0, x) = u_0(x) & x \in \mathbb{R}^d \end{array} \right. \quad (\text{Conservation})$$

Intuition:

The transport equation describes ...

Solution:

$$\begin{aligned} \partial_t u(t, \gamma(t)) &= \partial_1 u(t, \gamma(t)) + \dot{\gamma}(t) \cdot \nabla_2 u(t, \gamma(t)) \\ \dot{\gamma}(t) &= a(t, \gamma(t)) \end{aligned}$$

Conservation Laws:

The transport equation preserves the L_x^p -norm of the solution for $p \in [1, \infty]$

$$\|u(t, x)\|_{L_x^p(\mathbb{R}^d)} = \|u_0\|_{L_x^p(\mathbb{R}^d)}, \quad t \in I,$$

reflecting the conservation of the transported quantity along the flow.

4.5 Disspersive PDEs

Schrödinger Equation

$$\left| \begin{array}{ll} \partial_t u(t, x) = i\Delta u(t, x) & (t, x) \in I \times \mathbb{R}^d \\ u(0, x) = u_0(x) & x \in \mathbb{R}^d \end{array} \right. \quad (\text{Free Schrödinger})$$

Linear propagator is given by $e^{it\Delta}$

Power nonlinear Schrödinger

$$\left| \begin{array}{ll} \partial_t u = i\partial_x^2 u \pm |u|^{p-1}u & \\ u(0, x) = u_0(x), & x \in \mathbb{R}, t \in \mathbb{R} \end{array} \right. \quad (\text{PNLS})$$

Derivative nonlinear Schrödinger

$$\left| \begin{array}{ll} \partial_t u = i\partial_x^2 u + \lambda \partial_x(|u|^2 u) & \\ u(0, x) = u_0(x), & x \in \mathbb{R}, t \in \mathbb{R} \end{array} \right. \quad (\text{DNLS})$$

KdV Equation

Airy equation, Free KdV

$$\left| \begin{array}{ll} \partial_t u = -\partial_x^3 u & \\ u(0, x) = u_0(x), & x \in \mathbb{R}, t \in \mathbb{R} \end{array} \right. \quad (\text{Airy})$$

Linear propagator is given by $e^{-it\partial_x^3}$

Kortweg-de Vries equation

$$\left| \begin{array}{l} \partial_t u = -\partial_x^3 u - \frac{1}{2} \partial_x^2 u^2 \\ u(0, x) = u_0(x), \end{array} \right. \quad x \in \mathbb{R}, t \in \mathbb{R} \quad (\text{KdV})$$

4.6 Fluid PDEs

Let $\mathbf{u} : I \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ be a velocity field, $\nabla p : I \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ a pressure gradient field

$$\left| \begin{array}{l} \partial_t \mathbf{u}(t, x) + (\mathbf{u}(t, x) \cdot \nabla) \mathbf{u}(t, x) + \nabla p(t, x) = 0 \\ \nabla \cdot \mathbf{u}(t, x) = 0 \\ \mathbf{u}(0, x) = \mathbf{u}_0(x) \end{array} \right. \quad \begin{array}{l} (t, x) \in I \times \mathbb{R}^d, \\ \\ x \in \mathbb{R}^d \end{array} \quad (\text{Incompressible Euler})$$

Let $\nu > 0$ be kinematic viscosity coefficient

$$\left| \begin{array}{l} \partial_t \mathbf{u}(t, x) + (\mathbf{u}(t, x) \cdot \nabla) \mathbf{u}(t, x) + \nabla p(t, x) = \nu \Delta \mathbf{u}(t, x) \\ \nabla \cdot \mathbf{u}(t, x) = 0, \\ \mathbf{u}(0, x) = \mathbf{u}_0(x) \end{array} \right. \quad \begin{array}{l} (t, x) \in I \times \mathbb{R}^d \\ \\ x \in \mathbb{R}^d \end{array} \quad (\text{Incompressible Navier–Stokes})$$

Energy dissipation

$$\partial_t \int |\mathbf{u}| \, dx = -2\nu \int |\nabla \mathbf{u}| \, dx$$

4.7 Solid Mechanics

Small-deformation (linearized) regime. $\mathbf{u} : I \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is a displacement. The strain tensor is a symmetric gradient (symmetrized Jacobian)

$$e(\mathbf{u}) = \frac{1}{2}(\nabla \mathbf{u} + \nabla \mathbf{u}^\top)$$

Its diagonal entries are normal strains, off-diagonal are shear strains. Let v be a directional vector and h some small length. Then, the relative change of length is encoded in

$$h(e(\mathbf{u})v \mid v)$$

Strain tensor also is invariant under rigid motions of rotations and translations.

Cauchy stress tensor is a matrix for isotropic linear elastic solids, under Hooke's law

$$\sigma(\mathbf{u}) = 2\mu e(\mathbf{u}) + \lambda(\nabla \cdot \mathbf{u})\mathbf{1}_d$$

Let $\rho > 0$ be mass density, $\mu > 0$ Sheer modulus, $\lambda > 0$ first Lamé parameter

$$\left| \begin{array}{ll} \rho \partial_t^2 \mathbf{u}(t, x) - \nabla \cdot \sigma(\mathbf{u}(t, x)) = \mathbf{f}_{\text{ext}}(t, x) & (t, x) \in I \times \mathbb{R}^d \\ \mathbf{u}(0, x) = \mathbf{u}_0(x), \quad \partial_t \mathbf{u}(0, x) = \dot{\mathbf{u}}_0(x) & x \in \mathbb{R}^d \end{array} \right. \quad \text{(Linear Elasticity)}$$

Young's modulus

$$E = \frac{\mu(3\lambda + 2\mu)}{\lambda + \mu}$$

Poisson's coefficient

$$\nu = \frac{\lambda}{2(\lambda + \mu)}$$

5.0 Optimization

5.1 General Notions of Optimization

5.1 Theorem [First, Second Order Conditions]

Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ and $x^* \in \mathbb{R}^d$ be a local minimizer of f (unconstrained).

1) if $f \in C^1$, then

$$\nabla f(x^*) = 0$$

2) if $f \in C^2$, then at x^* , then Hessian is *PSD*

$$\mathcal{H}f(x^*) \in \mathcal{S}_+^n$$

5.2 Definition [Convexity]

$K \subset \mathbb{R}^d$ is said to be convex iff

$$(\forall x, y \in K, \forall \theta \in [0, 1]) : \quad \theta x + (1 - \theta)y \in K$$

5.3 Definition [Convexity]

$V \subset \mathbb{R}^d$ is said to be a cone iff

$$(\forall x \in V, \forall t \in \mathbb{R}_+) : \quad tx \in V$$

5.4 Definition [Lower Semi-Continuous Function]

A function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is lower semi-continuous (l.s.c.) iff

$$\liminf_{x \rightarrow x_0} f(x) \geq f(x_0), \quad \forall x_0 \in \mathbb{R}^d$$

Equivalently, iff $\text{epi}(f)$ is closed.

5.5 Definition [Convex Function]

Let $F : \mathbb{R}^d \rightarrow \overline{\mathbb{R}}$ be a function extended by infinity.

1) F is convex iff

$$F(\theta x + (1 - \theta)y) \leq \theta F(x) + (1 - \theta)F(y)$$

2) F is strictly convex iff

$$F(\theta x + (1 - \theta)y) < \theta F(x) + (1 - \theta)F(y)$$

3) F is γ -strongly convex iff

$$F(\theta x + (1 - \theta)y) \leq \theta F(x) + (1 - \theta)F(y) - \frac{\gamma}{2}\theta(1 - \theta) \|x - y\|_2^2$$

5.6 Proposition [Differential Characterization of Convexity]

5.7 Definition [Legendre Dual]

Let $F : \mathbb{R}^d \rightarrow \overline{\mathbb{R}}$ be a function extended by infinity. Its Legendre / Fenchel / Convex dual is the function $F^* : \mathbb{R}^d \rightarrow \overline{\mathbb{R}}$

$$F^*(y) = \sup_{x \in \mathbb{R}^d} (x \mid y) - F(x)$$

KKT Conditions

5.8 Theorem [Lagrange Multipliers, Equality Constraint]

Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ and $\mathbf{g} : \mathbb{R}^d \rightarrow \mathbb{R}^n$ be C^1 . Consider the constrained problem:

$$\left| \begin{array}{l} \arg \min_{x \in \mathbb{R}^d} f(x) \\ \mathbf{g}(x) = 0 \end{array} \right.$$

Define the **Lagrangian**

$$\begin{aligned} \mathcal{L} : \mathbb{R}^d \times \mathbb{R}^n &\rightarrow \mathbb{R} \\ (x, \lambda) &\mapsto f(x) + (\lambda \mid \mathbf{g}(x))_{\mathbb{R}^n} \end{aligned}$$

If

1) $x^\star$ is a local minimizer

2) $\nabla g_1(x_\star), \dots, \nabla g_n(x_\star)$ are linearly independent (equality constraint qualification)

Then, there exists $\lambda^\star \in \mathbb{R}^n$, such that the pair $(x^\star, \lambda^\star)$ satisfies the **Lagrangian stationarity**:

$$\nabla_x \mathcal{L}(x^\star, \lambda^\star) = 0$$

5.9 Theorem [Lagrange Multipliers, Inequality Constraint]

Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ and $\mathbf{h} : \mathbb{R}^d \rightarrow \mathbb{R}^m$ be C^1 . Consider the constrained problem:

$$\left| \begin{array}{l} \arg \min_{x \in \mathbb{R}^d} f(x) \\ \mathbf{h}(x) \leq 0 \end{array} \right.$$

Define the **Lagrangian**

$$\begin{aligned} \mathcal{L} : \mathbb{R}^d \times \mathbb{R}^n &\rightarrow \mathbb{R} \\ (x, \lambda) &\mapsto f(x) + (\lambda \mid \mathbf{h}(x))_{\mathbb{R}^m} \end{aligned}$$

If

1) $x^\star$ is a local minimizer

2) Inequality constraints qualifications are satisfied

Then, there exists $\lambda^\star \in \mathbb{R}^n$, such that the pair $(x^\star, \lambda^\star)$ satisfies the **Lagrangian stationarity**:

$$\nabla_x \mathcal{L}(x^\star, \lambda^\star) = 0$$

5.10 Theorem [Karush–Kuhn–Tucker (KKT) Conditions]

Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ and $\mathbf{g} : \mathbb{R}^d \rightarrow \mathbb{R}^n$ and $\mathbf{h} : \mathbb{R}^d \rightarrow \mathbb{R}^m$ be C^1 . Consider:

$$\left| \begin{array}{l} \arg \min_{x \in \mathbb{R}^d} f(x) \\ \mathbf{g}(x) = 0 \\ \mathbf{h}(x) \leq 0 \end{array} \right.$$

Define the **Lagrangian**

$$\begin{aligned}\mathcal{L} : \mathbb{R}^d \times \mathbb{R}^n \times \mathbb{R}^m &\rightarrow \mathbb{R} \\ (x, \lambda, \eta) &\mapsto f(x) + (\lambda \mid \mathbf{g}(x))_{\mathbb{R}^n} + (\eta \mid \mathbf{h}(x))_{\mathbb{R}^m}\end{aligned}$$

If

- 1) $x^\star$ is a local minimizer
- 2) Equality constraint qualifications are satisfied
- 3) Inequality constraint qualifications are satisfied

Then, there exist multipliers $\lambda^\star \in \mathbb{R}^n$ and $\eta^\star \in \mathbb{R}_+^m$ such that:

i) **Stationarity:**

$$\nabla_x \mathcal{L}(x^\star, \lambda^\star, \eta^\star) = 0$$

ii) **Primal feasibility:**

$$\mathbf{g}(x^\star) = 0, \quad \mathbf{h}(x^\star) \leq 0$$

iii) **Complementary slackness:**

$$\eta^\star \odot \mathbf{h}(x^\star) = 0$$

Lagrangian–Hamiltonian Duality

5.11 Definition [Hamiltonian]

Let $\mathcal{L}$ be a Lagrangian as above, define $H : \mathbb{R}^n \times \mathbb{R}_+^m \rightarrow \mathbb{R}$

$$H(\lambda, \eta) := \inf_{x \in \mathbb{R}^d} \mathcal{L}(x, \lambda, \eta)$$

5.12 Theorem [Weak Duality]

It always holds that

$$d^\star \leq p^\star$$

Where,

$$\left| \begin{array}{l} p^* := \min_{x \in \mathbb{R}^d} f(x) \\ \mathbf{g}(x) = 0 \\ \mathbf{h}(x) \leq 0 \end{array} \right| \quad \left| \begin{array}{l} d^* := \sup_{\substack{\lambda \in \mathbb{R}^n \\ \eta \in \mathbb{R}_+^m}} H(\lambda, \eta) \end{array} \right|$$

5.13 Theorem [Lagrangian Saddle Point]

Let $\mathcal{L}$ be a Lagrangian of an optimization problem. Then,

$$d^* = \sup_{\substack{\lambda \in \mathbb{R}^n \\ \eta \in \mathbb{R}_+^m}} \inf_{x \in \mathbb{R}^d} \mathcal{L}(x, \lambda, \eta) \leq p^* = \inf_{x \in \mathbb{R}^d} \sup_{\substack{\lambda \in \mathbb{R}^n \\ \eta \in \mathbb{R}_+^m}} \mathcal{L}(x, \lambda, \eta)$$

And for primal-dual optimizers (x^*, λ^*, η^*) are the saddle points of $\mathcal{L}$

$$\mathcal{L}(x^*, \lambda, \eta) \leq \mathcal{L}(x^*, \lambda^*, \eta^*) \leq \mathcal{L}(x, \lambda^*, \eta^*)$$

5.2 Algorithms

5.14 Proposition [Gradient Descent]

5.15 Proposition [Newton Algorithm]

6.0 Probability

6.1 Basic Definitions and Results

6.1 Definition [Probability Space]

Let $(\Omega, \mathcal{A})$ be a measurable space; P be a probability measure (finite and normalized to 1). Then $(\Omega, \mathcal{A}, P)$ is called a **probability space**.

$$P : \mathcal{A} \rightarrow [0, 1]$$

6.2 Definition [Random Variable]

Let $X : (\Omega, \mathcal{A}) \rightarrow (\mathbb{R}, \mathcal{B}(\mathbb{R}))$ be measurable, then it's called a real random variable.

6.3 Definition [Law / Distribution of a random variable]

For a random variable X we have

$$\begin{aligned} \mathcal{A} &\xrightarrow{X} \mathcal{B} \\ [0, 1] &\xleftarrow{P} \mathcal{A} \xleftarrow{X^{-1}} \mathcal{B} \end{aligned}$$

Now define

$$\begin{aligned} \mathcal{L}_X : \mathcal{B}(\mathbb{R}) &\rightarrow [0, 1] \\ A &\mapsto \mathcal{L}_X(A) := P \circ X^{-1}(A) = P(X^{-1}(A)) \end{aligned}$$

$\mathcal{L}_X$ is called a **law** (distribution) of a r.v. X

6.4 Definition [Cumulative Distribution Function]

Let $\mathcal{L}_X$ be a law of X

$$\begin{aligned} F_X : \mathbb{R} &\rightarrow [0, 1] \\ x &\mapsto \mathcal{L}_X((-\infty, x]) \end{aligned}$$

F_X is called the **Cumulative Distribution Function**

6.5 Definition [Density]

Let X be a r.v. with law $\mathcal{L}_X$, the **density** of X w.r.t. to a measure ν on $\mathcal{B}(\mathbb{R})$ is a function $f_X : \mathbb{R} \rightarrow \mathbb{R}_+$ such that

$$\mathcal{L}_X(A) = \int_A f_X d\nu, \quad \forall A \in \mathcal{B}(\mathbb{R})$$

If $\nu = \text{Leb}_{\mathbb{R}}$

$$\mathcal{L}_X(A) = \int_A f_X(t) dt, \quad \forall A \in \mathcal{B}(\mathbb{R})$$

6.6 Definition [Atom of a Random Variable]

Let X be a r.v., an element $a \in \mathbb{R}$ is called an atom of X if

$$P(X = a) \neq 0$$

6.7 Definition [Expectation]

Let $X \in L^1(\Omega : \mathbb{R}; P)$, then

$$\mathbb{E}[X] := \int_{\Omega} X dP = \int_{\Omega} X(\omega) dP(\omega)$$

Furthermore, by **Transfer Theorem** $g : \mathbb{R} \rightarrow \mathbb{R}$

$$\mathbb{E}[g(X)] = \int_{\Omega} g(X(\omega)) dP(\omega) = \int_{\mathbb{R}} g(t) d\mathcal{L}_X(t)$$

6.8 Definition [Variance]

Let $X \in L^2(\Omega : \mathbb{R}; P)$

$$\begin{aligned} \text{Var}[X] &= \mathbb{E}[X^2] - \mathbb{E}[X]^2 \\ &= \mathbb{E}[(X - \mathbb{E}[X])^2] \end{aligned}$$

6.9 Definition [Covariance]

Let $X, Y \in L^2(\Omega : \mathbb{R}; P)$

$$\begin{aligned} \text{cov}[X, Y] &= \mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y] \\ &= \mathbb{E}[(X - \mathbb{E}[X])(Y - \mathbb{E}[Y])] \end{aligned}$$

6.10 Definition [Covariance Matrix]

Let $\mathbf{X} = (X_1, \dots, X_d)^\top$ be a random vector in $L^2(\Omega : \mathbb{R}^d)$, its variance is defined as its covariance matrix $\mathbf{\Sigma} \in \mathbb{R}^{d \times d}$

$$\text{Var}[\mathbf{X}] := \mathbf{\Sigma} := \begin{bmatrix} \text{Var}[X_1] & \cdots & \text{cov}[X_1, X_d] \\ \vdots & \ddots & \vdots \\ \text{cov}[X_d, X_1] & \cdots & \text{Var}[X_d] \end{bmatrix} = (\text{cov}[X_i, X_j])_{i,j}$$

Covariance matrix is always in $\mathcal{S}_+^d$

6.11 Remark

If X takes values in $\mathbb{Z}_+$,

$$\begin{aligned}\mathbb{E}[X] &= \sum_{k \geq 0} k P(X = k) \\ \mathbb{E}[X] &= \sum_{k \geq 1} P(X \geq k)\end{aligned}$$

6.12 Definition [Convergence almost surely]

$(X_n)_{n \in \mathbb{N}}$ converges to X almost surely whenever

$$\begin{aligned}X_n &\xrightarrow[n \rightarrow \infty]{\text{a.s.}} X \\ P\left(\left\{\omega : \lim_{n \rightarrow \infty} X(\omega) = X_n(\omega)\right\}\right) &= 1\end{aligned}$$

6.13 Definition [Convergence in Probability]

$(X_n)_{n \in \mathbb{N}}$ converges to X in Probability whenever

$$\begin{aligned}X_n &\xrightarrow[n \rightarrow \infty]{P} X \\ P(\|X(\omega) - X_n(\omega)\| > \epsilon) &= 0\end{aligned}$$

6.14 Definition [Convergence in Distribution]

$(X_n)_{n \in \mathbb{N}}$ converges to X in Distribution whenever for all h bounded, continuous

$$\begin{aligned}X_n &\xrightarrow[n \rightarrow \infty]{(d)} X \\ \mathbb{E}[h(X_n)] &\xrightarrow[n \rightarrow \infty]{} \mathbb{E}[h(X)]\end{aligned}$$

6.15 Theorem [Borel-Canteli]

Let $(E_n)_n$ be a sequence of events

$$\sum_{n \in \mathbb{N}} P(E_n) < \infty \implies P(\limsup_{n \rightarrow \infty} E_n) = P\left(\bigcap_{n=1} \bigcup_{k=n} E_k\right) = 0.$$

Additionally, if E_n are independent,

$$\sum_{n \in \mathbb{N}} P(E_n) = \infty \implies P(\limsup_{n \rightarrow \infty} E_n) = P\left(\bigcap_{n=1} \bigcup_{k=n} E_k\right) = 1.$$

6.16 Definition [Characteristic Function]

Let X be a r.v. in $\mathbb{R}$

$$\begin{aligned} \varphi_X : \mathbb{R} &\rightarrow \mathbb{C} \\ t &\mapsto \mathbb{E}[\exp(itX)] = \mathbb{E}[e^{itX}] \end{aligned}$$

Let $\mathbf{X}$ be a r.v. in $\mathbb{R}^d$

$$\begin{aligned} \varphi_{\mathbf{X}} : \mathbb{R}^d &\rightarrow \mathbb{C} \\ t &\mapsto \mathbb{E}[\exp(it \cdot \mathbf{X})] = \mathbb{E}[e^{it^\top \mathbf{X}}] \end{aligned}$$

6.17 Proposition [Relationship Between Convergences]

Let $(X_n)_{n \in \mathbb{N}}$ be a sequence of r.v. in $\mathbb{R}$ and X be an r.v. in $\mathbb{R}$

$$\begin{aligned} a.s. &\implies P \implies (d) \\ L^p &\implies L^1 \implies P \end{aligned}$$

Additionally, if $X_n \xrightarrow[n \rightarrow \infty]{(d)} X = c$, where $c \in \mathbb{R}$, then

$$P \iff (d)$$

6.18 Theorem [Portmanteau]

Let $(X_n)_{n \in \mathbb{N}}$ be a sequence of r.v. in $\mathbb{R}$ and X be an r.v. in $\mathbb{R}$.

Let A be a set of atoms of X . Then,

$$X_n \xrightarrow[n \rightarrow \infty]{(d)} X \iff P(X_n \leq t) \xrightarrow[n \rightarrow \infty]{} P(X \leq t), \quad \forall t \in \mathbb{R}/A$$

6.19 Theorem [Continuous Mapping]

Let $(\mathbf{X}_n)_{n \in \mathbb{N}}$ be a sequence of r.v. in $\mathbb{R}^d$ and $\mathbf{X}$ be an r.v. in $\mathbb{R}^d$

Let $f : \mathbb{R}^d \rightarrow \mathbb{R}^p$ be a continuous function.

Then,

$$\begin{aligned}\mathbf{X}_n &\xrightarrow[n \rightarrow \infty]{a.s.} \mathbf{X} \implies f(\mathbf{X}_n) \xrightarrow[n \rightarrow \infty]{a.s.} f(\mathbf{X}) \\ \mathbf{X}_n &\xrightarrow[n \rightarrow \infty]{P} \mathbf{X} \implies f(\mathbf{X}_n) \xrightarrow[n \rightarrow \infty]{P} f(\mathbf{X}) \\ \mathbf{X}_n &\xrightarrow[n \rightarrow \infty]{(d)} \mathbf{X} \implies f(\mathbf{X}_n) \xrightarrow[n \rightarrow \infty]{(d)} f(\mathbf{X})\end{aligned}$$

6.20 Theorem [Lévy]

Let $(X_n)_{n \in \mathbb{N}}$ be a sequence of r.v. in $\mathbb{R}$ and X be an r.v. in $\mathbb{R}$

Let $\varphi_{X_n}(t)$ denote the characteristic function of X_n .

If

$$\varphi_{X_n}(t) \xrightarrow[n \rightarrow \infty]{} \varphi(t), \quad \forall t \in \mathbb{R}$$

Then,

$$\begin{aligned}\varphi(t) &= \varphi_X(t) \in C^0 \\ X_n &\xrightarrow[n \rightarrow \infty]{(d)} X\end{aligned}$$

6.21 Theorem [Slutsky]

Let $(\mathbf{X}_n)_{n \in \mathbb{N}}$, $(\mathbf{Y}_n)_{n \in \mathbb{N}}$ be sequences of r.v. in $\mathbb{R}^d$, $\mathbb{R}^p$.

Let $\mathbf{X}$ be a r.v. in $\mathbb{R}^d$, let $\mathbf{c} \in \mathbb{R}^p$ be a fixed vector.

Then,

$$\begin{pmatrix} \mathbf{X}_n \\ \mathbf{Y}_n \end{pmatrix} \xrightarrow[n \rightarrow \infty]{(d)} \begin{pmatrix} \mathbf{X} \\ \mathbf{c} \end{pmatrix} \iff \begin{cases} \mathbf{X}_n \xrightarrow[n \rightarrow \infty]{(d)} \mathbf{X} \\ \mathbf{Y}_n \xrightarrow[n \rightarrow \infty]{(d)} \mathbf{c} \end{cases}$$

Theorem gives ($\iff$).

6.22 Proposition [Convergence in Distribution]

Let $(\mathbf{X}_n)_{n \in \mathbb{N}}$, $(\mathbf{Y}_n)_{n \in \mathbb{N}}$ be sequences of r.v. in $\mathbb{R}^d$, $\mathbb{R}^p$.

Let $\mathbf{X}$ be a r.v. in $\mathbb{R}^d$, let $\mathbf{Y} \in \mathbb{R}^p$.

If $\mathbf{X}$ is independent of $\mathbf{Y}$ and $\mathbf{X}_n$ is independent of $\mathbf{Y}_n$ for all $n \in \mathbb{N}$. Then,

$$\begin{pmatrix} \mathbf{X}_n \\ \mathbf{Y}_n \end{pmatrix} \xrightarrow[n \rightarrow \infty]{(d)} \begin{pmatrix} \mathbf{X} \\ \mathbf{Y} \end{pmatrix} \iff \begin{cases} \mathbf{X}_n \xrightarrow[n \rightarrow \infty]{(d)} \mathbf{X} \\ \mathbf{Y}_n \xrightarrow[n \rightarrow \infty]{(d)} \mathbf{Y} \end{cases}$$

6.23 Theorem [R.v. Generator]

Let Z be a real r.v. with density w.r.t. $\text{Leb}_{\mathbb{R}}$. Let F be CDF of Z . Then, $F(Z) \sim \mathcal{U}(0, 1)$

6.2 Limit Theorems

6.24 Theorem [Law of Large Numbers]

Let $X_1, \dots, X_n, \dots$ be i.i.d. random variables with 1 moment; let

$$\bar{X}_n := \frac{1}{n} \sum_{k=1}^n X_k$$

Then,

$$\bar{X}_n \xrightarrow[n \rightarrow \infty]{\text{a.s.}} \mathbb{E}[X_1]$$

6.25 Theorem [Central Limit Theorem]

Let $X_1, \dots, X_n, \dots$ be i.i.d. random variables with 2 moments.

Then,

$$\sqrt{n}(\bar{X}_n - \mathbb{E}[X_1]) \xrightarrow[n \rightarrow \infty]{(d)} \mathcal{N}(0, \text{Var}[X_1])$$

6.26 Theorem [Multivariate Central Limit Theorem]

Let $\mathbf{X}_1, \dots, \mathbf{X}_n, \dots$ be i.i.d. random variables in $\mathbb{R}^d$ with 2 moments.

Then,

$$\sqrt{n}(\bar{\mathbf{X}}_n - \mathbb{E}[\mathbf{X}_1]) \xrightarrow[n \rightarrow \infty]{(d)} \mathcal{N}_d(\mathbf{0}, \Sigma)$$

6.27 Theorem [Delta-Method]

Let $(T_n)_{n \geq 1}$ be a sequence of r.v. with

$$\rho_n(T_n - \theta) \xrightarrow[n \rightarrow \infty]{(d)} \mathcal{N}(0, \sigma^2)$$

for some $\theta \in \mathbb{R}, \sigma^2 > 0, \rho_n$ increasing sequence.

We would say that T_n is asymptotically normal around θ with rate ρ_n and asymptotic variance σ^2 (though it may have nothing to do with the actual limiting variance of T_n).

Let $g : \mathbb{R} \rightarrow \mathbb{R}$ be differentiable at θ with $g'(\theta) \neq 0$. Then,

$$\rho_n(g(T_n) - g(\theta)) \xrightarrow[n \rightarrow \infty]{(d)} \mathcal{N}(0, \sigma^2 \cdot (g'(\theta))^2)$$

In practice, usually, $T_n = \bar{X}_n, \theta = \mathbb{E}[X_1], \sigma^2 = \text{Var}[X_1], \rho_n = \sqrt{n}$.

6.28 Theorem [Multivariate Delta-Method]

Let $(\mathbf{T}_n)_{n \geq 1}$ be a sequence of r.v. in $\mathbb{R}^d$ with

$$\rho_n(\mathbf{T}_n - \boldsymbol{\theta}) \xrightarrow[n \rightarrow \infty]{(d)} \mathcal{N}_d(\mathbf{0}, \boldsymbol{\Sigma})$$

for some $\boldsymbol{\theta} \in \mathbb{R}^d, \boldsymbol{\Sigma} \in \mathcal{S}_+^d$

Let $g : \mathbb{R}^d \rightarrow \mathbb{R}^p$ be differentiable at $\boldsymbol{\theta}$ with $\mathcal{J}_g(\boldsymbol{\theta}) \neq 0$. Then,

$$\rho_n(g(\mathbf{T}_n) - g(\boldsymbol{\theta})) \xrightarrow[n \rightarrow \infty]{(d)} \mathcal{N}_d(\mathbf{0}, \mathcal{J}_g(\boldsymbol{\theta}) \boldsymbol{\Sigma} \mathcal{J}_g^\top(\boldsymbol{\theta})),$$

where $\mathcal{J}_g(\boldsymbol{\theta}) = [\mathcal{J}g](\boldsymbol{\theta})$ is Jacobian of g evaluated at $\boldsymbol{\theta}$

$$\mathcal{J}_g(\mathbf{x}) := \begin{pmatrix} \partial_{x_1} g_1(\mathbf{x}) & \cdots & \partial_{x_d} g_1(\mathbf{x}) \\ \vdots & \ddots & \vdots \\ \partial_{x_1} g_p(\mathbf{x}) & \cdots & \partial_{x_d} g_p(\mathbf{x}) \end{pmatrix} = (\partial_{x_j} g_i(\mathbf{x}))_{i,j} \in \mathbb{R}^{p \times d}$$

6.3 Gaussian and Chi-squared

6.29 Definition [Gaussian (Normal) Distribution]

Univariate Gaussian Distribution $\mathcal{N}(\mu, \sigma^2)$ for some $\mu \in \mathbb{R}, \sigma^2 > 0$ is a distribution on

$\mathcal{B}(\mathbb{R})$ with the following density $f : \mathbb{R} \rightarrow \mathbb{R}_+$, w.r.t. $\text{Leb}_{\mathbb{R}}$

$$f(x) = \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{1}{2} \frac{(x - \mu)^2}{\sigma^2}\right)$$

6.30 Proposition

6.31 Definition [Gaussian Vector]

Let $\mathbf{X} = (X_1, \dots, X_d)^\top$ be a random vector. It is called a Gaussian vector if any linear combination of $X_1, \dots, X_d$ is a univariate Gaussian r.v.

6.32 Proposition

Let $X_1, \dots, X_d$ be independent Gaussian random variables, then $\mathbf{X} = (X_1, \dots, X_d)^\top$ is a Gaussian vector

6.33 Proposition [Multivariate Gaussian Distribution]

Let $\mathbf{X}$ be a Gaussian vector. It has 2 moments, and we can denote $\boldsymbol{\mu} = \mathbb{E}[\mathbf{X}]$, $\boldsymbol{\Sigma} = \text{Var}[\mathbf{X}]$. Then, its characteristic function is

$$\varphi_{\mathbf{X}}(\mathbf{t}) = \mathbb{E}[e^{i\mathbf{t}^\top \mathbf{X}}] = \exp(i\mathbf{t}^\top \boldsymbol{\mu} - \frac{1}{2} \mathbf{t}^\top \boldsymbol{\Sigma} \mathbf{t})$$

and the distribution of X can be denoæted as $\mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$

6.34 Proposition

Let $X \sim \mathcal{N}_d(\boldsymbol{\mu}, \boldsymbol{\Sigma})$, for some $\boldsymbol{\mu} \in \mathbb{R}^d$, $\boldsymbol{\Sigma} \in \mathcal{S}_+^d$.

The following holds

$$X \text{ has a density w.r.t. } \text{Leb}_{\mathbb{R}^d} \iff \boldsymbol{\Sigma} \in \mathcal{S}_{++}^d$$

and the density $f : \mathbb{R}^d \rightarrow \mathbb{R}_+$ is

$$f(\mathbf{x}) = (2\pi)^{-d/2} \det(\boldsymbol{\Sigma})^{-1/2} \exp\left(-\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})^\top \boldsymbol{\Sigma}^{-1}(\mathbf{x} - \boldsymbol{\mu})\right)$$

6.35 Proposition

Let $\mathbf{X}$ be a Gaussian vector of size $d \geq 2$. Write

$$X = \begin{pmatrix} Y \\ Z \end{pmatrix} \quad \text{Var}[X] = \begin{bmatrix} \text{Var}[Y] & \text{cov}[Y, Z] \\ \text{cov}[Z, Y] & \text{Var}[Z] \end{bmatrix}$$

The following holds

$$Y \text{ is independent of } Z \iff \text{cov}[Y, Z] = 0$$

6.36 Definition [Chi-squared]

Let $Z_1, \dots, Z_d \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(0, 1)$, then the distribution of the random variable

$$\chi_d^2 := \sum_{i=1}^d Z_i^2$$

is called "chi-squared distribution with d degrees of freedom".

6.37 Theorem [Cochran]

Let $X_1, \dots, X_n \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(\mu, \sigma^2)$; $\mu \in \mathbb{R}, \sigma^2 > 0, n \geq 2$.

Denote $\hat{\mu}_n := \bar{X}_n$, $\hat{\sigma}_n^2 := \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X}_n)^2$.

Then,

- 1) $\hat{\mu}_n \sim \mathcal{N}(\mu, \sigma^2/n)$
- 2) $n\hat{\sigma}_n^2/\sigma^2 \sim \chi_{n-1}^2$
- 3) $\hat{\mu}_n$ is independent of $\hat{\sigma}_n^2$

6.38 Definition [Student-t distribution]

Let $d \geq 1$; $U \sim \mathcal{N}(0, 1)$, $V \sim \chi_d^2$ and U independent of V .

Then, the Student-t distribution with d -degrees of freedom is the distribution of

$$T = \frac{U}{\sqrt{V/d}}$$

$T \sim t_d$.

6.39 Example

Let $X_1, \dots, X_n \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(\mu, \sigma^2)$, $\mu \in \mathbb{R}$, $\sigma^2 > 0$, $n \geq 2$; $\hat{\sigma}_n := \sqrt{\hat{\sigma}_n^2}$. Then,

$$\sqrt{n-1} \frac{\bar{X}_n - \mu}{\hat{\sigma}_n} \sim t_{n-1}$$

6.4 Matringales

6.40 Definition [Conditional Expectation]

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, $X \in L^1(\Omega)$ a random variable, and $\mathcal{G} \subset \mathcal{F}$ a sub- σ -algebra. The **conditional expectation** of X given $\mathcal{G}$, denoted $\mathbb{E}[X \mid \mathcal{G}]$, is the unique (up to a.s. equality) $\mathcal{G}$ -measurable random variable satisfying

$$\int_G \mathbb{E}[X \mid \mathcal{G}] d\mathbb{P} = \int_G X d\mathbb{P}, \quad \forall G \in \mathcal{G}.$$

6.41 Definition [Martingale]

Let $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$ be a filtered probability space. A stochastic process $(M_t)_{t \geq 0}$ adapted to $(\mathcal{F}_t)$ is a **martingale** if

$$\mathbb{E}[|M_t|] < \infty, \quad \text{and} \quad \mathbb{E}[M_t \mid \mathcal{F}_s] = M_s \quad \text{a.s. for all } 0 \leq s \leq t.$$

6.5 Stochastic Calculus

6.42 Definition [Brownian Motion]