** Collision Induced Absorption (CIA) references**

These numbers can be found in the data headers of the HITRAN CIA files, see readme file for details

- [1] K.M. Smith, D.A. Newnham, Near-infrared absorption cross sections and integrated absorption intensities of molecular oxygen (O₂, O₂ -O₂, and O₂ -N₂), J. Geophys. Res. Atmos. 105 (2000) 7383–7396. doi:10.1029/1999JD901171.
- [2] Y.I. Baranov, W.J. Lafferty, G.T. Fraser, Investigation of collision-induced absorption in the vibrational fundamental bands of O_2 and N_2 at elevated temperatures, J. Mol. Spectrosc. 233 (2005) 160–163. doi:10.1016/j.jms.2005.06.008.
- [3] W.J. Lafferty, A.M. Solodov, A. Weber, W.B. Olson, J.-M. Hartmann, Infrared collision-induced absorption by N_2 near 4.3 μ m for atmospheric applications: measurements and empirical modeling, Appl. Opt. 35 (1996) 5911. doi:10.1364/AO.35.005911.
- [4] H. Tran, C. Boulet, J.-M. Hartmann, Line mixing and collision-induced absorption by oxygen in the A band: Laboratory measurements, model, and tools for atmospheric spectra computations, J. Geophys. Res. 111 (2006) D15210. doi:10.1029/2005JD006869.
- [5] M. Vangvichith, H. Tran, J.-M. Hartmann, Line-mixing and collision induced absorption for O₂–CO₂ mixtures in the oxygen A-band region, J. Quant. Spectrosc. Radiat. Transf. 110 (2009) 2212–2216. doi:10.1016/J.JQSRT.2009.06.002.
- [6] M. Abel, L. Frommhold, X. Li, K.L.C. Hunt, Infrared absorption by collisional H_2 –He complexes at temperatures up to 9000 K and frequencies from 0 to 20 000 cm⁻¹, J. Chem. Phys. 136 (2012) 044319. doi:10.1063/1.3676405.
- [7] M. Abel, L. Frommhold, X. Li, K.L.C. Hunt, Collision-Induced Absorption by H_2 Pairs: From Hundreds to Thousands of Kelvin, J. Phys. Chem. A. 115 (2011) 6805–6812. doi:10.1021/jp109441f.
- [8] G.D. Greenblatt, J.J. Orlando, J.B. Burkholder, A.R. Ravishankara, Absorption measurements of oxygen between 330 and 1140 nm, J. Geophys. Res. 95 (1990) 18577. doi:10.1029/JD095iD11p18577.
- [9] C. Hermans, Measurement of absorption cross sections and spectroscopic molecular parameters, (2011).http://spectrolab.aeronomie.be/o2.htm.
- [10] B. Maté, C. Lugez, G.T. Fraser, W.J. Lafferty, Absolute intensities for the O_2 1.27 μ m continuum absorption, J. Geophys. Res. Atmos. 104 (1999) 30585–30590. doi:10.1029/1999JD900824.
- [11] A. Borysow, L. Frommhold, Collision-induced rototranslational absorption spectra of N_2 - N_2 pairs for temperatures from 50 to 300 K, Astrophys. J. 311 (1986) 1043. doi: 10.1086/164841.
- [12] A. Borysow, L. Frommhold, Theoretical collision-induced rototranslational absorption spectra for modeling Titan's atmosphere H₂-N₂ pairs, Astrophys. J. 303 (1986) 495. doi:10.1086/164096.
- [13] A. Borysow, C. Tang, Far Infrared CIA Spectra of N₂-CH₄ Pairs for Modeling of Titan's Atmosphere, Icarus. 105 (1993) 175–183. doi:10.1006/icar.1993.1117.

- [14] J. Borysow, L. Frommhold, G. Birnbaum, Collison-induced rototranslational absorption spectra of H_2 -He pairs at temperatures from 40 to 3000 K, Astrophys. J. 326 (1988) 509. doi:10.1086/166112.
- [15] G. Orton, M. Gustafsson, M. Burgdorf, V. Meadows, Revised ab initio models for H₂–H₂ collision-induced absorption at low temperatures, Icarus. 189 (2007) 544–549. doi:10.1016/j.icarus.2007.02.003.
- [16] A. Borysow, L. Frommhold, Theoretical collision-induced rototranslational absorption spectra for the outer planets H₂-CH₄ pairs, Astrophys. J. 304 (1986) 849. doi:10.1086/164221.
- [17] M. Gustafsson, L. Frommhold, The H_2 –H infrared absorption bands at temperatures from 1000 K to 2500 K, Astron. Astrophys. 400 (2003) 1161–1162. doi:10.1051/0004-6361:20030100.
- [18] M. Gustafsson, L. Frommhold, Infrared Absorption Spectra of Collisionally Interacting He and H Atoms, Astrophys. J. 546 (2001) 1168–1170. doi:10.1086/318311.
- [19] A. Borysow, L. Frommhold, Collision-induced rototranslational absorption spectra of CH₄-CH₄ pairs at temperatures from 50 to 300 K, Astrophys. J. 318 (1987) 940. doi:10.1086/165426.
- [20] M. Gruszka, A. Borysow, Roto-Translational Collision-Induced Absorption of CO₂ for the Atmosphere of Venus at Frequencies from 0 to 250 cm⁻¹, at Temperatures from 200 to 800 K, Icarus. 129 (1997) 172−177. doi:10.1006/icar.1997.5773.
- [21] R.E. Samuelson, N.R. Nath, A. Borysow, Gaseous abundances and methane supersaturation in Titan's troposphere, Planet. Space Sci. 45 (1997) 959–980. doi:10.1016/S0032-0633(97)00090-1.
- [22] B.J. Drouin, D.C. Benner, L.R. Brown, M.J. Cich, T.J. Crawford, V.M. Devi, A. Guillaume, J.T. Hodges, E.J. Mlawer, D.J. Robichaud, F. Oyafuso, V.H. Payne, K. Sung, E.H. Wishnow, S. Yu, Multispectrum analysis of the oxygen A-band, J. Quant. Spectrosc. Radiat. Transf. 186 (2017) 118–138. doi:10.1016/j.jqsrt.2016.03.037.
- [23] Y.I. Baranov, W.J. Lafferty, G.T. Fraser, Infrared spectrum of the continuum and dimer absorption in the vicinity of the O_2 vibrational fundamental in O_2/CO_2 mixtures, J. Mol. Spectrosc. 228 (2004) 432–440. doi:10.1016/j.jms.2004.04.010.
- [24] J.M. Hartmann, C. Boulet, G.C. Toon, Collision-induced absorption by N₂ near 2.16μm: Calculations, model, and consequences for atmospheric remote sensing, J. Geophys. Res. 122 (2017) 2419–2428. doi:10.1002/2016JD025677.
- [25] R.H. Taylor, A. Borysow, L. Frommhold, Concerning the rototranslational absorption spectra of He-CH₄ pairs, J. Mol. Spectrosc. 129 (1988) 45–58. doi:10.1016/0022-2852(88)90257-3.
- [26] E. Bar-Ziv, S. Weiss, Translational Spectra Due to Collision-Induced Overlap Moments in Mixtures of He with CO₂, N₂, CH₄, and C₂H₆, J. Chem. Phys. 57 (1972) 34–37. doi:10.1063/1.1677970.
- [27] T. Karman, M.A.J. Koenis, A. Banerjee, D.H. Parker, I.E. Gordon, A. Van Der Avoird, W.J. Van Der Zande, G.C. Groenenboom, O₂ –O₂ and O₂ –N₂ collision-induced absorption mechanisms unravelled, Nat. Chem. (2018) 1–6. doi:10.1038/s41557-018-0015-x.

- [28] R. Wordsworth, Y. Kalugina, S. Lokshtanov, A. Vigasin, B. Ehlmann, J. Head, C. Sanders, H. Wang, Transient reducing greenhouse warming on early Mars, Geophys. Res. Lett. 44 (2017) 665–671. doi:10.1002/2016GL071766.
- [29] F.R. Spiering, W.J. van der Zande, Collision induced absorption in the $a^1\Delta(v=2) \leftarrow X^1\Sigma_g^-(v=0)$ band of molecular oxygen, Phys. Chem. Chem. Phys. 14 (2012) 9923. doi:10.1039/c2cp40961e.
- [30] F.R. Spiering, M.B. Kiseleva, N.N. Filippov, H. Naus, B. van Lieshout, C. Weijenborg, W.J. van der Zande, Line mixing and collision induced absorption in the oxygen A-band using cavity ring-down spectroscopy., J. Chem. Phys. 133 (2010) 114305. doi:10.1063/1.3460924.
- [31] T. Karman, E. Miliordos, K.L.C. Hunt, G.C. Groenenboom, A. van der Avoird, Quantum mechanical calculation of the collision-induced absorption spectra of N₂ −N₂ with anisotropic interactions, J. Chem. Phys. 142 (2015) 084306. doi:10.1063/1.4907917.
- [32] R. Thalman, R. Volkamer, Temperature dependent absorption cross-sections of O₂–O₂ collision pairs between 340 and 630 nm and at atmospherically relevant pressure, Phys. Chem. Chem. Phys. 15 (2013) 15371. doi:10.1039/c3cp50968k.
- [33] Estimated from N_2 - N_2 data from Refs. [2,3] using Eq. (7) of Ref. [3], which is derived using data from V. Menoux, R.L. Doucen, C. Boulet, A. Roblin, A.M. Bouchardy, Collision-induced absorption in the fundamental band of N_2 : temperature dependence of the absorption for N_2 - N_2 and N_2 - N_2 pairs., Appl. Opt. 32 (1993) 263–268. doi: 10.1364/AO.32.000263.
- [34] L.N. Fletcher, M. Gustafsson, G.S. Orton, Hydrogen Dimers in Giant-planet Infrared Spectra, Astrophys. J. Suppl. Ser. 235 (2018) 24. doi:10.3847/1538-4365/aaa07a.
- [35] J.-M. Hartmann, C. Boulet, D.D. Tran, H. Tran, Y. Baranov, Effect of humidity on the absorption continua of CO_2 and N_2 near 4 μ m: Calculations, comparisons with measurements, and consequences for atmospheric spectra, J. Chem. Phys. 148 (2018) 054304. doi:10.1063/1.5019994.
- [36] E.H. Wishnow, K. Sung, The far-infrared collision-induced spectrum of nitrogen over the temperature range 78-129 K, J. Quant. Spectrosc. Radiat. Transf. in prepara (2018).
- [37] Y.I. Baranov, A.A. Vigasin, Collision-Induced Absorption by CO_2 in the Region of v_1 , $2v_2$, J. Mol. Spectrosc. 193 (1999) 319–325. doi: 10.1006/JMSP.1998.7743.
- [38] Y.I. Baranov, G.T. Fraser, W.J. Lafferty, A.A. Vigasin, Collision-Induced Absorption in The CO₂ Fermi Triad for Temperatures from 211K to 296 K, in: Weakly Interact. Mol. Pairs Unconv. Absorbers Radiat. Atmos., Springer Netherlands, Dordrecht, 2003: pp. 149–158. doi:10.1007/978-94-010-0025-3 12.
- [39] Y.I. Baranov, Collision-induced absorption in the region of the $v_2 + v_3$ band of carbon dioxide, J. Mol. Spectrosc. 345 (2018) 11–16. doi:10.1016/J.JMS.2017.11.005.