

# Nonlinearity

## Numerical Flow Simulation

# Linearity

- General conservation equation

$$\frac{\partial(\rho\phi)}{\partial t} + \text{div}(\rho\phi\mathbf{u}) = \text{div}(\Gamma \text{grad}(\phi)) + S$$

- Linear in  $\phi$  if:

- coefficients independent of  $\phi$ ,
- source term independent of/linear in  $\phi$ .

- Example: steady diffusion  $\frac{\partial}{\partial x} \left( k \frac{\partial T}{\partial x} \right) + S = 0$  with  $\left| \begin{array}{l} k = \text{cst} \text{ or } k(x) \\ S = \text{cst} \text{ or } S = S(x) \\ \text{or } S = S_c + S_l T \end{array} \right.$

- Discretization yields a **linear** algebraic system:

$$\mathbf{A}\phi = \mathbf{b} \quad \rightarrow \quad \phi = \mathbf{A}^{-1}\mathbf{b}$$

# Nonlinearity

- General conservation equation

$$\frac{\partial(\rho\phi)}{\partial t} + \text{div}(\rho\phi\mathbf{u}) = \text{div}(\Gamma \text{grad}(\phi)) + S$$

- Nonlinear in  $\phi$  if:
  - Coefficients depend on  $\phi$
  - Other terms nonlinear in  $\phi$

- Example: Navier-Stokes equations

$$\frac{\partial \rho}{\partial t} + \boxed{\text{div}(\rho\mathbf{u})} = 0$$

Nonlinear if compressible  
(both  $\rho$  and  $\mathbf{u}$  unknown)

Always nonlinear


$$\boxed{\frac{\partial(\rho\mathbf{u})}{\partial t}} + \boxed{\text{div}(\rho\mathbf{u}\mathbf{u})} = \text{div} \left[ \left( -p - \frac{2}{3}\mu \text{div}(\mathbf{u}) \right) \mathbf{I} + 2\mu\mathbf{d} \right]$$

- Discretization yield a **nonlinear** system:  $\mathbf{A}(\phi) = \mathbf{b} \rightarrow \phi = ?$
- We only know how to solve linear systems  $\rightarrow$  must linearize.


# Linearization: Picard's method

$$\mathbf{A}(\phi) = \mathbf{b} \rightarrow \phi = ?$$

- If the system can be written in the “quasi-linear” form  $\tilde{\mathbf{A}}(\phi)\phi = \mathbf{b}$ 
  - take a guess solution  $\phi^*$  to evaluate the matrix  $\tilde{\mathbf{A}}(\phi^*)$
  - solve  $\tilde{\mathbf{A}}(\phi^*)\phi = \mathbf{b}$
  - take  $\phi$  as new guess,
  - iterate until convergence.



Sketch: in the scalar case,  $\tilde{A}(\phi^*)\phi$  is a line that goes through the 2 points  $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$  and  $\begin{pmatrix} \phi^* \\ \tilde{A}(\phi^*)\phi^* \end{pmatrix} = \begin{pmatrix} \phi^* \\ A(\phi^*) \end{pmatrix}$ , which defines a 3<sup>rd</sup> point  $\begin{pmatrix} \phi_{new}^* \\ b \end{pmatrix}$ .




# Linearization: Newton's method

$$\mathbf{A}(\phi) = \mathbf{b} \quad \rightarrow \quad \phi = ?$$

- Linearize with 1<sup>st</sup>-order Taylor expansion:

$$\mathbf{A}(\phi^* + \delta\phi) \approx \mathbf{A}(\phi^*) + \left. \frac{\partial \mathbf{A}}{\partial \phi} \right|_{\phi^*} \delta\phi \approx \mathbf{b}$$


- take a guess solution to evaluate the Jacobian matrix  $\mathbf{J}(\phi^*) = \left. \frac{\partial \mathbf{A}}{\partial \phi} \right|_{\phi^*}$
- solve for the increment  $\delta\phi$
- take  $\phi = \phi^* + \delta\phi$  as new guess,
- iterate until convergence.



- Alternatively, can write  $\mathbf{A}(\phi^*) + \mathbf{J}(\phi^*)(\phi - \phi^*) \approx \mathbf{b}$  and solve directly for  $\phi$ .
- Faster convergence than Picard's method, but more complicated to implement and costly to solve.

# Possible problems

- May fail:
  - Can converge to “another” solution (when multiple solutions)
  - Can oscillate / diverge
  
- Remedies:
  - Choose a good initial guess
  - Use under-relaxation



# Under-relaxation

- Instead of taking  $\phi$  or  $\phi^* + \delta\phi$  as new guess,  
take  $\omega\phi + (1 - \omega)\phi^*$  or  $\phi^* + \omega\delta\phi$ , with  $\omega < 1$
- Can help to stabilize the iterative process.
- Generally, slows down convergence.
- Suitable value of the relaxation factor is case-dependent.



# Examples

- Convective term (momentum eq.)  $\mathbf{C}(\phi) = \rho \mathbf{u} \mathbf{u}$

Picard

$$\mathbf{C}(\phi) \approx (\rho^* \mathbf{u}^*) \mathbf{u}$$

Newton

$$\mathbf{C}(\phi^* + \delta \phi) \approx \rho^* \mathbf{u}^* \mathbf{u}^* + (\mathbf{u}^* \mathbf{u}^*) \delta \rho + (2\rho^* \mathbf{u}^*) \delta \mathbf{u}$$

$$\text{or} \quad -2\rho^* \mathbf{u}^* \mathbf{u}^* + (\mathbf{u}^* \mathbf{u}^*) \rho + (2\rho^* \mathbf{u}^*) \mathbf{u}$$

- Nonlinear source term (steady diffusion eq.), e.g.  $S(T) = 4 - 5T^3$

Picard

$$S(T) \approx 4 - (5T^{*2})T$$

Newton


$$S(T^* + \delta T) \approx 4 - 5T^{*3} - (15T^{*2})\delta T$$

$$\text{or} \quad 4 + 10T^{*3} - (15T^{*2})T$$




# Summary

Linear system,  
direct methods



Linear system,  
iterative methods



Nonlinear system,  
iterative methods

