

Abdelouahed Hamdi

## Convergence Results on Proximal Method of Multipliers in Nonconvex Programming

Bericht des Journal of Public Health

### Kurzfassung

**Aim:** The aim of this study was to examine whether the correlates of physical activity (PA) differed as a function of two different PA recommendations: (1) the Fitness Recommendation, i.e., the improvement of cardiovascular fitness by  $3 \times 20$  min/week of vigorously intense PA and (2) the health-enhancing PA (HEPA) concept such as the emphasis on the important benefits to health and fitness by performing  $5 \times 60$  min/week of moderately intense PA. **Subjects and Methods:** A cross-sectional design was used. In a sample of 524 adolescents (50.2% males, mean age = 15.0), we quantified different (demographic/biological, psychological, behavioural) self-reported correlates of PA as well as compliance with the PA recommendations. Sex-specific multivariate logistic regression models were used to identify correlates of fulfilling the two PA recommendations. **Results:** Multivariate analyses revealed differences between the correlates and PA-specific Recommendations. For boys, only the Fitness Recommendation was significantly associated with psychosocial variables (self-efficacy and behavioural change strategies) and membership in a sports club. For girls, membership in a sports club was significantly associated with both PA recommendations. However, this association was positive with respect to the Fitness Recommendation and negative with respect to the HEPA Recommendation. **Conclusion:** Our results provided some preliminary indication that correlates differ depending on specific recommendations in 14–16-year-old boys and girls. Moreover some variability with regard to sex was observed.