

A mathematical modelling of rheumatoid arthritis

F. Apparailly¹, P. Azerad², O. Bortolotti¹, G. Courties¹, O. Iosifescu², A. Parmeggiani³

¹IRMB, UMR1183, Université de Montpellier and INSERM, France

²IMAG, UMR5149, Université de Montpellier, France

³L2C, UMR5221, Université de Montpellier, France

October 10, 2025

We model the interaction between three populations of cells in the context of arthritis, namely the neutrophils, the tissue resident macrophages (SubLining Tissue Resident Macrophages) and the Monocyte Macrophages. The SL-TRM macrophage cells reside in the outer border of the joint. The Momac (monocytes-macrophages) cells are coming from the reservoir of monocytes in the marrow. Basically the crisis scenario is the following: The arthritis crisis is triggered by an external signal (injection of immune serum containing antigens) that activates the immune system and provokes an inflammatory response. The antigens are recognized by the macrophages L-TRM (lining tissue resident macrophages) cells that secrete inflammatory chemokines and cytokines. These cytokines activate the fibroblasts which in turn recruit the neutrophils cells from the blood stream into the joint. The cytokines also recruit the Momac cells from the blood stream into the joint. There is a positive feedback since the inflammation is amplified by the neutrophils that are recruited in the joint by the chemokines secreted by the Momac cells. The macrophages SL-TRM cells have a protective role since they inhibit the inflammation but they are depleted during the crisis because they are penalized by the proliferation of Momac cells.

1 Model

Basically the crisis scenario is the following: The arthritis crisis is triggered by an external signal (injection of immune serum containing antigens) that activates the immune system and provokes an inflammatory response. The antigens are recognized by the macrophages L-TRM (lining tissue resident macrophages) cells that secrete inflammatory chemokines and cytokines. These cytokines activate the fibroblasts which in turn recruit the neutrophils cells from the blood stream into the joint. The cytokines also recruit the Momac cells from the blood stream into the joint. There is a positive feedback since the inflammation is amplified by the neutrophils that are recruited in the joint by the chemokines secreted by the Momac cells. The macrophages SL-TRM cells have a protective role since they inhibit the inflammation but they are depleted during the crisis because they are inhibited by the proliferation of Momac cells.

1.1 Mathematical Model

Let us denote by $T(t)$ the population of SL-TRM cells and by $M(t)$ the population of Momac cells at time t . The cells T (SL-TRM) and M (Momac) interact with each other in the following way. The Momac cells M are recruited from the blood stream at a constant rate a . They die at a rate δ_M and they transform into SL-TRM cells at a rate b . The SL-TRM cells T die at a rate δ_T . The interaction between the two populations is described by the following system of differential equations

The inflammatory state $S(t)$ is enhanced by neutrophils N and inhibited by SL-TRM macrophages T .

Inflammation further increases neutrophils (factor α), increases Momac (factor β), and decreases SL-TRM (factor γ).

Notice that we do not keep track of the population of fibroblasts, but rather the population of neutrophils, which are recruited by inflammatory chemokines.

The variable $N(t)$ designates the population of neutrophils at time t . Finally $S(t)$ is the inflammatory signal at time t . The model is given by the following system of ordinary differential equations:

$$\begin{cases} \frac{dS}{dt} = (a N - b T)S \\ \frac{dN}{dt} = \alpha S N - \delta_N N(N - 1) \\ \frac{dM}{dt} = \beta S - \delta_M M(M - 1) \\ \frac{dT}{dt} = -\gamma S T + \delta_M M(1 - T) \end{cases} \quad (1)$$

where $a, b, \alpha, \beta, \gamma, \delta_N, \delta_M$ are positive parameters. To make things easier to understand, we summarize the model in the following diagram (see Figure 1.1).

Remark (Comments and questions on the model):

- to be discussed with our immunologists collaborators

2 Analysis of the model

2.1 Equilibria points

Equilibria points of the system are found by setting the right-hand side of the system to zero, i.e. by solving the following system of equations: 01 The equilibrium points (S^*, N^*, M^*, T^*) satisfy the following system:

$$(aN^* - bT^*)S^* = 0 \quad (2)$$

$$\alpha S^* N^* - \delta_N N^*(N^* - 1) = 0 \quad (3)$$

$$\beta S^* - \delta_M M^*(M^* - 1) = 0 \quad (4)$$

$$-\gamma S^* T^* + \delta_M M^*(1 - T^*) = 0 \quad (5)$$

This gives the following equilibria points. A first natural equilibrium point is given by $S^* = 0, N^* = 1, M^* = 1, T^* = 1$. This point corresponds to the disease-free state, where there is no inflammation ($S^* = 0$) and the populations of neutrophils, Momac,



Figure 1: Schematic representation of the model

and SL-TRM are at their baseline levels ($N^* = 1, M^* = 1, T^* = 1$). There are other equilibria points with $S^* = 0$ namely $S^* = 0, N^* = 0, M^* = 0, T^* = t$, with arbitrary t ; $S^* = 0, N^* = 0, M^* = 1, T^* = 1$; $S^* = 0, N^* = 1, M^* = 0, T^* = t$, with arbitrary t .

For equilibria points with $S^* \neq 0$, from equation (2) we have $aN^* - bT^* = 0$, hence

$$T^* = \frac{a}{b}N^*. \quad (6)$$

Factoring N^* from equation (3), we obtain:

$$N^* (\alpha S^* - \delta_N(N^* - 1)) = 0$$

This gives us two possibilities: either $N^* = 0$ or $\alpha S^* - \delta_N(N^* - 1) = 0$. If $N^* = 0$, then from equation (2) we have $a \cdot 0 - bT^* = 0$, hence $T^* = 0$. From equation (5), we get $M^* = 0$. Then from equation (4), we have $\beta S^* - \delta_M \cdot 0 = 0$, hence $S^* = 0$. This contradicts our assumption that $S^* \neq 0$. Therefore, the case $N^* = 0$ is not possible for equilibria with $S^* \neq 0$. Therefore $\alpha S^* - \delta_N(N^* - 1) = 0$, so we can express S^* in terms of N^* :

$$S^* = \frac{\delta_N}{\alpha}(N^* - 1) \quad (7)$$

We substitute (6)-(7) into equation (4) and (5) and obtain a 2 by 2 system for M^* and N^* : By substituting $T^* = \frac{a}{b}N^*$ and $S^* = \frac{\delta_N}{\alpha}(N^* - 1)$ into the equilibrium equations for

M^* and T^* , we obtain the following system for N^* and M^* :

$$\begin{cases} \beta \frac{\delta_N}{\alpha} (N^* - 1) - \delta_M M^* (M^* - 1) = 0 \\ -\gamma \frac{\delta_N}{\alpha} (N^* - 1) \cdot \frac{a}{b} N^* + \delta_M M^* \left(1 - \frac{a}{b} N^*\right) = 0 \end{cases}$$

This system allows us to determine the equilibrium values N^* and M^* when $S^* \neq 0$. Indeed we can express N^* in terms of M^* from the first equation:

$$N^* = 1 + \frac{\alpha \delta_M}{\beta \delta_N} M^* (M^* - 1) \quad (8)$$

Substituting this expression for N^* into the second equation, we obtain an equation for M^* :

$$M^* = \frac{a \gamma \delta_N}{b \alpha \delta_M} \frac{N^* (N^* - 1)}{(1 - \frac{a}{b} N^*)} \quad (9)$$

Geometrically, the equilibria points with $S^* \neq 0$ correspond to the intersection points of the parabola defined by equations (8) and the hyperbola defined by equation (9) in the (N^*, M^*) plane. The number of such intersection points (and hence the number of equilibria with $S^* \neq 0$) depends on the parameters of the model. We can find the asymptotes of the hyperbola and the apex of the parabola to determine the possible number of intersection points in the first quadrant. We rewrite the equation (9) of the hyperbola as follows:

$$M^* = -\frac{\gamma \delta_N}{\alpha \delta_M} (N^* + b/a - 1) + \frac{(b/a)^2 (1 - b/a)}{(N^* - b/a)} \quad (10)$$

From this expression, we see that the hyperbola has a vertical asymptote at $N^* = b/a$ and an oblique asymptote $M^* = -\frac{\gamma \delta_N}{\alpha \delta_M} (N^* + b/a - 1)$. The parabola (8) is horizontal in the N, M plane, has its apex at $N_{min} = 1 - \frac{\alpha \delta_M}{4\beta \delta_N}$ and $M = 1/2$ and is symmetric with respect to the horizontal line $M = 1/2$ and opens eastwards. We plot some typical cases in the following figure. Notice that we need only 4 *non dimensional* parameters a/b , α/β , γ/α , and δ_N/δ_M instead of the 7 original parameters.

There is a smarter way¹ to find the number of equilibria points with $S^* \neq 0$ leading to a cubic equation in T^* . We express N^* in terms of T^* from equation (6):

$$N^* = \frac{b}{a} T^* \quad (11)$$

Substituting this expression for N^* into equation (7), we obtain:

$$S^* = \frac{\delta_N}{\alpha} \left(\frac{b}{a} T^* - 1 \right) \quad (12)$$

From equation (4) and equation (5), we derive $\delta_M M^* = \frac{\beta S^*}{M^* - 1}$ and $\delta_M M^* = \frac{\gamma S^* T^*}{1 - T^*}$. Equating these two expressions for $\delta_M M^*$, we obtain:

$$\frac{\beta S^*}{M^* - 1} = \frac{\gamma S^* T^*}{1 - T^*}$$

¹thanks to O. Iosifescu



Figure 2: equilibria in the (N^*, M^*) plane, left $a < b$, right $a > b$.

Assuming $S^* \neq 0$, we can simplify this equation to:

$$\frac{\beta}{M^* - 1} = \frac{\gamma T^*}{1 - T^*}$$

From this, we can express M^* in terms of T^* :

$$M^* = 1 + \frac{\beta(1 - T^*)}{\gamma T^*} = 1 - \frac{\beta}{\gamma} + \frac{\beta}{\gamma T^*} \quad (13)$$

Substituting the expression for M^* and S^* into the last equation (5), we obtain

$$-\gamma \frac{\delta_N}{\alpha} \left(\frac{b}{a} T^* - 1 \right) T^* + \delta_M \left(1 - \frac{\beta}{\gamma} + \frac{\beta}{\gamma T^*} \right) (1 - T^*) = 0 \quad (14)$$

Since $T^* \neq 0$, we can multiply this equation by T^* and rearrange the terms to obtain a cubic equation for T^* :

$$\frac{\gamma}{\alpha} \frac{\delta_N}{\delta_M} \frac{b}{a} (T^*)^3 + \left(1 - \frac{\gamma}{\alpha} \frac{\delta_N}{\delta_M} - \frac{\beta}{\gamma} \right) (T^*)^2 + \left(2 \frac{\beta}{\gamma} - 1 \right) T^* - \frac{\beta}{\gamma} = 0 \quad (15)$$

Rewriting this equation in the standard unitary form $(T^*)^3 + p(T^*)^2 + qT^* + r = 0$, we have:

$$(T^*)^3 + \frac{\alpha}{\gamma} \frac{\delta_M}{\delta_N} \frac{a}{b} \left(1 - \frac{\beta}{\gamma} - \frac{\gamma}{\alpha} \frac{\delta_N}{\delta_M} \right) (T^*)^2 + \frac{\alpha}{\gamma} \frac{\delta_M}{\delta_N} \frac{a}{b} \left(2 \frac{\beta}{\gamma} - 1 \right) T^* - \frac{\alpha}{\gamma} \frac{\delta_M}{\delta_N} \frac{a}{b} \frac{\beta}{\gamma} = 0 \quad (16)$$

Notice that the coefficients of this cubic equation depend only on the 4 *non dimensional* parameters a/b , α/β , γ/α , and δ_N/δ_M . Denoting these dimensionless parameters by $A = a/b$, $B = \beta/\gamma$, $C = \alpha/\gamma$, and $D = \delta_M/\delta_N$, we can rewrite the cubic equation as follows:

$$(T^*)^3 + ACD \left(1 - B - \frac{1}{CD}\right) (T^*)^2 + ACD(2B - 1)T^* - ABCD = 0 \quad (17)$$

Remark 2.1 *There is an obvious solution in the particular case $A = a/b = 1$, namely $T^* = 1$. This solution corresponds to the disease-free equilibrium point $S^* = 0, N^* = 1, M^* = 1, T^* = 1$. But this solution is not relevant for the case $S^* \neq 0$ that we are considering here.*

From basic algebra, we can show that there are at most 3 such intersection points, hence at most 3 equilibria points with $S^* \neq 0$. We can be more precise because the biological feasibility imposes the roots to be real nonnegative numbers. Since $ABCD > 0$, there is at least one positive root. But we can be more precise using *Descartes' rule of signs*. It states that the number of positive real roots of a polynomial (counting multiplicities) is equal to the number of sign changes between consecutive nonzero coefficients, or less than that by an even number. In our case, the coefficients of the cubic polynomial in (17) are:

$$1, \quad ACD \left(1 - B - \frac{1}{CD}\right), \quad ACD(2B - 1), \quad -ABCD$$

So the number of sign changes depends on the signs of the second and third coefficients. We have the following cases:

- If $B + \frac{1}{CD} > 1$ and $B > 1/2$, then there are 3 sign changes and the cubic has one or three positive roots.
- If $B + \frac{1}{CD} < 1$ and $B > 1/2$, then there is one sign change and the cubic has only one positive roots.
- If $B + \frac{1}{CD} < 1$ and $B < 1/2$, then there is one sign changes and the cubic has only one positive root.
- If $B + \frac{1}{CD} > 1$ and $B < 1/2$, then there is one sign change and the cubic has only one positive root.

We can plot some typical cases in the following figure (see Figure 3).

We can use Cardano's method to find the roots of the cubic equation (17). We rewrite it in the standard form:

$$T^3 + pT^2 + qT + r = 0$$

where

$$p = ACD \left(1 - B - \frac{1}{CD}\right), \quad q = ACD(2B - 1), \quad r = -ABCD$$

We perform the change of variable $T = x - p/3$ to eliminate the quadratic term and obtain the depressed cubic equation:

$$x^3 + \mathcal{A}x + \mathcal{B} = 0$$




Figure 3: Cubic equation in T^* , left $B < 1/2$, right $B > 1/2$.

where

$$\mathcal{A} = q - \frac{p^2}{3}, \quad \mathcal{B} = \frac{2p^3}{27} - \frac{pq}{3} + r$$

The discriminant of this equation is given by:

$$\Delta = \left(\frac{\mathcal{B}}{2}\right)^2 + \left(\frac{\mathcal{A}}{3}\right)^3$$

The nature of the roots depends on the sign of the discriminant Δ :

- If $\Delta > 0$, there is one real root and two complex conjugate roots.
- If $\Delta = 0$, all roots are real and at least two are equal.
- If $\Delta < 0$, all roots are real and unequal.

Hence we can characterize the case when we have three distinct real roots by the condition $\Delta < 0$, which is equivalent to:

$$\left(\frac{\mathcal{A}}{3}\right)^3 < -\left(\frac{\mathcal{B}}{2}\right)^2$$

This condition can be rewritten in terms of the original parameters using computer algebra as follows:

$$\left(\frac{\mathcal{A}}{3}\right)^3 < -\left(\frac{\mathcal{B}}{2}\right)^2$$

where

$$\mathcal{A} = A \left(-A(CD(B-1) + 1)^2 + 3CD(2B-1) \right) / 3$$

$$\mathcal{B} = A \left(-2A^2(CD(B-1)+1)^3 + 9ACD(2B-1)(CD(B-1)+1) - 27BCD \right) / 27$$

$$\Delta = \frac{A^2}{2916} \left(-4A \left[A(CD(B-1)+1)^2 - 3CD(2B-1) \right]^3 + \left[2A^2(CD(B-1)+1)^3 - 9ACD(2B-1)(CD(B-1)+1) + 27BCD \right]^2 \right)$$

The real roots can be found using Cardano's formulas². If $\Delta \geq 0$, the real root is given by:

$$x = \sqrt[3]{-\frac{\mathcal{B}}{2} + \sqrt{\Delta}} + \sqrt[3]{-\frac{\mathcal{B}}{2} - \sqrt{\Delta}}$$

If $\Delta < 0$, the three real roots are given by:

$$x_k = 2\sqrt{-\frac{\mathcal{A}}{3}} \cos \left(\frac{1}{3} \arccos \left(\frac{3\mathcal{B}}{2\mathcal{A}} \sqrt{-\frac{3}{\mathcal{A}}} \right) - \frac{2k\pi}{3} \right), \quad k = 0, 1, 2$$

Here we plot a case with three distinct positive roots (see Figure 4). Namely we take $A = 0.9$, $B = 7$, $C = 1.1$, and $D = 1.1$.




Figure 4: Cubic equation with three distinct real roots.

Finally, we can recover the roots T^* of the original cubic equation using the relation $T^* = x + p/3$.

2.2 Stability analysis

Stability analysis of the equilibria point (S^*, N^*, M^*, T^*) is done by linearizing the system around this point. The Jacobian matrix of the system is given by:

$$J = \begin{pmatrix} aN - bT & aS & 0 & -bS \\ \alpha N & \alpha S - \delta_N(2N - 1) & 0 & 0 \\ \beta & 0 & -\delta_M(2M - 1) & 0 \\ -\gamma T & 0 & \delta_M(1 - T) & -\gamma S - \delta_M M \end{pmatrix}$$

²probably wrong

We analyze first the stability of the disease-free equilibrium point $(S^*, N^*, M^*, T^*) = (0, 1, 1, 1)$. At this point, the Jacobian matrix simplifies to:

$$J|_{(S^*, N^*, M^*, T^*)=(0,1,1,1)} = \begin{pmatrix} a-b & 0 & 0 & 0 \\ \alpha & -\delta_N & 0 & 0 \\ \beta & 0 & -\delta_M & 0 \\ -\gamma & 0 & 0 & -\delta_M \end{pmatrix}$$

We notice immediately that the eigenvalues of this matrix are $a-b$, $-\delta_N$ and $-\delta_M$. The last two are negative since $\delta_N, \delta_M > 0$. The remaining eigenvalue is $a-b$. Thus the disease-free equilibrium is stable if $a < b$ and unstable if $a > b$. This means that if the inflammatory effect of Neutrophils cells controlled by parameter a is less than the curing effect of the SL-TRM macrophages controlled by parameter b , the system will return to the disease-free state. Conversely, if $a > b$, the system may move away from this equilibrium, potentially leading to a chronic inflammatory state. The stability of other equilibria points can be analyzed similarly by evaluating the Jacobian matrix at those points and determining the eigenvalues.

For $S^* = 0, N^* = 0, M^* = 0, T^* = t$, with arbitrary t , the Jacobian matrix becomes:

$$J|_{(S^*, N^*, M^*, T^*)=(0,0,0,t)} = \begin{pmatrix} -bt & 0 & 0 & 0 \\ 0 & \delta_N & 0 & 0 \\ \beta & 0 & \delta_M & 0 \\ -\gamma t & 0 & \delta_M(1-t) & 0 \end{pmatrix}$$

The eigenvalues of this matrix are $-bt$, δ_N , δ_M , and 0. Since δ_N and δ_M are positive, this equilibrium point is unstable. Even if the immune signal will decrease exponentially fast, the populations of neutrophils and momac will grow out of bound. This unstable equilibrium will not be observed in practice.

For $S^* = 0, N^* = 0, M^* = 1, T^* = 1$, the Jacobian matrix is:

$$J|_{(S^*, N^*, M^*, T^*)=(0,0,1,1)} = \begin{pmatrix} -b & 0 & 0 & 0 \\ 0 & \delta_N & 0 & 0 \\ \beta & 0 & -\delta_M & 0 \\ -\gamma & 0 & 0 & -\delta_M \end{pmatrix}$$

The eigenvalues of this matrix are $-b$, δ_N , $-\delta_M$, and $-\gamma$. Since there is a positive eigenvalue, this equilibrium point is unstable. Neutrophils can grow out of bound. This equilibrium will not be observed in practice neither.

For $S^* = 0, N^* = 1, M^* = 0, T^* = t$, with arbitrary t , the Jacobian matrix is

$$J|_{(S^*, N^*, M^*, T^*)=(0,1,0,t)} = \begin{pmatrix} -bt & 0 & 0 & 0 \\ \alpha & -\delta_N & 0 & 0 \\ \beta & 0 & \delta_M & 0 \\ -\gamma t & 0 & \delta_M(1-t) & 0 \end{pmatrix}$$

There is a positive eigenvalue δ_M , hence this equilibrium point is unstable. Momac can grow out of bound. This equilibrium will not be observed in practice neither.

3 Numerical simulations

Here we reproduce one typical simulations of the model using the following parameters: $\alpha = 1, \beta = 1, \gamma = 1, \delta_N = 1, \delta_M = 1, a = 0.35, b = 1$, duration=20 days and initial conditions $S(0) = 1, N(0) = 1, M(0) = 1, T(0) = 1$.




Figure 5:

We need to have realistic values for the parameters to feed the model.

The Python Jupyter notebook used to perform the simulations is available at <https://github.com/azerad/immunomath>.

3.1 Bifurcation analysis

We perform a bifurcation analysis with respect to the *non dimensional* parameters a/b , α/β , γ/α , and δ_N/δ_M . We vary a in the interval $[0, 1]$ and b in the interval $[0.1, 2]$. For each pair (a, b) , we simulate the system for 20 days and we record the value of S at the final time. The results are shown in the following heatmap.

Acknowledgments

This work was supported by the IMMUN4CURE project of IHU Montpellier.

References

- [1] Jit, M., et al. (2005). Mathematical modelling of rheumatoid arthritis. *Rheumatology*, 44, 323–331. <https://doi.org/10.1093/rheumatology/keh491>
- [2] Baker, M., et al. (2013). Mathematical modelling of cytokine-mediated inflammation in rheumatoid arthritis *Mathematical Medicine and Biology*, 30, 311–337. <https://doi.org/10.1093/imammb/dqs026>
- [3] Baker, M., et al. (2017). Mathematical modelling of cytokines, MMPs and fibronectin fragments in osteoarthritic cartilage *Journal of Mathematical Biology*, 75, 985–1024 (2017). <https://doi.org/10.1007/s00285-017-1104-y>

- [4] Baker, M., et al. (2018). Mathematical models of inflammation and its resolution. *Immunology Letters*, 196, 1–10. <https://doi.org/10.1016/j.imlet.2018.01.005>
- [5] Zhou, X., et al. (2018). Circuit Design Features of a Stable Two-Cell System. *Cell*, 172, 744–757. <https://doi.org/10.1016/j.cell.2018.01.015>
- [6] Odisharia, K., Odisharia, V., Tsereteli, P., and Janikashvili, N. (2019). On the Mathematical Model of Drug Treatment of Rheumatoid Arthritis In G. Jaiani and D. Natroshvili (eds.). Mathematics, Informatics, and their Applications in Natural Sciences and Engineering. *Springer Proceedings in Mathematics & Statistics*, 2019. https://doi.org/10.1007/978-3-030-10419-1_10
- [7] Handel, A., et al (2020). Simulation modelling for immunologists *Nature Reviews Immunology*, 20, 186–195. <https://doi.org/10.1038/s41577-019-0235-3>
- [8] Macfarlane, F. R., Chaplain, M. A. J., & Eftimie, R. (2022). Mathematical models of immune-mediated diseases: A review. *Mathematics*, 10(3), 456.
- [9] Ugolkov, Y., et al. (2024). Mathematical modeling in autoimmune diseases: from theory to clinical application *Frontiers in Immunology*, 15, 1371620. <https://doi.org/10.3389/fimmu.2024.1371620>