

Factors influencing patient engagement in mental health chatbots: A thematic analysis of findings from a systematic review of reviews

DIGITAL HEALTH
Volume 10: 1–10
© The Author(s) 2024
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/20552076241247983
journals.sagepub.com/home/dhj



Mohsen Khosravi¹ and Ghazaleh Azar²

Abstract

Introduction: Mental health disorders affect millions of people worldwide. Chatbots are a new technology that can help users with mental health issues by providing innovative features. This article aimed to conduct a systematic review of reviews on chatbots in mental health services and synthesized the evidence on the factors influencing patient engagement with chatbots.

Methods: This study reviewed the literature from 2000 to 2024 using qualitative analysis. The authors conducted a systematic search of several databases, such as PubMed, Scopus, ProQuest, and Cochrane database of systematic reviews, to identify relevant studies on the topic. The quality of the selected studies was assessed using the Critical Appraisal Skills Programme appraisal checklist and the data obtained from the systematic review were subjected to a thematic analysis utilizing the Boyatzis's code development approach.

Results: The database search resulted in 1494 papers, of which 10 were included in the study after the screening process. The quality assessment of the included studies scored the papers within a moderate level. The thematic analysis revealed four main themes: chatbot design, chatbot outcomes, user perceptions, and user characteristics.

Conclusion: The research proposed some ways to use color and music in chatbot design. It also provided a systematic and multidimensional analysis of the factors, offered some insights for chatbot developers and researchers, and highlighted the potential of chatbots to improve patient-centered and person-centered care in mental health services.

Keywords

Mental health, artificial intelligence, chatbots, patient engagement, patient-centered care

Submission date: 24 August 2023; Acceptance date: 29 March 2024

Introduction

The prevalence of mental health disorders is a significant challenge for society, as they affect a large number of individuals worldwide. As an example, according to the World Health Organization, depression alone affects more than 264 million people of all ages, and anxiety disorders are estimated to affect 3.76% of the global population.¹ Additionally, according to the World Health Organization, mental disorders rank among the most common causes of disability worldwide.²

According to the World Health Organization, mental health refers to “a condition of well-being that enables the individual to recognize and utilize his or her own potential, to manage the normal challenges of life, to work effectively

and productively, and to contribute positively to his or her community.”³

The availability and quality of mental health services and treatment vary widely across countries and cultures in

¹Department of Healthcare Management, School of Management and Medical Informatics, Shiraz University of Medical Sciences, Shiraz, Iran

²Department of Consultation and Mental Health, Yasuj University of Medical Sciences, Yasuj, Iran

Corresponding author:

Mohsen Khosravi, Department of Healthcare Management, School of Management and Medical Informatics, Shiraz University of Medical Sciences, Zand St, Shiraz 7134814336, Iran.
Email: mohsenkhosravi@live.com



the world. According to the World Health Organization, major depression as a mental illness is the most common cause of disability in terms of years lived with disability and the fourth most common cause of disability-adjusted life years globally.⁴ Moreover, the Health Canada Editorial Board on Mental Illnesses in Canada reports that more than one-fifth of Canadians will experience a mental illness in their lifetime, and the global economic impact of mental health in 2010 was estimated at 2.5 trillion U.S. dollars.⁵ Meanwhile, the prevalence of suicide is rising in many countries, such as the United States, indicating the urgency of developing new solutions and innovations in mental health.⁵

Shown as a major challenge in mental health services, the current clinical workforce is inadequate to address these needs; The number of psychiatrists per population is very low, with an average of 9 per 100,000 people in developed countries and as few as 0.1 per 1,000,000 people in some in lower-income regions. Meanwhile, the current and projected demand for care exceeds the available capacity, which necessitates the exploration of digital technology as a potential solution.^{4,6}

Interestingly, chatbots, also known as conversational agents or multipurpose virtual assistants, are an emerging technology that may facilitate the engagement and adherence of users with mental health issues.⁷ They have been deployed across various domains for a multitude of purposes. Specifically, in the realm of mental health, they have been utilized for therapeutic interventions, training programs, educational initiatives, counseling sessions, and screening procedures.^{8–10} Furthermore, they may reduce the stigma associated with seeking mental health-related advice and enhance the user experience of mobile mental health apps.¹¹ Moreover, they have been investigated for their effectiveness in promoting self-disclosure and expressive writing.^{7,12} Chatbots have also provided various types of social support, such as appraisal, informational, emotional, and instrumental support, to young people with mental health issues. While they have been designed to educate underprivileged communities on mental health and stigmatized topics.^{13,14}

There is emerging evidence of user engagement with chatbots for supporting various mental health issues and preliminary indications of positive health outcomes in the physical and mental health domains.¹⁵ The engagement of patients with chatbots in mental health services is a crucial factor for the successful implementation and adoption of this emerging technology. However, there is a lack of evidence-based guidance on how to design and evaluate chatbots for mental health purposes.¹⁶ Therefore, this article aims to conduct a systematic review of the existing reviews on the topic and identify the factors influencing the engagement of patients with chatbots in mental health services to synthesize the existing evidence and present it in a single document. The conclusions drawn from this research offer significant

implications for manufacturers, healthcare professionals, and policymakers who are strategizing to enhance the applicability of chatbots in the field of mental health services.

Methods

This study involved a systematic review of reviews by employing qualitative analysis of literature published between 2000 and 2024. The research question was: “what are the factors influencing patient engagement with mental health chatbots?” Furthermore, this study employed the Joanna Briggs Institute approach to execute a qualitative systematic review, which encompasses several stages.¹⁷ Initially, the authors identified and selected relevant studies by conducting a comprehensive search across multiple databases and implementing a rigorous screening process. Subsequently, the identified studies were critically evaluated using the Critical Appraisal Skills Programme (CASP) quality appraisal checklist. The authors then extracted pertinent data from the studies using a preestablished checklist. Following this, a thematic analysis was conducted on the data to synthesize the findings. Finally, the results were reported and analyzed in a comprehensive manner, providing a thorough understanding of the subject matter.

Data collection and search method

The authors conducted a systematic search of several databases, such as PubMed, Scopus, ProQuest, and Cochrane database of systematic reviews, to identify relevant studies on the topic. To identify relevant studies that met the essential criteria, we categorized the search terms into three themes, namely Mental health, Engagement, and Chatbot. We commenced our search by using broad terms to enhance sensitivity and incorporated synonyms using the “OR” operator. To ensure specificity and minimize irrelevant studies, we used the “AND” operator in our search strategy. Furthermore, Medical Subject Headings terms such as “mental health” were used in order to improve the search strategy. Our search strategy is presented in Table 1. The search was conducted on the 26th of January 2024 and was registered in GitHub with the following link: <https://github.com/mohsenkhosravi3913/Factors-Influencing-Patient-Engagement-in-Mental-Health-Chatbots.git>.

Inclusion and exclusion standards

The inclusion criteria for this study were articles published in English between 2000 and 2024 that specifically addressed the factors influencing patient engagement in mental health chatbots. Articles were excluded if they: (A) did not cover a chatbot corresponding to mental health services, (B) lacked a text describing the factors of patient engagement with a chatbot intervention in healthcare, or (C) weren't a review paper in terms of methodology.

Table 1. The search methodology utilized to perform the systematic review.

No.	Theme	Keywords
1	Mental health	Depress* OR anxi* OR anxiety OR mental health OR mental illness OR mental disorder OR mental wellbeing OR behavioral health OR obsess* OR obsessive behavior OR panic OR phobia OR schizophrenia* OR PTSD OR substance disorder* OR drug abuse* OR ADHD* OR bipolar* OR mood disorder* OR psychosis OR psychotic disorder* OR autism OR addiction* OR melancholia* OR eating disorder* OR anorexia*
2	Engagement	engag* OR Usability OR user experience OR satisfaction OR usage OR use OR using OR adopt* OR accept* OR perspective* OR percept* OR involve* OR barrier*
3	Chatbot	Chatbot* OR bot* OR conversational agent* OR conversational AI OR virtual agent* OR virtual AI
Scope	2000–2024, [English], PubMed, Scopus, ProQuest, Cochrane database of systematic reviews	

Screening and data retrieval

The systematic review was conducted in several stages. Firstly, the authors independently assessed the title of all articles obtained from the databases multiple times. In this procedure, we selected papers that appeared to discuss the factors influencing patient engagement with mental health chatbots. Any papers that seemed irrelevant were subsequently removed from further stages of the screening process. Secondly, the abstracts of selected articles were screened similar to the previous stage, and then the full text of chosen articles was thoroughly assessed. Furthermore, to ensure that no relevant studies were missed, the authors also reviewed the references cited in the articles. Finally, articles that met the validity criteria were selected for inclusion in the study.

The authors used a form to extract necessary information from selected articles, which was then summarized and combined using MAXQDA 12 software.¹⁸ MAXQDA is a software for qualitative data analysis. It provides tools for managing, transcribing, and analyzing data through coding. It also allows note-taking, data summarization, and visualization of patterns.¹⁹ The form utilized by the authors comprised several sections, including the year of the study's publication, its settings (the place in which the study was conducted), the associated journal, its indexed databases, the study type in terms of methodological approach, the type of chatbot(s) studied, corresponding disorders, and the number of final studies included within the study. During this process, the segments of the manuscript text that pertained to each section of the form were identified, coded, and subsequently incorporated into the final form. Furthermore, the authors verified the results at all stages to ensure reliability and minimize bias.

Quality appraisal of final articles using the CASP checklist

The quality of selected studies was evaluated using the CASP appraisal checklist, which covers multiple study designs. The checklist helps assess the validity, relevance, bias, and applicability of research studies.²⁰

The CASP checklist comprises ten questions that evaluate articles based on various factors, including result validity, study quality, and result applicability.²⁰ We used a scoring system where each question was assigned 2 points for yes, 1 point for cannot tell, and 0 points for no. The maximum score was 16, corresponding to three quality levels: low, medium, and high. Only articles with an average score of at least nine were included.²¹ The total score for each study was computed, and the score for each study type was presented in the results section. After screening and evaluating the quality of articles using the CASP checklist, only the final articles were included in our review.

Data analysis

The aim of this phase was to identify the primary themes of factors that affect patient engagement with mental health chatbots. The authors conducted a thematic analysis of the items extracted from the contents of the final articles obtained through the systematic review.

The authors followed Boyatzis's code development approach to perform the analysis.²² In this methodology, the authors initially read the text of the papers to familiarize themselves with the data; subsequently, they generated initial codes based on the objective of the analysis; the next step involved searching for themes by categorizing these codes into groups; following this, they reviewed the themes and began the process of defining and naming them; the final step was to produce the report. In this study, the authors extracted the factors affecting patient engagement with mental health chatbots separately from the final articles and compiled them into a single table using Microsoft Excel. The analysis resulted in mega-themes, themes and subthemes, based on the research question's framework. Furthermore, to improve the validity and reliability of the outcome and reduce the risk of error or bias, the authors repeated the same steps of the thematic analysis. They also consulted with each other to resolve any disagreements during the process, if needed.

Results

The following sections present the results from the systematic review of reviews, which consisted of three main steps: searching the databases, assessing the quality of the final articles, and conducting a thematic analysis of the data extracted from the final articles.

Systematic review

The execution of the systematic review of reviews was conducted in adherence to the PRISMA guidelines, which are specifically designed for reporting systematic reviews.²³ Figure 1 illustrates the results of the search within the

databases. The search strategy yielded 1494 references from the databases. After removing 12 duplicates, the remaining 1482 references were screened by their titles, abstracts, and full texts. Based on the inclusion and exclusion criteria, 1472 references were excluded from the study. Consequently, 10 papers were selected as the final articles for the study.

The characteristics of the final 10 studies are delineated in Appendix 1 (Bibliography of Final Articles). The studies employed different study types such as scoping reviews and systematic reviews, and were published between 2017 and 2023. Rule-based chatbots and machine learning-based chatbots were the most frequently used types of chatbots in these studies. Moreover, the studies addressed a broad




Figure 1. Results of the search within the databases.




Figure 2. Number of included studies within the final articles of this study.

range of disorders including depression, autism, anxiety, schizophrenia, posttraumatic stress disorder, substance use disorder, stress, mindfulness, and multiple other disorders. Additionally, the studies explored the potential of chatbots for promoting mental well-being, treating cooccurring depression and anxiety, and managing multiple health behaviors. Furthermore, the studies were conducted in diverse settings including the United States, Japan, Australia, China, Germany, the United Kingdom and multiple other countries. The number of final articles reviewed by the studies included in the current study varied between 10 and 54, indicating heterogeneity in the scope of the studies (Figure 2).

Quality assessment of final articles

The quality assessment was done using the CASP Systematic Review Checklist, which consisted of 10 questions that evaluated the validity, results, and applicability of the review.²⁰ Each question had been scored as yes (2 points), can't tell (1 point), or no (0 point), and the final score was calculated as the sum of the scores for each question. The data revealed that the majority of the 10 articles had the same final score of 9/16, which indicated a moderate level of validity, but also some limitations and gaps. The main areas where the articles scored poorly were the inclusion of all important and relevant studies, the assessment of the quality of the included studies, the consideration of all important outcomes, and the benefits, harms, and costs of the intervention. Moreover, the level of bias in the included studies was demonstrated to be reasonable. Further information regarding the quality assessment of included papers can be seen in Appendix 2 (quality assessment of final articles).

Thematic analysis

Table 2 presents the results of the thematic analysis of the data extracted from the final articles included in the study.

The thematic analysis summarized the factors that influence user engagement with chatbots, conversational agents, or virtual health assistants for mental health or clinical psychology into two broad mega-themes of “chatbot factors” and “user factors.” The analysis identified four main themes: chatbot design, chatbot outcomes, user perceptions, and user characteristics. Each theme had several sub-themes that synthesized the findings from the literature.

Chatbot design. This theme comprised subthemes such as purpose, platform, response generation, dialog initiative, input and output modalities, embodiment, targeted disorders, personalization, interactivity, verbal and nonverbal behavior, anonymity, and intensity and duration. These subthemes described the characteristics and features of the chatbot that can affect user engagement. For instance, the purpose of the chatbot, whether it is for therapy, training, or screening, can influence engagement. The platform on which the chatbot is implemented, such as stand-alone software or a mobile app, can affect the engagement with the chatbots. Furthermore, the method used by the chatbot to generate responses, such as rule-based or machine learning-based, can influence the quality of the engagement.

Chatbot outcomes. This theme included subthemes such as effectiveness, measurement, support, security, privacy, and responsiveness. The effectiveness of the chatbot in improving mental health outcomes can influence engagement. Moreover, the use of appropriate outcome measurement instruments to evaluate the effectiveness of the chatbot can affect user engagement. Meanwhile, the availability of support for the chatbot app, including customer service and technical assistance, can affect user engagement.

User perceptions. This theme consisted of subthemes such as usefulness, ease of use, comparison, content, enjoyability, trustworthiness, attractiveness, acceptability, understandability, empathy, and affordability. The perceived usefulness of the chatbot in providing mental health support can influence engagement. Meanwhile, the ease of use of the chatbot can affect engagement. Furthermore, the comparison with other forms of mental health support can affect user's overall engagement.

User characteristics. This theme is comprised of subthemes such as age, gender, and mental health condition. These subthemes described the demographic, personal and health condition of the users which can influence engagement of the users from the services acquired via the chatbots.

Discussion

The results indicated that user engagement with the mental health services delivered by chatbots is influenced by several categories, such as chatbot design, chatbot

Table 2. Thematic analysis of the data extracted from the final articles.

Mega-theme	Theme	Subtheme	Reference
Chatbot factors	Chatbot design	Purpose: The purpose of the chatbot, whether it is for therapy, training, or screening, can influence engagement.	24
		Platform: The platform on which the chatbot is implemented, such as stand-alone software or a mobile app, can affect engagement.	24,25
		Response generation: The method used by the chatbot to generate responses, such as rule-based or machine learning-based, can influence the quality of user engagement.	24–26
		Dialog initiative: Whether the chatbot controls and leads the conversation or allows the user to take the initiative can affect engagement.	24
		Input and output modalities: The input and output modalities used by the chatbot, such as written language only or a combination of written, spoken, and visual languages, can influence engagement.	24,25
		Embodiment: The inclusion of a virtual representation of the chatbot (the use of a human-like avatar or voice) can affect user engagement.	24,27,28
		Targeted disorders: The specific mental health disorders targeted by the chatbot, such as depression or autism, can influence user engagement.	24
		Personalization: Personalization of the virtual health assistant to the user's needs and preferences can improve engagement.	24,26,29–31
		Interactivity: Interactive features, such as the ability to ask questions and receive feedback, can enhance user engagement.	24,26,28
		Verbal and nonverbal behavior: The ability of the chatbot to simulate key properties of human face-to-face conversation, such as verbal and nonverbal behavior, can enhance user engagement.	28,31
		Anonymity: The anonymity provided by mental health chatbots can enhance users.	29
		Intensity and duration: The characteristics of the chatbot intervention, such as its duration and intensity, can affect engagement.	26,30,32
	Chatbot outcomes	Effectiveness: The effectiveness of the chatbot in improving mental health outcomes can influence engagement.	29,32
		Measurement: The use of appropriate outcome measurement instruments to evaluate the effectiveness of the chatbot can affect user engagement.	32
		Support: The availability of support for the chatbot app, including customer service and technical assistance, can affect user engagement.	27,28,32,33
		Security: The security measures implemented by the chatbot app including its ability to manage crisis situations can affect user engagement.	27,28,32
		Privacy: Concerns about privacy were mentioned in negative reviews of chatbot apps.	29
		Responsiveness: The responsiveness of the chatbot to user input can influence user engagement.	33

(continued)

Table 2. Continued.

Mega-theme	Theme	Subtheme	Reference
User factors	User perceptions	Usefulness: The perceived usefulness of the chatbot in providing mental health support can influence engagement.	28,29,32,33
		Ease of use: The ease of use of the chatbot can affect engagement.	28,29,32,33
		Comparison: Comparisons with other forms of mental health support can affect user's overall engagement.	32,33
		Content: The quality and relevance of the content provided by the chatbot can influence engagement.	26,27,33
		Enjoyability: The enjoyability of interacting with the chatbot can affect engagement.	31,33
		Trustworthiness: The perceived trustworthiness of the chatbot can influence user engagement.	27,33
		Attractiveness: The visual attractiveness of the chatbot can affect user engagement.	33
		Acceptability: The acceptability of the chatbot as a form of mental health support can influence engagement.	26,29,33
		Understandability: The ability of the chatbot to provide understandable responses can affect engagement.	33
		Empathy: The ability of the virtual health assistant to demonstrate empathy and understanding can enhance user engagement.	26,27,30,31
		Affordability: The affordability of mental health chatbots, including their cost and availability of insurance coverage, can affect engagement.	27,28,32,33
	User characteristics	Age and gender: demographic characteristics of the user such as age and gender can influence engagement.	25,32
		Mental health condition: The mental health condition of the user, can influence engagement.	25,32

outcomes, user perceptions, and user characteristics. This section discussed these categories and their implications for chatbot development and evaluation.

Chatbot design

The findings of the study declared several chatbot design factors, such as purpose, platform, response generation, dialog initiative, input and output modalities, embodiment, targeted disorders, personalization, interactivity, verbal and nonverbal behavior, anonymity, and intensity and duration. The findings implied that these factors have an influence on how patients perceive and interact with chatbots in mental health services.

The literature reveals the impact of color on mood, anxiety, aggression, and depression, which may affect engagement with mental health chatbots. For instance, the color blue has been associated with higher levels of

trustworthiness and mental alertness in relation to digital platforms.³⁴ Hence, investigating the optimal and effective colors for the design of mental health chatbots can be a crucial step toward enhancing not only the engagement with the chatbots in the design category, but also the attitude of the users toward the chatbots in the perception category.

The literature also indicates that the incorporation of music in mental health chatbots can be a potential strategy to enhance both the engagement of patients with the chatbots in the design category and the treatment of some mental health disorders in an efficient and effective way in the outcome category.^{35,36} In this regard, the implementation of music therapy which is a description of a systematic intervention process in which the clients are assisted to enhance their health, by using musical experiences and the emerging relationships from them as dynamic agents of change can be a strategic policy by the chatbot designers.³⁷

Chatbot outcomes

The findings of the study manifested several chatbot outcome factors, such as effectiveness, measurement, support, security, privacy, and responsiveness as factors influencing patient engagement with chatbots in mental health services.

Concerning the effectiveness of chatbots, it is said that chatbots should follow the principles of other Behavioral Intervention Technologies (BITs) by adopting an efficiency model of support, i.e. a conceptual framework that captures the interaction between information and intervention in an optimal way. This model suggests that decisions should be made based on the analysis of why people may not benefit from BITs.³⁸ Moreover, chatbots that provide rapid diagnoses may undermine diagnostic practice, which requires practical wisdom and collaboration among different specialists as well as close communication with patients.³⁹

Concerning the security and safety of chatbots, the interactions between patients and chatbots should be rigorously monitored, as inaccurate chatbot responses may lead to unintended harm. Particularly, as chatbots increase their conversational flexibility, there may be more potential errors associated with natural language understanding or response generation. Hence, using unbounded chatbots should be accompanied by careful supervision of patient and chatbot interactions, and of safety functions.⁴⁰

In summary, the evidence regarding the factors influencing the engagement of patients with chatbots in the outcome category seems to be insufficient for any definitive conclusion. Therefore, it seems that further research on this category is needed to obtain a comprehensive view of the factors influencing patient engagement with mental health chatbots in the outcome category.

User perceptions

The findings of the study declared factors such as usefulness, ease of use, comparison, content, enjoyability, trustworthiness, attractiveness, acceptability, understandability, empathy, and affordability influencing patient's engagement with mental health chatbots.

Maintaining users' preferences and demands to enhance their perception of chatbot usage has the potential to realize the long-standing concept of patient-centeredness particularly person-centeredness and person therapy in clinical services. Person-centered therapy, developed by Carl Rogers, is a nondirective approach that emphasizes the client's potential for self-actualization and self-healing. The therapist's attitude, relationship, and empathy are the key factors in the therapeutic process.³⁹

It is noteworthy that various international organizations, such as the G-20 summit, a prominent international forum, have recognized the need for patient-centered or person-centered care. They have emphasized the importance of this approach as a strategic policy for developing and implementing healthcare plans.⁴¹

In such context, healthcare chatbots can facilitate the implementation of person-centered or patient-centered care within the healthcare service delivery networks which have been shown to enhance not only the clinical criteria of healthcare services but also the criteria related to user perception of healthcare services, such as user engagement.^{42–44} Therefore, healthcare chatbots can become the primary source of referral when users need healthcare services; This phenomenon indicates the significance of taking into account the factors affecting patient engagement within the process of delivering healthcare services via chatbots.

User characteristics

The findings showed several demographic and health status factors such as age, gender, and mental health condition influencing patient engagement with mental health chatbots.

The World Health Organization and other organizations have advocated for a feminist global health agenda that aims to address gender inequalities and barriers in health care and to achieve gender equity and empowerment for women and girls in the provision of healthcare services.⁴⁵ For instance, considering that the female population in certain regions of the world, especially Muslims in the Middle East may have distinct values and preferences, the development of chatbots that respect gender preferences could be a significant advancement in such a feminist agenda in the health care systems.⁴⁶ Since, the gender of healthcare providers, the discomfort and insecurity of being in mixed-gender settings, and the gender of the medical staff are all important aspects of person-centered care from the perspective of Muslim women who, as adherents of Islam, generally avoid any direct contact with the opposite gender due to their beliefs.⁴⁷

Furthermore, chatbots are deemed an unsuitable intervention for addressing health conditions that have high perceived stigma and severity, according to the evidence. As a result, healthcare customers may not use this technology as an independent intervention and should not rely on it as a substitute for a trustworthy health information source from a health professional. However, chatbots could be a useful adjunct for enhancing doctor–patient communication for conditions with lower perceived stigma and severity.⁴⁸ However, future studies should determine the scope of these conditions and explore how chatbots could encourage more disclosure of sensitive information to health professionals who could then provide more relevant healthcare services.

Limitations and implications of the study

The study had some limitations, such as the lack of consideration of the tradeoffs or conflicts between different categories of the factors influencing patient engagement with chatbots. The study also didn't account for the variability or diversity of user perceptions across different groups,

contexts, or cultures, and didn't include other potential user characteristics that may affect user engagement with chatbots, which were not found through the evidence from the search.

The study also had some implications, such as the need for chatbot developers to consider user preferences and needs when designing chatbots, to adopt a holistic and multidimensional approach to measure and optimize chatbot outcomes, to conduct user research and testing to understand and address user perceptions, and to provide chatbots that are accessible and inclusive for different users. Furthermore, the study reviewed the literature on the impact of color and music on mental health and suggested some strategies for incorporating these factors in chatbot design. Finally, based on the findings of the study, it is suggested that future original researches are required to explore and describe unobserved domains of patient engagement within mental health chatbots.

Conclusions

This research investigated the factors affecting patient engagement with mental health chatbots, and found four categories of factors: chatbot design, chatbot outcomes, user perceptions, and user characteristics. It also examined the literature on the role of color and music in mental health and suggested some ways to integrate these factors in chatbot design. Furthermore, the research offered some practical and theoretical insights for chatbot developers and researchers. Moreover, it highlighted the potential of chatbots to facilitate person-centered or patient-centered care in mental health services and to improve the clinical and perceptual criteria of healthcare outcomes. Finally, further research is needed to obtain a comprehensive view of the factors influencing patient engagement with chatbots in mental health services, and there is a need for more collaboration and communication between chatbot developers, researchers, health professionals, and users.

Acknowledgments: We would like to acknowledge the Bing AI chatbot for its contribution to rewriting the manuscript in terms of English grammar and wording of the text.

Contributorship: Ghazaleh theorized the project and cooperated in writing the introduction and discussion sections of the manuscript. Mohsen performed the search within the databases, conducted the thematic analysis of the findings and wrote the text of the manuscript.

Consent statement: Not applicable; Since the study is a review in methodology and no individual (patient) has been included in current research.

Declaration of conflicting interests: The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Ethical approval: Not applicable to this methodology of research.

Funding: The authors received no financial support for the research, authorship, and/or publication of this article.

Guarantor: The guarantor for the whole process of the manuscript is the corresponding author.

ORCID ID: Mohsen Khosravi  <https://orcid.org/0000-0002-0576-7660>

Supplemental material: Supplementary material for this article is available online.

References

1. Elfleijn J. Share of the population worldwide who suffered from anxiety disorders from 1990 to 2017. Statista, 2020.
2. Rudd BN and Beidas RS. Digital mental health: the answer to the global mental health crisis? *JMIR Ment Health* 2020; 7: e18472.
3. World Health Organization. Mental health: a state of well-being. World Health Organization, 2014.
4. Murray CJ, Vos T, Fau - Lozano R, et al. Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. (1474-547X (Electronic)).
5. Trautmann S, Rehm J and Wittchen HU. The economic costs of mental disorders: Do our societies react appropriately to the burden of mental disorders? (1469-3178 (Electronic)).
6. Oladeji BD and Gureje O. Brain drain: a challenge to global mental health. *BJPsych Int* 2016; 13: 61–63.
7. Lee Y-C, Yamashita N and Huang Y. Designing a chatbot as a mediator for promoting deep self-disclosure to a real mental health professional. *Proc ACM Hum-Comput Interact* 2020; 4: 1–27.
8. Haque MDR and Rubya S. An overview of chatbot-based mobile mental health apps: insights from app description and user reviews. *JMIR mHealth uHealth* 2023; 11: e44838.
9. Abd-Alrazaq AA, Rababeh A, Alajlani M, et al. Effectiveness and safety of using chatbots to improve mental health: systematic review and meta-analysis. *J Med Internet Res* 2020; 22: e16021.
10. Wilson L and Marasoiu M. The development and use of chatbots in public health: scoping review. *JMIR Hum Factors* 2022; 9: e35882.
11. Oh YJ, Zhang J, Fang M-L, et al. A systematic review of artificial intelligence chatbots for promoting physical activity, healthy diet, and weight loss. *Int J Behav Nutr Phys Act* 2021; 18: 160.
12. Sannon S, Stoll B, Difranzo D, et al. How personification and interactivity influence stress-related disclosures to conversational agents. 2018, pp.285–288.
13. Rahman R, Rahman M, Tripto N, et al. AdolescentBot: understanding opportunities for Chatbots in combating adolescent sexual and reproductive health problems in Bangladesh. 2021, pp.1–15.
14. Yadav D, Malik P, Dabas K, et al. Feedpal: understanding opportunities for chatbots in breastfeeding education of women in India. *Proc ACM Hum-Comput Interact* 2019; 3: Article 170.

15. Pham KT, Nabizadeh A and Selek S. Artificial intelligence and chatbots in psychiatry. *Psychiatric Quarterly* 2022; 93: 249–253.
16. Shan Y, Zhang J, Li Z, et al. Mental health assessment for the chatbots. arXiv preprint arXiv:220105382. 2022.
17. Santos WMD, Secoli SR and Püschel VAA. The Joanna Briggs Institute approach for systematic reviews. *Rev Lat Am Enfermagem* 2018; 26: e3074.
18. Software V. MAXQDA 2022 Berlin, Germany 2021.
19. MAXQDA. Why MAXQDA? <https://www.maxqda.com/why-maxqda> (2023).
20. Programme CAS. CASP checklist: 10 questions to help you make sense of a systematic review. 2018.
21. Hannes K, Lockwood C and Pearson A. A comparative analysis of three online appraisal instruments' ability to assess validity in qualitative research. *Qual Health Res* 2010; 20: 1736–1743.
22. Boyatzis RE. *Transforming qualitative information: thematic analysis and code development*. Thousand Oaks, CA, USA: Sage Publications, Inc, 1998, pp.184–xvi.
23. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *Br Med J* 2021; 372: n71.
24. Abd-Alrazaq AA, Alajlani M, Alalwan AA, et al. An overview of the features of chatbots in mental health: a scoping review. *Int J Med Inform* 2019; 132: 103978.
25. Li H, Zhang R, Lee YC, et al. Systematic review and meta-analysis of AI-based conversational agents for promoting mental health and well-being. *NPJ Digit Med* 2023; 6: 236.
26. Gaffney H, Mansell W and Tai S. Conversational agents in the treatment of mental health problems: mixed-method systematic review. *JMIR Ment Health* 2019; 6: e14166.
27. Curtis RG, Bartel B, Ferguson T, et al. Improving user experience of virtual health assistants: scoping review. *J Med Internet Res* 2021; 23: e31737.
28. Provoost S, Lau HM, Ruwaard J, et al. Embodied conversational agents in clinical psychology: a scoping review. *J Med Internet Res* 2017; 19: e151.
29. Vaidyam AN, Wisniewski H, Halamka JD, et al. Chatbots and conversational agents in mental health: a review of the psychiatric landscape. *Can J Psychiatry* 2019; 64: 456–464.
30. He Y, Yang L, Qian C, et al. Conversational agent interventions for mental health problems: systematic review and meta-analysis of randomized controlled trials. *J Med Internet Res* 2023; 25: e43862.
31. Xue J, Zhang B, Zhao Y, et al. Evaluation of the current state of chatbots for digital health: scoping review. *J Med Internet Res* 2023; 25: e47217.
32. Jabir AI, Martinengo L, Lin X, et al. Evaluating conversational agents for mental health: scoping review of outcomes and outcome measurement instruments. *J Med Internet Res* 2023; 25: e44548.
33. Abd-Alrazaq AA, Alajlani M, Ali N, et al. Perceptions and opinions of patients about mental health chatbots: scoping review. *J Med Internet Res* 2021; 23: e17828.
34. Elliot AJ. Color and psychological functioning: a review of theoretical and empirical work. *Front Psychol* 2015; 6: 368.
35. Nizamie SH and Tikka SK. Psychiatry and music. *Indian J Psychiatry* 2014; 56: 128–140.
36. Golden TL, Springs S, Kimmel HJ, et al. The use of music in the treatment and management of serious mental illness: a global scoping review of the literature. *Front Psychol* 2021; 12: 649840.
37. Bruscia KE. Defining music therapy. (No Title). 1989. 14–17.
38. Schueller S, Tomasino K and Mohr D. Integrating human support into behavioral intervention technologies: the efficiency model of support. *Clin Psychol: Sci Pract* 2016; 24: 27.
39. Parviainen J and Rantala J. Chatbot breakthrough in the 2020s? An ethical reflection on the trend of automated consultations in health care. *Med Health Care Philos* 2022; 25: 61–71.
40. Laranjo L, Dunn AG, Tong HL, et al. Conversational agents in healthcare: a systematic review. *J Am Med Inform Assoc* 2018; 25: 1248–1258.
41. GHM D. G20 Health Ministers' Declaration. Riyadh, Saudi Arabia, 2020.
42. Edvardsson D, Fetherstonhaugh D, Fau - McAuliffe L, et al. Job satisfaction amongst aged care staff: exploring the influence of person-centered care provision. 2011(1741-203X (Electronic)).
43. Willard-Grace R, Hessler D, Fau-Rogers E, et al. Team structure and culture are associated with lower burnout in primary care. *J Am Board Fam Med* 2014; 27: 229–238. (1558-7118 (Electronic)) .
44. Edvardsson D, Sandman PO and Borell L. Implementing national guidelines for person-centered care of people with dementia in residential aged care: effects on perceived person-centeredness, staff strain, and stress of conscience. 2014(1741-203X (Electronic)).
45. Davies SE, Harman S, Manjoo R, et al. Why it must be a feminist global health agenda. *Lancet* 2019; 393: 601–603.
46. Tackett S, Young JH, Putman S, et al. Barriers to healthcare among Muslim women: a narrative review of the literature. *Womens Stud Int Forum* 2018; 69: 190–194.
47. Inhom MC and Serour GI. Islam, medicine, and Arab-Muslim refugee health in America after 9/11. *Lancet* 2011; 378: 935–943.
48. Ayers JW, Poliak A, Dredze M, et al. Comparing physician and artificial intelligence chatbot responses to patient questions posted to a public social media forum. *JAMA Intern Med* 2023; 183: 589–596.