

Homework 7

ALECK ZHAO

March 30, 2017

1. Let R be a ring, and let σ be an automorphism of R . Show that $\{a \in R \mid \sigma(a) = a\}$ is a subring of R , and a subfield if R is a field.

Proof. Call the subset S . Any automorphism must fix 1, so $1 \in S$. Now if $a, b \in S$, we have

$$\begin{aligned}\sigma(a + b) &= \sigma(a) + \sigma(b) = a + b \\ \sigma(ab) &= \sigma(a)\sigma(b) = ab\end{aligned}$$

so $a + b, ab \in S$, so S is indeed a subring. Now, if R is a field, then for all nonzero $a \in R$,

$$1 = \sigma(1) = \sigma\left(a \cdot \frac{1}{a}\right) = \sigma(a)\sigma\left(\frac{1}{a}\right)$$

Now, if $a \in S$, then $\sigma(a) = a$, so

$$\sigma\left(\frac{1}{a}\right) = \frac{1}{\sigma(a)} = \frac{1}{a}$$

so $\frac{1}{a} \in S$ as well, and thus S is a field. □

2. Let F be a finite field with p^n elements for p a prime. Show that each element $a \in F$ has a p th root in F , i.e. there exists $b \in F$ such that $b^p = a$. Is b unique? By contrast, for $K := F(x)$ the fraction field of the polynomial ring $F[x]$, show that x has no p th root in K .

Section 6.4: Finite Fields

8. Find $[\mathbb{F}_{p^n} : \mathbb{F}_{p^m}]$ where $m \mid n$.
18. (a) Show that a monic irreducible polynomial $f \in F[x]$ has no repeated root in any splitting field over F if and only if $f \not\equiv 0$ in $F[x]$.
 (b) If $\text{char } F = 0$, show that no irreducible polynomial has a repeated root in any splitting field over F .
19. If $\text{char } F = p$, show that a monic irreducible polynomial $f \in F[x]$ has a repeated root in some splitting field if and only if $f = g(x^p)$ for some $g \in F[x]$. (Hint: Ex 18)
21. Let p be a prime and write $f = x^p - x - 1$. Show that the splitting field of f over \mathbb{F}_p is $\mathbb{F}_p(u)$, where u is any root of f . (Hint: Compute $f(u + a)$, $a \in \mathbb{F}_p$)
22. (a) Let f be a monic irreducible polynomial of degree n in $\mathbb{F}_p[x]$. Show that f divides $x^{p^n} - x$ in $\mathbb{F}_p[x]$. (Hint: First work over $\mathbb{F}_p(u)$, $f(u) = 0$. Use the uniqueness in Theorem 4 § 4.1.)
 (b) Show that the degree of each monic irreducible divisor f of $x^{p^n} - x$ is a divisor of n . (Hint: Theorem 5)
 (c) Factor $x^8 - x$ into irreducibles in $\mathbb{F}_2[x]$.

Section 4.5: Symmetric Polynomials

14. Given $\sigma \in S_n$, define $\theta_\sigma : R[x_1, \dots, x_n] \rightarrow R[x_1, \dots, x_n]$ by $\theta_\sigma[f(x_1, \dots, x_n)] = f(x_{\sigma 1}, \dots, x_{\sigma n})$.

(a) Show that θ_σ is a ring automorphism of $R[x_1, \dots, x_n]$.

Proof. First we show this is a ring homomorphism. Clearly $\theta_\sigma(1) = 1$. Now, for $f, g \in R[x_1, \dots, x_n]$,

$$\begin{aligned} \theta_\sigma[f(x_1, \dots, x_n) + g(x_1, \dots, x_n)] &= \theta_\sigma[(f + g)(x_1, \dots, x_n)] \\ &= (f + g)(x_{\sigma 1}, \dots, x_{\sigma n}) \\ &= f(x_{\sigma 1}, \dots, x_{\sigma n}) + g(x_{\sigma 1}, \dots, x_{\sigma n}) \\ &= \theta_\sigma(f) + \theta_\sigma(g) \\ \theta_\sigma[f(x_1, \dots, x_n) \cdot g(x_1, \dots, x_n)] &= \theta_\sigma[(fg)(x_1, \dots, x_n)] \\ &= (fg)(x_{\sigma 1}, \dots, x_{\sigma n}) \\ &= f(x_{\sigma 1}, \dots, x_{\sigma n}) \cdot g(x_{\sigma 1}, \dots, x_{\sigma n}) \\ &= \theta_\sigma(f) \cdot \theta_\sigma(g) \end{aligned}$$

Now if

$$\theta_\sigma(f) = f(x_{\sigma 1}, \dots, x_{\sigma n}) = g(x_{\sigma 1}, \dots, x_{\sigma n}) = \theta_\sigma(g)$$

then consider σ^{-1} and its associated $\theta_{\sigma^{-1}}$. Then applying $\theta_{\sigma^{-1}}$ to both of these polynomials,

$$\begin{aligned} \theta_{\sigma^{-1}}[f(x_{\sigma 1}, \dots, x_{\sigma n})] &= f(x_{\sigma^{-1}\sigma 1}, \dots, x_{\sigma^{-1}\sigma n}) = f(x_1, \dots, x_n) \\ \theta_{\sigma^{-1}}[g(x_{\sigma 1}, \dots, x_{\sigma n})] &= g(x_{\sigma^{-1}\sigma 1}, \dots, x_{\sigma^{-1}\sigma n}) = g(x_1, \dots, x_n) \end{aligned}$$

so θ_σ is injective. Now, for any $f(x_1, \dots, x_n)$, we have

$$\theta_\sigma[f(x_{\sigma^{-1}1}, \dots, x_{\sigma^{-1}n})] = f(x_1, \dots, x_n)$$

so θ_σ is surjective. Thus, θ_σ is a bijective ring homomorphism from $R[x_1, \dots, x_n]$ to itself, so it is a ring automorphism. \square

(b) Show that $\sigma \mapsto \theta_\sigma$ is a group homomorphism $S_n \rightarrow \text{aut } R[x_1, \dots, x_n]$, which is injective.

Proof. Let $\sigma, \tau \in S_n$. Then consider $\theta_{\sigma\tau}$. For some $f(x_1, \dots, x_n) \in R[x_1, \dots, x_n]$, we have

$$\begin{aligned} \theta_{\sigma\tau}[f(x_1, \dots, x_n)] &= f(x_{\sigma\tau 1}, \dots, x_{\sigma\tau n}) \\ &= \theta_\sigma[f(x_{\tau 1}, \dots, x_{\tau n})] \\ &= \theta_\sigma(\theta_\tau[f(x_1, \dots, x_n)]) \\ &= (\theta_\sigma \circ \theta_\tau)[f(x_1, \dots, x_n)] \end{aligned}$$

so $(\sigma\tau) \mapsto \theta_{\sigma\tau} = \theta_\sigma \circ \theta_\tau$ and this is indeed a group homomorphism. Suppose $\theta_\sigma = \theta_\tau$ for some $\sigma, \tau \in S_n$. \square

i++i

(c) If $G \subseteq \text{aut } R[x_1, \dots, x_n]$ is a subgroup, show that $S_G = \{f \mid \theta(f) = f, \forall \theta \in G\}$ is a subring of $R[x_1, \dots, x_n]$.