

## Homework 5

ALECK ZHAO

May 4, 2018

### 1 Analytical (15 points)

**1) (15 points)** The probability density function of most Markov Random Fields cannot be factorized as the product of a few conditional probabilities. This question explores some MRFs that can be factorized in this way.

Consider the graph structure in Figure 1. From this graph, we know that  $X_2$  and  $X_3$  are conditionally




Figure 1: The Original Undirected Graph

independent given  $X_1$ . We can draw the corresponding directed graph as Figure 2. This suggests the following




Figure 2: The Converted Directed Graph

factorization of the joint probability:

$$P(X_1, X_2, X_3) = P(X_3|X_1)P(X_2|X_1)P(X_1)$$

Now consider the following graphical model in Figure 3.




Figure 3: An Undirected Graph

As before, we can read the conditional independence relations from the graph.

- (a) Following the example above, write a factorization of the joint distribution:

$$P(X_1, X_2, X_3, X_4, X_5, X_6, X_7).$$

*Solution.* We can let  $x_1$  be the root of a tree, so then the factorization would be

$$p(x_1)p(x_2 | x_1)p(x_3 | x_2)p(x_4 | x_2)p(x_5 | x_3)p(x_6 | x_3)p(x_7 | x_4)$$

□

- (b) Is this factorization unique, meaning, could you have written other factorizations that correspond this model? If the factorization is unique, explain why it is unique. If it is not unique, provide an alternate factorization.

*Solution.* This factorization is unique because the graph is a tree structure with no cycles or cliques. □

- (c) What is it about these examples that allows them to be factored in this way? Show an example MRF that cannot be factored in this way.

*Solution.* These examples are both tree structures without cliques, so there exist conditional independences. Consider the complete graph on 3 nodes: there is no conditional independence given only a single node, so there is no way to simplify the factorization. □