

Precalculus Practice Problems: Final

Alan Zhou

2024-2025

The focus of these review problems is on the material covered in Weeks 25 through 35, but keep in mind that prior material can still appear on the exam.

Contents

1	Matrices in 2D	2
1.1	Review Problems	2
1.2	Challenge Problems	4
1.3	Answers	5

1 Matrices in 2D

1.1 Review Problems

Review problems are meant to cover “standard” definitions and calculations as well as the use of some important results.

Throughout, $\hat{\mathbf{i}} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $\hat{\mathbf{j}} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$ are the standard unit vectors while $\mathbf{0} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$ is the zero vector.

We also let $\mathbf{I} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ be the (2×2) identity matrix and $\mathbf{0} = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$ be the zero matrix.

1. *Vector calculations.* Let $\mathbf{u} = \begin{pmatrix} 2 \\ 3 \end{pmatrix}$ and $\mathbf{v} = \begin{pmatrix} 4 \\ -1 \end{pmatrix}$. Compute each of the following.
 - (a) $\mathbf{u} + \mathbf{v}$
 - (b) $2\mathbf{v}$
 - (c) $\mathbf{u} \cdot \mathbf{v}$ and $\mathbf{v} \cdot \mathbf{u}$
 - (d) $\|\mathbf{u}\|$, $\|\mathbf{v}\|$, and $\|\mathbf{u} + \mathbf{v}\|$
 - (e) The angle between \mathbf{u} and \mathbf{v} (in terms of an inverse trig function)
 - (f) $\text{proj}_{\mathbf{v}}(\mathbf{u})$ and $\text{proj}_{\mathbf{u}}(\mathbf{v})$
2. *Applying matrices to vectors.* Let $\mathbf{A} = \begin{pmatrix} 2 & 4 \\ 1 & 1 \end{pmatrix}$ and $\mathbf{v} = \begin{pmatrix} 5 \\ 2 \end{pmatrix}$.
 - (a) Compute $\mathbf{A}\mathbf{v}$
 - (b) Find a vector \mathbf{u} for which $\mathbf{A}\mathbf{u} = \mathbf{v}$, or show that none exists.
3. *Matrix operations.* Let $\mathbf{A} = \begin{pmatrix} 2 & 4 \\ 1 & 1 \end{pmatrix}$ and $\mathbf{B} = \begin{pmatrix} -3 & 4 \\ 5 & -7 \end{pmatrix}$. Compute each of the following.
 - (a) $\mathbf{A} + \mathbf{B}$
 - (b) $-3\mathbf{A}$
 - (c) \mathbf{AB}
 - (d) \mathbf{BA}
 - (e) \mathbf{B}^T (the transpose of \mathbf{B})
4. *Geometric transformations.* Write down matrices for each of the following.
 - (a) Dilation about the origin by a factor of 4
 - (b) Horizontal dilation by a factor of 3 and vertical dilation by a factor of 2
 - (c) Rotation about the origin by $\pi/4$ counterclockwise
 - (d) Projection onto the line $y = (3/2)x$
 - (e) Reflection across the line $y = (3/2)x$

5. *Matrix determinants.* Let $A = \begin{pmatrix} 2 & 4 \\ 1 & 1 \end{pmatrix}$ and $B = \begin{pmatrix} -3 & 4 \\ 5 & -7 \end{pmatrix}$. Compute each of the following.

- (a) $\det A$ and $\det B$
- (b) $\det(AB)$
- (c) $\det(A^T)$
- (d) $\det(A + B)$
- (e) The area of the ellipse formed by applying A to the unit circle

6. *Matrix inverses.* Let $A = \begin{pmatrix} 2 & 4 \\ 1 & 1 \end{pmatrix}$ and $B = \begin{pmatrix} -3 & 4 \\ 5 & -7 \end{pmatrix}$. Compute each of the following.

- (a) A^{-1} and B^{-1}
- (b) $A^{-1}B^{-1}$ and $B^{-1}A^{-1}$
- (c) $(AB)^{-1}$
- (d) $(A^T)^{-1}$
- (e) $(A + B)^{-1}$
- (f) $\det(A^{-1})$

7. *Shear transformations.* A **horizontal shear** is given by a matrix of the form $\begin{pmatrix} 1 & k \\ 0 & 1 \end{pmatrix}$.

- (a) Describe the image of the unit square with vertices $(0, 0)$, $(1, 0)$, $(1, 1)$, and $(0, 1)$ when the horizontal shear $\begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}$ is applied.
- (b) By what factor does a horizontal shear multiply areas?
- (c) Find real constants a, b, k, θ for which

$$\begin{pmatrix} 4 & 1 \\ 3 & 7 \end{pmatrix} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix} \begin{pmatrix} 1 & k \\ 0 & 1 \end{pmatrix}.$$

(The constant θ can be expressed in terms of an inverse trig function.)

1.2 Challenge Problems

Challenge problems are meant to provide optional extensions of the ideas from class.

8. The **trace** of a square matrix is the sum of its main diagonal entries,

$$\operatorname{tr} \begin{pmatrix} a & b \\ c & d \end{pmatrix} = a + d.$$

- (a) For the matrices A, B in problems 3, 5, 6, compute $\operatorname{tr} A$, $\operatorname{tr} B$, $\operatorname{tr}(A + B)$, and $\operatorname{tr}(AB)$.
 - (b) Show that for any 2×2 matrices P and Q , we have $\operatorname{tr}(PQ) = \operatorname{tr}(QP)$.
 - (c) In general, must it be true that $\operatorname{tr}(ABC) = \operatorname{tr}(ACB)$?
9. Two matrices A, B are **similar**, written $A \sim B$, if there is an invertible P with $B = P^{-1}AP$.
- (a) Show that the only matrix similar to I is I .
 - (b) Show that if $A \sim B$, then $\det A = \det B$ and $\operatorname{tr} A = \operatorname{tr} B$.
 - (c) Let $A = \begin{pmatrix} 3 & 1 \\ 2 & 2 \end{pmatrix}$. There is exactly one diagonal matrix $D = \begin{pmatrix} d_1 & 0 \\ 0 & d_2 \end{pmatrix}$ with $d_1 \geq d_2$ for which $D \sim A$. Find D .
10. If A is a square matrix, the **characteristic polynomial** of A is defined by

$$f_A(X) = \det(A - XI).$$

- (a) Compute the characteristic polynomial $f_A(X)$ of the matrix $A = \begin{pmatrix} 3 & 1 \\ 2 & 2 \end{pmatrix}$.
- (b) Find the two roots $\lambda_1 \geq \lambda_2$ of $f_A(X)$.
- (c) Find non-zero vectors $\mathbf{v}_1, \mathbf{v}_2$ for which $A\mathbf{v}_j = \lambda_j\mathbf{v}_j$ for $j = 1, 2$. (In general, if $A\mathbf{v} = \lambda\mathbf{v}$ and $\mathbf{v} \neq \mathbf{0}$, we call \mathbf{v} an **eigenvector** of A corresponding to the **eigenvalue** λ .)
- (d) Let P be the matrix whose columns are \mathbf{v}_1 and \mathbf{v}_2 . Compute $P^{-1}AP$.
- (e) *Cayley-Hamilton theorem.* Suppose $f_A(X) = a_0 + a_1X + a_2X^2$. (The values of a_0, a_1, a_2 are known from part (a).) Compute

$$a_0I + a_1A + a_2A^2.$$

1.3 Answers

1. (a) $\begin{pmatrix} 6 \\ 2 \end{pmatrix}$
(b) $\begin{pmatrix} 8 \\ -2 \end{pmatrix}$
(c) Both are 5. In general, $\mathbf{a} \cdot \mathbf{b} = \mathbf{b} \cdot \mathbf{a}$.
(d) $\|\mathbf{u}\| = \sqrt{13}$
 $\|\mathbf{v}\| = \sqrt{17}$
 $\|\mathbf{u} + \mathbf{v}\| = \sqrt{40} = 2\sqrt{10}$
(e) $\arccos\left(\frac{5}{\sqrt{221}}\right)$
(f) $\text{proj}_{\mathbf{v}}(\mathbf{u}) = \begin{pmatrix} 20/17 \\ -5/17 \end{pmatrix}$
 $\text{proj}_{\mathbf{u}}(\mathbf{v}) = \begin{pmatrix} 10/13 \\ 15/13 \end{pmatrix}$
2. (a) $\begin{pmatrix} 18 \\ 7 \end{pmatrix}$
(b) Let $\mathbf{u} = \begin{pmatrix} a \\ b \end{pmatrix}$. Then

$$\mathbf{A}\mathbf{u} = \begin{pmatrix} 2 & 4 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 2a + 4b \\ a + b \end{pmatrix},$$

so we require $2a + 4b = 5$ and $a + b = 2$. The solution to this system is that $a = 3/2$ and $b = 1/2$, so then $\mathbf{u} = \begin{pmatrix} 3/2 \\ 1/2 \end{pmatrix}$.

Remark: We can also compute $\mathbf{u} = \mathbf{A}^{-1}\mathbf{v}$ once we have \mathbf{A}^{-1} (see Problem 6).

3. (a) $\begin{pmatrix} -1 & 8 \\ 6 & -6 \end{pmatrix}$
(b) $\begin{pmatrix} -6 & -12 \\ -3 & -3 \end{pmatrix}$
(c) $\begin{pmatrix} 14 & -20 \\ 2 & -3 \end{pmatrix}$
(d) $\begin{pmatrix} -2 & -8 \\ 3 & 13 \end{pmatrix}$
(e) $\begin{pmatrix} -3 & 5 \\ 4 & -7 \end{pmatrix}$
4. (a) $4\mathbf{I} = \begin{pmatrix} 4 & 0 \\ 0 & 4 \end{pmatrix}$
(b) $\begin{pmatrix} 3 & 0 \\ 0 & 2 \end{pmatrix}$

- (c) $\begin{pmatrix} \cos(\pi/4) & -\sin(\pi/4) \\ \sin(\pi/4) & \cos(\pi/4) \end{pmatrix} = \begin{pmatrix} 1/\sqrt{2} & -1/\sqrt{2} \\ 1/\sqrt{2} & 1/\sqrt{2} \end{pmatrix}$
- (d) $P = \begin{pmatrix} 4/13 & 6/13 \\ 6/13 & 9/13 \end{pmatrix}$
- (e) $2P - I = \begin{pmatrix} -5/13 & 12/13 \\ 12/13 & 5/13 \end{pmatrix}$
5. (a) $\det A = -2$
 $\det B = 1$
- (b) $\det(AB) = \det(A) \cdot \det(B) = -2$
- (c) $\det(A^T) = \det A = -2$
- (d) $\det(A + B) = \det \begin{pmatrix} -1 & 8 \\ 6 & -6 \end{pmatrix} = -42$
- (e) $|\det A| \cdot (\text{unit circle area}) = 2\pi$
6. (a) $A^{-1} = \frac{1}{\det A} \begin{pmatrix} 1 & -4 \\ -1 & 2 \end{pmatrix} = \begin{pmatrix} -1/2 & 2 \\ 1/2 & -1 \end{pmatrix}$
 $B^{-1} = \frac{1}{\det B} \begin{pmatrix} -7 & -4 \\ -5 & -3 \end{pmatrix} = \begin{pmatrix} -7 & -4 \\ -5 & -3 \end{pmatrix}$
- (b) $A^{-1}B^{-1} = \begin{pmatrix} -13/2 & -4 \\ 3/2 & 1 \end{pmatrix}$
 $B^{-1}A^{-1} = \begin{pmatrix} 3/2 & -10 \\ 1 & -7 \end{pmatrix}$
- (c) $(AB)^{-1} = B^{-1}A^{-1} = \begin{pmatrix} 3/2 & -10 \\ 1 & -7 \end{pmatrix}$
- (d) $(A^T)^{-1} = (A^{-1})^T = \begin{pmatrix} -1/2 & 1/2 \\ 2 & -1 \end{pmatrix}$
- (e) $(A + B)^{-1} = \frac{1}{\det(A + B)} \begin{pmatrix} -6 & -8 \\ -6 & -1 \end{pmatrix} = \begin{pmatrix} 1/7 & 4/21 \\ 1/7 & 1/42 \end{pmatrix}$
- (f) $\det(A^{-1}) = 1/\det A = -1/2$
7. (a) A parallelogram with vertices $(0, 0), (1, 0), (3, 1), (2, 1)$
- (b) $\det \begin{pmatrix} 1 & k \\ 0 & 1 \end{pmatrix} = 1$
- (c) Multiplying the right two matrices, $\begin{pmatrix} 4 & 1 \\ 3 & 7 \end{pmatrix} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} a & ak \\ 0 & b \end{pmatrix}$. Looking at the image of vector $\hat{\mathbf{i}}$, we need $\begin{pmatrix} a \\ 0 \end{pmatrix}$ to rotate to $\begin{pmatrix} 4 \\ 3 \end{pmatrix}$. This can be achieved with a rotation by $\theta = \arccos(4/5)$ and $a = 5$. To find b , taking the determinant on both sides and noting that rotations have determinant 1, we require $ab = 25$, so $b = 5$. Finally, to get k , we need $\begin{pmatrix} 5k \\ 5 \end{pmatrix}$ to rotate to $\begin{pmatrix} 1 \\ 7 \end{pmatrix}$. Comparing lengths and noting that $\begin{pmatrix} 5k \\ 5 \end{pmatrix}$ must be in the first quadrant, $k = 1$.

8. (a) $\text{tr } \mathbf{A} = 3$
 $\text{tr } \mathbf{B} = -10$
 $\text{tr}(\mathbf{A} + \mathbf{B}) = \text{tr } \mathbf{A} + \text{tr } \mathbf{B} = -7$
 $\text{tr}(\mathbf{AB}) = 11$

- (b) Let $\mathbf{P} = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ and $\mathbf{Q} = \begin{pmatrix} e & f \\ g & h \end{pmatrix}$. Then

$$\mathbf{PQ} = \begin{pmatrix} ae + bg & af + bh \\ ce + dg & cf + dh \end{pmatrix} \quad \text{and} \quad \mathbf{QP} = \begin{pmatrix} ae + cf & be + df \\ ag + ch & bg + dh \end{pmatrix},$$

so $\text{tr}(\mathbf{PQ}) = \text{tr}(\mathbf{QP}) = ae + bg + cf + dh$.

- (c) In general, the answer is **no**. For example, let

$$\mathbf{A} = \begin{pmatrix} 2 & 4 \\ 1 & 1 \end{pmatrix}, \quad \mathbf{B} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad \mathbf{C} = \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix}.$$

Then

$$\mathbf{ABC} = \begin{pmatrix} 2 & 4 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 2 & 1 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 4 & 6 \\ 2 & 2 \end{pmatrix},$$

$$\mathbf{ACB} = \begin{pmatrix} 2 & 4 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 2 & 2 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 4 & 8 \\ 2 & 3 \end{pmatrix},$$

so $\text{tr}(\mathbf{ABC}) = 6$ while $\text{tr}(\mathbf{ACB}) = 7$.

9. (a) Suppose $\mathbf{I} \sim \mathbf{B}$. Then there is an invertible matrix \mathbf{P} such that $\mathbf{B} = \mathbf{P}^{-1}\mathbf{I}\mathbf{P}$, but the right hand side simplifies to $\mathbf{P}^{-1}\mathbf{P} = \mathbf{I}$.
(b) If $\mathbf{A} \sim \mathbf{B}$ with $\mathbf{B} = \mathbf{P}^{-1}\mathbf{A}\mathbf{P}$, then

$$\det \mathbf{B} = \det(\mathbf{P}^{-1}\mathbf{A}\mathbf{P}) = \det(\mathbf{P})^{-1} \cdot \det \mathbf{A} \cdot \det \mathbf{P} = \det \mathbf{A}.$$

For the trace, Problem 8b gives us

$$\text{tr } \mathbf{B} = \text{tr}(\mathbf{P}^{-1}(\mathbf{A}\mathbf{P})) = \text{tr}((\mathbf{A}\mathbf{P})\mathbf{P}^{-1}) = \text{tr } \mathbf{A}.$$

- (c) We have $\det \mathbf{A} = 4$ and $\text{tr } \mathbf{A} = 5$, so

$$\det \mathbf{D} = d_1 d_2 = 4 \quad \text{and} \quad \text{tr } \mathbf{D} = d_1 + d_2 = 5.$$

This is satisfied by $d_1 = 4$ and $d_2 = 1$, so $\mathbf{D} = \begin{pmatrix} 4 & 0 \\ 0 & 1 \end{pmatrix}$.

10. placeholder