

Fiddler on the Proof - Sorting Hat

Alex Zhu

November 12, 2024

This is a solution to a Fiddler on the Proof [puzzle](#).

To make things easier to follow, let us suppose that Logwarts' four houses are:

- Graphindor (hence referred to as \mathcal{G})
- Riemannclaw (hence referred to as \mathcal{R})
- Hexapuff (hence referred to as \mathcal{H})
- Scalerin (hence referred to as \mathcal{S})

Suppose that for some arbitrary round S , the probability of a student being sorted into \mathcal{G} is P_s .

Now let us work through round $S + 1$.

If the student S was placed into \mathcal{G} , then, by the rules, student $S + 1$ cannot be a \mathcal{G} . So we can disregard this case.

On the other hand, suppose student S was placed into one of the other housees - let's denote that as \mathcal{X} , which may be any of $\mathcal{R}, \mathcal{H}, \mathcal{S}$.

- This student has a $1/4$ chance of choosing \mathcal{G} , which is allowed.
- However, this student also has a $1/4$ chance of choosing \mathcal{X} again, which is not allowed. So that's a $1/3$ chance of the hat placing them in \mathcal{G} against their will, for a total of $1/4 * 1/3 = 1/12$ chance of \mathcal{G} given that the previous student S was not placed into \mathcal{G} .
- The remaining $1/2$ chance is that the student picks one of the other two non- \mathcal{G} and non- \mathcal{X} . This is allowed and doesn't contribute to the odds of student $S + 1$ ending up in \mathcal{G} .

So, if student S is placed into \mathcal{X} , the chances of student $S + 1$ being placed into \mathcal{G} are $1/4 + 1/12 = 1/3$.

We know the odds of student S being placed into \mathcal{X} to be $1 - P_s$, and we can disregard the P_s case as impossible for student $S + 1$ to be a \mathcal{G} , so that means that:

$$P_{s+1} = 1/3(1 - P_s)$$

Let's fruther define B_s as the chances that our protagonist Barry Plotter gets his choice, given that he is S_{th} in line.

Since Barry chooses \mathcal{G} , he will get his wish as long as student $S - 1$ doesn't pick \mathcal{G} . So:

$$B_s = 1 - P_{s-1}$$

We can plug this into Excel (okay, Google Sheets) to get our results:

Round	P_s	B_s
1	1.0000	n/a (Barry can't be first)
2	0.0000	0.0000
3	0.3333	1.0000
4	0.2222	0.6667
5	0.2593	0.7778
6	0.2469	0.7407
7	0.2510	0.7531
8	0.2497	0.7490
9	0.2501	0.7503
10	0.2500	0.7499

So our solution is $B_{10} = 74.9886\%$.