

CLASS TOPIC

Continuous and normal distributions

What this class fixes

- Know the central question of the topic.
- Use the correct notation and units.
- Avoid mechanical calculation without interpretation.

W05**Statistics**

Clean formulas · aligned layout ·
corrected terminology



CLASS TOPIC

Continuous and normal distributions


Economic question first

- Start from a real decision: compare, describe, forecast or test.
- Translate the question into variables before using formulas.
- Define the population, sample and measurement unit.

FORMULA

$$P(a \leq X \leq b) = \int_a^b f(x) dx$$

Total area under $f(x)$ equals 1



CLASS TOPIC

Continuous and normal distributions


Key definitions

- Population: full group of interest.
- Sample: observed part of the population.
- Statistic: number computed from a sample;
parameter: number for the population.

FORMULA

$$P(a \leq X \leq b) = \int_a^b f(x) dx$$

Total area under $f(x)$ equals 1



CLASS TOPIC

Continuous and normal distributions


Main formula box

- All formulas are kept in one merged formula area on this slide.
- Read every symbol before calculating.
- Use parentheses carefully.

FORMULA 1

$$P(a \leq X \leq b) = \int_a^b f(x) dx$$

Total area under $f(x)$ equals 1



CLASS TOPIC

Continuous and normal distributions

Mini example

- Write the data clearly.
- Substitute values step by step.
- Round only at the final reporting stage.

01 Data: 4, 7, 8, 9, 12

02 Step 1: compute the needed statistic.

03 Step 2: write the result with units.

04 Step 3: explain it in one sentence.

FORMULA

$$P(a \leq X \leq b) = \int_a^b f(x) dx$$


Total area under $f(x)$ equals 1

CLASS TOPIC

Continuous and normal distributions

Visual intuition

- A graph should answer a question, not decorate the slide.
- Always label axes and units.
- Look for center, spread, shape and unusual values.

**FORMULA**

$$P(a \leq X \leq b) = \int_a^b f(x) dx$$

CLASS TOPIC

Continuous and normal distributions

Second formula box


- This formula is separated from explanations but kept in one clean box.
- No broken equation fragments or floating text boxes.
- Notation is consistent across English and Russian versions.

FORMULA 2

$$Z = (X - \mu) / \sigma$$

$$X \sim N(\mu, \sigma^2)$$

$$P(X \leq x) = P(Z \leq (x - \mu) / \sigma)$$



CLASS TOPIC

Continuous and normal distributions


Common mistakes

- Mixing sample and population symbols.
- Using a formula without checking conditions.
- Reporting a number without units or interpretation.

FORMULA

$$P(a \leq X \leq b) = \int_a^b f(x) dx$$

Total area under $f(x)$ equals 1



CLASS TOPIC

Continuous and normal distributions

R / spreadsheet check

- Use software to calculate, but keep the statistical logic visible.
- Check variable names, missing values and units first.
- Reproducible work is easier to grade and correct.

R / EXCEL

```
# example pattern  
mean(x); sd(x); summary(x)  
plot(x, y); cor(x, y)
```

FORMULA

$$P(a \leq X \leq b) = \int_a^b f(x) dx$$


Total area under $f(x)$ equals 1

CLASS TOPIC

Continuous and normal distributions

Interpretation sentence

- Do not stop at the number.
- Explain what the result means in the original context.
- Use cautious language when evidence is limited.

**FORMULA**

$$P(a \leq X \leq b) = \int_a^b f(x) dx$$

CLASS TOPIC

Continuous and normal distributions


Conditions and assumptions

- Check whether the method fits the data type.
- Check independence and sample size when needed.
- If assumptions fail, explain the limitation.

FORMULA

$$P(a \leq X \leq b) = \int_a^b f(x) dx$$

Total area under $f(x)$ equals 1



CLASS TOPIC

Continuous and normal distributions


Class exercise

- Solve one calculation by hand.
- Then verify it with software.
- Write a two-sentence conclusion.

FORMULA

$$P(a \leq X \leq b) = \int_a^b f(x) dx$$

Total area under $f(x)$ equals 1




CLASS TOPIC

Continuous and normal distributions

How to report

- Report method, estimate, uncertainty and interpretation.
- Use tables only when they make comparison easier.
- Separate statistical evidence from causal claims.

**FORMULA**

$$P(a \leq X \leq b) = \int_a^b f(x) dx$$

CLASS TOPIC

Continuous and normal distributions

Final checklist

- Question is clear.
- Variables and units are defined.
- Formula is correct.
- Result is interpreted.
- Limitations are stated.

FORMULA

$$P(a \leq X \leq b) = \int_a^b f(x) dx$$

Total area under $f(x)$ equals 1

