



# Paradoxical sets and the Axiom of Choice

Azul Lihuen Fatalini


Joint work with Prof. Ralf Schindler

Axiom  
of Choice



Paradoxical  
sets

# My research



# Set Theory

Sets

Consistency

Cardinality

ZFC

$\emptyset, \cap, \cup, \subseteq$

Axiom of Choice

Paradoxes

Intersectionality

Models

Venn diagrams

Continuum  
hypothesis

Infinity

Zorn's lemma


Induction

Forcing

$\aleph_0$


# Set Theory

•  $\mathfrak{c} = |\mathbb{R}|$   
↑  
"continuum"



# Set Theory


•  $\mathfrak{c} = |\mathbb{R}|$   
↑  
"continuum"  $\rightsquigarrow$  Continuum hypothesis



# Set Theory


- $\mathfrak{c} = |\mathbb{R}|$

↑  
"continuum"  $\rightsquigarrow$  Continuum hypothesis



# Set Theory


- Model of  $ZF(c)$  = "universe"







# Set Theory

- Model of  $ZF(C)$  = "universe"  
↓  
Axiom of Choice





# Set Theory

- Model of  $ZF(C)$  = "universe"  $\rightarrow$  There are several  
Axiom of Choice




There are  
several




# Choice principles

Axiom of Choice



Zorn's lemma



Well-ordering principle



" $\mathbb{R}$  can be well-ordered"

Axiom  
of Choice



Paradoxical  
sets

# Paradoxical sets

- $p \subseteq \mathbb{R}^n \quad n = 1, 2, 3.$

[  $p$  satisfies some counterintuitive property.

[ The existence of such a  $p$  involves Axiom of Choice

# Paradoxical sets

- $p \subseteq \mathbb{R}^n$      $n = 1, 2, 3$ .

[  $p$  satisfies some counterintuitive property.

[ The existence of such a  $p$  involves Axiom of Choice



a well-order  
of the reals



allows induction  
"on the reals"

# Set Theory

Sets

Consistency

Cardinality

ZFC

$\emptyset, \cap, \cup, \subseteq$

Axiom of Choice

Paradoxes

Intersectionality

Models

Venn diagrams

Continuum  
hypothesis

Infinity

Zorn's lemma


Induction

Forcing

$\aleph_0$




# My research



# Contribution

Negative answer  $\rightarrow$  Model of  $ZF + \neg C + \exists P$






# Theorem (ZFC) (Conway-Croft / Kharazishvili)


$\mathbb{R}^3$  can be partitioned in unit circles.

Let  $\mathbb{R}^3 = \{P_\alpha\}_{\alpha \in \mathbb{C}}$ .


(0)



(1)



(2)



## Theorem

There is a model of

$ZF + \exists$  partition of  $\mathbb{R}^3$  in unit circles  
+ no well order of the reals.


# Set Theory

How does the proof look like?



$$(M, \mathbb{P}, g \in \mathbb{P}) \longrightarrow \boxed{\text{Forcing}} \rightsquigarrow M[g]$$

# Strategy

Negative answer  $\rightarrow$  Model of  $ZF + \neg C + \exists P$






- Strategy:
- Construct a model (of  $ZF + \neg C$ ) which contains a nice structure of inner models satisfying AC.
  - $P$  will be a limit of the partial paradoxical sets.



$M[g]$   
 $\downarrow$   
 $M'$







# Contribution

	Hamel basis	Mazurkiewicz	Partition of $\mathbb{R}^3$ in unit circles
$DC + \neg WO(\mathbb{R})$	✓ [4]	✓ [4]	✓
$DC + \neg U_f(\omega)$	✓	✓	?
$\neg AC_\omega$	✓ [3]	✓ [2]	✓

# References

- [1] P. Bankston and R. Fox, Topological partitions of euclidean space by spheres, *The American Mathematical Monthly*, 92 (1985), p. 423.
- [2] M. Beriashvili and R. Schindler, Mazurkiewicz sets with no well-ordering of the reals, *Georgian Mathematical Journal*. 2022;29(3): 343-345. <https://doi.org/10.1515/gmj-2022-2142>
- [3] M. Beriashvili, R. Schindler, L. Wu, and L. Yu, Hamel bases and well-ordering the continuum, in *Proc. Amer. Math. Soc* (Vol. 146, pp. 3565-3573), (2018).
- [4] J. Brendle, F. Castiblanco, R. Schindler, L. Wu, and L. Yu, A model with everything except for a well-ordering of the reals, (2018).
- [5] J. H. Conway and H. T. Croft, Covering a sphere with congruent great-circle arcs, *Mathematical Proceedings of the Cambridge Philosophical Society*, 60 (1964), pp. 787-800.
- [6] M. Jonsson and J. Wästlund, Partitions of  $\mathbb{R}^3$  into curves, *Mathematica Scandinavica*, 83. 10.7146/math.scand.a-13850 , (1998).
- [7] A. B. Kharazishvili, Partition of the three-dimensional space into congruent circles, *Bull. Acad. Sci. GSSR*, v. 119, n. 1, 1985 (in Russian).
- [8] A. B. Kharazishvili and T. S. Tetunashvili, On some coverings of the euclidean plane with pairwise congruent circles, *American Mathematical Monthly*, 117 (2010), pp. 414-423.
- [9] J. Schilhan, Maximal sets without choice, (2022).
- [10] A. Szulkin,  $\mathbb{R}^3$  is the union of disjoint circles, (1983).
- [11] Z. Vidnyánszky, Transfinite inductions producing coanalytic sets, *Fundamenta Mathematicae*. 224. 10.4064/fm224-2-2 , (2012)

Theorem (ZFC) (Conway-Croft / Kharazishvili)

$\mathbb{R}^3$  can be partitioned in unit circles.

Theorem (ZFC) (Conway-Croft / Kharazishvili)

$\mathbb{R}^3$  can be partitioned in unit circles.


Let  $\mathbb{R}^3 = \{P_\alpha\}_{\alpha < \mathfrak{c}}$ .

Theorem (ZFC) (Conway-Croft / Kharazishvili)

$\mathbb{R}^3$  can be partitioned in unit circles.

Let  $\mathbb{R}^3 = \{P_\alpha\}_{\alpha \in \mathbb{C}}$ .

(0)




Theorem (ZFC) (Conway-Croft / Kharazishvili)


$\mathbb{R}^3$  can be partitioned in unit circles.

Let  $\mathbb{R}^3 = \{P_\alpha\}_{\alpha \in \mathbb{C}}$ .

(0)



(1)






# Theorem (ZFC) (Conway-Croft / Kharazishvili)


$\mathbb{R}^3$  can be partitioned in unit circles.

Let  $\mathbb{R}^3 = \{P_\alpha\}_{\alpha \in \mathbb{C}}$ .


(0)



(1)





(2)




## Theorem (ZF) (Szulkin)

$\mathbb{R}^3$  can be partitioned in circles.



# Algebra



## Theorem (F., Schindler; ~2024)

Let  $V$  be a model of ZFC and let  $S$  be a finite set of mutually generic Cohen reals over  $V$ .

Then the transcendence degree of  $\mathbb{R}^{V[S]}$  over  $\bigcup_{\substack{T \subseteq S \\ |T|=|S|-1}} \mathbb{R}^{V[T]}$  is  $\mathbb{C}$ .