

INDICATION: , Iron deficiency anemia.,PROCEDURE:  
,Colonoscopy with terminal ileum  
examination.,POSTOPERATIVE DIAGNOSIS:, Normal  
examination.,WITHDRAWAL TIME: , 15 minutes.,SCOPE: ,  
CF-H180AL.,MEDICATIONS: , Fentanyl 100 mcg and versed  
10 mg.,PROCEDURE DETAIL: ,Following the preprocedure  
patient assessment the procedure, goals, risks including  
bleeding, perforation, missed polyp rate as well as side effects  
of medications and alternatives were reviewed. Questions  
were answered. Pause preprocedure was  
performed.,Following titrated intravenous sedation the flexible  
video endoscope was introduced into the rectum and  
advanced to the cecum without difficulty. The ileocecal valve  
looked normal. Preparation was fair allowing examination of  
85% of mucosa after washing and cleaning with tap water  
through the scope. The terminal ileum was intubated through  
the ileocecal valve for a 5 cm extent. Terminal ileum mucosa  
looked normal.,Then the scope was withdrawn while  
examining the mucosa carefully including the retroflexed  
views of the rectum. No polyp, no diverticulum and no  
bleeding source was identified.,The patient was assessed  
upon completion of the procedure. Okay to discharge once  
criteria met. ,RECOMMENDATIONS:, Follow up with primary  
care physician.