

PREOPERATIVE DIAGNOSES:, Increased intracranial pressure and cerebral edema due to severe brain injury.,POSTOPERATIVE DIAGNOSES: , Increased intracranial pressure and cerebral edema due to severe brain injury.,PROCEDURE:, Burr hole and insertion of external ventricular drain catheter.,ANESTHESIA: , Just bedside sedation.,PROCEDURE: , Scalp was clipped. He was prepped with ChloroPrep and Betadine. Incisions are infiltrated with 1% Xylocaine with epinephrine 1:200000. He did receive antibiotics post procedure. He was draped in a sterile manner.,Incision made just to the right of the right mid pupillary line 10 cm behind the nasion. A self-retaining retractor was placed. Burr hole was drilled with the cranial twist drill. The dura was punctured with a twist drill. A brain needle was used to localize the ventricle that took 3 passes to localize the ventricle. The pressure was initially high. The CSF was clear and colorless. The CSF drainage rapidly tapered off because of the brain swelling. With two tries, the ventricular catheter was then able to be placed into the ventricle and then brought out through a separate stab wound, the depth of catheter is 7 cm from the outer table of the skull. There was intermittent drainage of CSF after that. The catheter was secured to the scalp with #2-0 silk suture and the incision was closed with Ethilon suture. The patient tolerated the procedure well. No complications. Sponge and needle counts were correct. Blood loss is minimal. None replaced.