

PREOPERATIVE DIAGNOSES: ,1. Nasolabial mesiolabial fold.,2. Mid glabellar fold.,POSTOPERATIVE DIAGNOSES: ,1. Nasolabial mesiolabial fold.,2. Mid glabellar fold.,TITLE OF PROCEDURES: ,1. Perlane injection for the nasolabial fold.,2. Restylane injection for the glabellar fold.,ANESTHESIA: ,Topical with Laseraine.,COMPLICATIONS: , None.,PROCEDURE: , The patient was evaluated preop and noted to be in stable condition. Chart and informed consent were all reviewed preop. All risks, benefits, and alternatives regarding the procedure have been reviewed in detail with the patient. This includes risk of bleeding, infection, scarring, need for further procedure, etc. The patient did sign the informed consent form regarding the Perlane and Restylane. She is aware of the potential risk of bruising. The patient has had Cosmederm in the past and had had a minimal response with this. Please note Laseraine had to be applied 30 minutes prior to the procedure. The excess Laseraine was removed with a sterile alcohol swab.,Using the linear threading technique, I injected the deep nasolabial fold. We used 2 mL of the Perlane for injection of the nasolabial mesiolabial fold. They were carefully massaged into good position at the end of the procedure. She did have some mild erythema noted.,I then used approximately 0.4 mL of the Restylane for injection of the mid glabellar site. She has a resting line of the mid glabella that did not respond with previous Botox injection. Once this was filled, the Restylane was massaged into the proper tissue plane. Cold compressors were applied

afterwards. She is scheduled for a recheck in the next one to two weeks, and we will make further recommendations at that time. Post Restylane and Perlane precautions have been reviewed with the patient as well.