

PREOPERATIVE DIAGNOSIS:, Visually significant posterior capsule opacity, right eye.,POSTOPERATIVE DIAGNOSIS:, Visually significant posterior capsule opacity, right eye.,OPERATIVE PROCEDURES: ,YAG laser posterior capsulotomy, right eye.,ANESTHESIA: , Topical anesthesia using tetracaine ophthalmic drops.,INDICATIONS FOR SURGERY: , This patient was found to have a visually significant posterior capsule opacity in the right eye. The patient has had a mild decrease in visual acuity, which has been a gradual change. The posterior capsule opacity was felt to be related to the decline in vision. The risks, benefits, and alternatives (including observation) were discussed. I feel the patient had a good understanding of the proposed procedure and informed consent was obtained.,DESCRIPTION OF PROCEDURE: , The patient was identified and the procedure was verified. Pupil was dilated per protocol. Patient was positioned at the YAG laser. Then, \*\*\* of energy were used to perform a circular posterior laser capsulotomy through the visual axis. A total of \*\* shots were used. Total energy was \*\*. The patient tolerated the procedure well and there were no complications. The lens remained well centered and stable. Postoperative instructions were provided. Alphagan P ophthalmic drops times two were instilled prior to his dismissal.,Post-laser intraocular pressure measured \*\* mmHg. Postoperative instructions were provided and the patient had no further questions.