

INDICATION: , Iron deficiency anemia.,PROCEDURE:
,Colonoscopy with terminal ileum
examination.,POSTOPERATIVE DIAGNOSIS:, Normal
examination.,WITHDRAWAL TIME: , 15 minutes.,SCOPE: ,
CF-H180AL.,MEDICATIONS: , Fentanyl 100 mcg and versed
10 mg.,PROCEDURE DETAIL: ,Following the preprocedure
patient assessment the procedure, goals, risks including
bleeding, perforation, missed polyp rate as well as side effects
of medications and alternatives were reviewed. Questions
were answered. Pause preprocedure was
performed.,Following titrated intravenous sedation the flexible
video endoscope was introduced into the rectum and
advanced to the cecum without difficulty. The ileocecal valve
looked normal. Preparation was fair allowing examination of
85% of mucosa after washing and cleaning with tap water
through the scope. The terminal ileum was intubated through
the ileocecal valve for a 5 cm extent. Terminal ileum mucosa
looked normal.,Then the scope was withdrawn while
examining the mucosa carefully including the retroflexed
views of the rectum. No polyp, no diverticulum and no
bleeding source was identified.,The patient was assessed
upon completion of the procedure. Okay to discharge once
criteria met. ,RECOMMENDATIONS:, Follow up with primary
care physician.