

PREOPERATIVE DIAGNOSIS: ,  
Phimosis.,POSTOPERATIVE DIAGNOSIS: ,  
Phimosis.,PROCEDURE: , Reduction of  
paraphimosis.,ANESTHESIA: ,General inhalation anesthetic  
with 0.25% Marcaine, penile block and ring block about 20 mL  
given.,FLUIDS RECEIVED: , 100 mL.,SPECIMENS:, No  
tissues sent to pathology.,COUNTS: , Sponge and needle  
counts were not necessary.,TUBES/DRAINS: , No tubes or  
drains were used.,FINDINGS: , Paraphimosis with moderate  
swelling.,INDICATIONS FOR OPERATION: , The patient is a  
15-year-old boy who had acute alcohol intoxication had his  
foreskin retracted with a Foley catheter placed at another  
institution. When they removed the catheter they forgot to  
reduce the foreskin and he developed paraphimosis. The plan  
is for reduction.,DESCRIPTION OF OPERATION: , The  
patient was taken to the operating room where surgical  
consent, operative site, and patient identification were  
verified. Once he was anesthetized, with manual pressure  
and mobilization of the shaft skin we were able to reduce the  
paraphimosis. Using Betadine and alcohol cleanse, we then  
did a dorsal penile block and a ring block by surgeon with  
0.25% Marcaine, 20 mL were given. He did quite well after the  
procedure and was transferred to the recovery room in stable  
condition.