

PROCEDURE:, Colonoscopy.,PREOPERATIVE DIAGNOSIS:  
, Follow up adenomas.,POSTOPERATIVE DIAGNOSES:,1.  
Two colon polyps, removed.,2. Small internal hemorrhoids.,3.  
Otherwise normal examination of cecum.,MEDICATIONS: ,  
Fentanyl 150 mcg and Versed 7 mg slow IV  
push.,INDICATIONS: , This is a 60-year-old white female with  
a history of adenomas. She does have irregular bowel  
habits.,FINDINGS: , The patient was placed in the left lateral  
decubitus position and the above medications were  
administered. The colonoscope was advanced to the cecum  
as identified by the ileocecal valve, appendiceal orifice, and  
blind pouch. The colonoscope was slowly withdrawn and a  
careful examination of the colonic mucosa was made,  
including a retroflexed view of the rectum. There was a 4 mm  
descending colon polyp, which was removed with jumbo  
forceps, and sent for histology in bottle one. There was a 10  
mm pale, flat polyp in the distal rectum, which was removed  
with jumbo forceps, and sent for histology in bottle 2. There  
were small internal hemorrhoids. The remainder of the  
examination was normal to the cecum. The patient tolerated  
the procedure well without complication.,IMPRESSION:,1.  
Two colon polyps, removed.,2. Small internal hemorrhoids.,3.  
Otherwise normal examination to cecum.,PLAN: , I will await  
the results of the colon polyp histology. The patient was told  
the importance of daily fiber.