

PREOPERATIVE DIAGNOSIS: ,
Phimosis.,POSTOPERATIVE DIAGNOSIS: ,
Phimosis.,PROCEDURE: , Reduction of
paraphimosis.,ANESTHESIA: ,General inhalation anesthetic
with 0.25% Marcaine, penile block and ring block about 20 mL
given.,FLUIDS RECEIVED: , 100 mL.,SPECIMENS:, No
tissues sent to pathology.,COUNTS: , Sponge and needle
counts were not necessary.,TUBES/DRAINS: , No tubes or
drains were used.,FINDINGS: , Paraphimosis with moderate
swelling.,INDICATIONS FOR OPERATION: , The patient is a
15-year-old boy who had acute alcohol intoxication had his
foreskin retracted with a Foley catheter placed at another
institution. When they removed the catheter they forgot to
reduce the foreskin and he developed paraphimosis. The plan
is for reduction.,DESCRIPTION OF OPERATION: , The
patient was taken to the operating room where surgical
consent, operative site, and patient identification were
verified. Once he was anesthetized, with manual pressure
and mobilization of the shaft skin we were able to reduce the
paraphimosis. Using Betadine and alcohol cleanse, we then
did a dorsal penile block and a ring block by surgeon with
0.25% Marcaine, 20 mL were given. He did quite well after the
procedure and was transferred to the recovery room in stable
condition.