
a
Six full annotated RNA readouts
ER folding (9)
ERAD (67)
ER targeting (34)
ISR (20)
Reactome UPR (69)
GO ER-UPR (41)
0.28* 0.72* 0.05 0.41 0.20
0.22* 0.72* 0.55* -0.07 0.20
0.11 0.79* 0.25 -0.24 0.12
0.24* 0.11 -0.38 -0.02 0.16
0.17* 0.84* 0.49 -0.00 0.59*
0.18* 0.84* 0.29 -0.05 0.72*
RPE1 438 targets
Adamson 30 targets
Melanoma 19 targets
IFNg 19 targets
Co-culture 17 targets
Spearman rho
-1 0 1
*BH q < 0.05 across all 80 readout/context tests.
b
Matched random-readout calibration
ER folding (9)
ERAD (67)
ER targeting (34)
ISR (20)
Reactome UPR (69)
GO ER-UPR (41)
Reactome disjoint (30)
GO disjoint (32)
0.47 0.35 0.50 0.06 0.25
0.86 0.34 0.08 0.92 0.49
0.99 0.07 0.32 1.00 0.42
0.27 0.99 0.99 0.68 0.52
0.92 0.02 0.22 0.82 0.06
0.84 0.02 0.51 0.77 0.01
1.00 0.37 0.26 0.97 0.24
0.38 0.55 0.36 0.92 0.02
RPE1 438 targets
Adamson 30 targets
Melanoma 19 targets
IFNg 19 targets
Co-culture 17 targets
Upper-tail fraction
0.0 0.5 1.0
100 size/expression/detection-matched nulls per readout.
Real and null scores share fixed-library normalization.
c
Parent UPR definitions: full and Hallmark-disjoint
RPE1 438 targets
Adamson 30 targets
Melanoma 19 targets
IFNg 19 targets
Co-culture 17 targets
Reactome UPR (69)
Reactome disjoint (30)
GO ER-UPR (41)
GO disjoint (32)
-1 0 1
Source–context Spearman correlation
95% whole-gene bootstrap intervals; all outcomes were previously examined.